



**INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)**

<p>(51) International Patent Classification <sup>5</sup> :  <b>C07K 5/06, 5/02, A61K 38/55, C07C 257/18, 257/16, C07D 239/14, 211/26, 205/04, 207/09</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 94/29336</b></p> <p>(43) International Publication Date: <b>22 December 1994 (22.12.94)</b></p>
<p>(21) International Application Number: <b>PCT/SE94/00535</b></p> <p>(22) International Filing Date: <b>2 June 1994 (02.06.94)</b></p> <p>(30) Priority Data:  <b>9301916-4      3 June 1993 (03.06.93)      SE</b></p>	<p>(81) Designated States: <b>AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b></p>	
<p>(71) Applicant (for all designated States except US): <b>ASTRA AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE).</b></p> <p>(72) Inventors; and  (75) Inventors/Applicants (for US only): <b>ANTONSSON, Karl, Thomas [SE/SE]; Torkels väg, P1 3503, S-437 34 Lindome (SE). BYLUND, Ruth, Elvy [SE/SE]; Forellgatan 60, S-426 58 Västra Frölunda (SE). GUSTAFSSON, Nils, David [SE/SE]; Ånghagavägen, P1 3758, S-430 41 Kullavik (SE). NILSSON, Nils, Olov, Ingemar [SE/SE]; Rågvägen 50, S-430 33 Fjärås (SE).</b></p> <p>(74) Agent: <b>SAMUELSSON, Britta; Astra Aktiebolag, Patent Dept., S-151 85 Södertälje (SE).</b></p>	<p><b>Published</b>  <i>With international search report.</i>  <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p><b>(54) Title: NEW PEPTIDE DERIVATIVES</b></p>		
<p><b>(57) Abstract</b></p> <p>The invention relates to new competitive inhibitors of trypsin-like serine proteases, their synthesis, pharmaceutical compositions containing the compounds as active ingredients, and the use of the compounds as thrombin inhibitors, anticoagulants and anti-inflammatory inhibitors for prophylaxis and treatment of related diseases, according to the formulas (I): A<sup>1</sup>-A<sup>2</sup>-NH-(CH<sub>2</sub>)<sub>n</sub>-B and (V): A<sup>1</sup>-A<sup>2</sup>-NH-(CH<sub>2</sub>)<sub>n</sub>-B-D wherein A<sup>1</sup> represents a structural fragment of formulas (IIa), (IIb), (IIc), (IIe), A<sup>2</sup> represents a structural fragment of formulas (IIIa), (IIIb), (IIIc), B represents a structural fragment of formulas (IVa), (IVb), (IVc), (IVd). Further described are novel compounds, the new use of compounds and especially new structural fragment in synthesis of pharmaceutical compounds.</p>		
<p>The figure displays twelve chemical structures labeled (IIa) through (IVd). Structures (IIa), (IIb), and (IIc) are linear peptide fragments with side chains R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> respectively. Structure (IId) is a cyclic peptide derivative. Structures (IIIa), (IIIb), and (IIIc) are cyclic peptide derivatives with side chains R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> respectively. Structures (IVa), (IVb), (IVc), and (IVd) are structural fragments, including a cyclic amine, a benzene ring, a pyridine ring, and a pyrazole ring, respectively.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**New peptide derivatives**

5 This invention relates to new competitive inhibitors of  
trypsin-like serine proteases, especially thrombin and  
kininogenases such as kallikrein, their synthesis,  
pharmaceutical compositions containing the compounds as  
active ingredients, and the use of the compounds as  
10 thrombin inhibitors and anticoagulants and as  
antiinflammatory inhibitors, respectively.  
The invention also relates to novel use of compounds as  
starting materials in synthesis of a serine protease  
inhibitor. Furthermore the invention relates to a novel  
structural fragments in serine protease inhibitors.

15

**BACKGROUND**

Blood coagulation is the key process involved in both  
haemostasis (i.e. prevention of blood loss from a  
20 damaged vessel) and thrombosis (i.e. the pathological  
occlusion of a blood vessel by a blood clot).  
Coagulation is the result of a complex series of  
enzymatic reactions, where one of the final steps is  
conversion of the proenzyme prothrombin to the active  
25 enzyme thrombin.

Thrombin plays a central role in coagulation. It  
activates platelets, it converts fibrinogen into fibrin  
monomers, which polymerise spontaneously into  
30 filaments, and it activates factor XIII, which in turn  
crosslinks the polymer to insoluble fibrin. Thrombin  
further activates factor V and factor VIII in a  
positive feedback reaction. Inhibitors of thrombin are  
therefore expected to be effective anticoagulants by  
35 inhibition of platelets, fibrin formation and fibrin  
stabilization. By inhibiting the positive feedback

mechanism they are expected to exert inhibition early in the chain of events leading to coagulation and thrombosis.

5 Kininogenases are serine proteases that act on kininogens to produce kinins (bradykinin, kallidin, and Met-Lys-bradykinin). Plasma kallikrein, tissue kallikrein, and mast cell tryptase represent important kininogenases.

10

Kinins (bradykinin, kallidin) are generally involved in inflammation. For example, the active inflammation process is associated with increased permeability of the blood vessels resulting in extravasation of plasma into the tissue. The ensuing plasma exudate contains all the protein systems of circulating blood. The plasma-derived kininogens inevitably will be interacting with different kallikreins, forming kinins continually as long as the active plasma exudation process is ongoing. Plasma exudation occurs independent of the mechanisms that are involved in the inflammation, whether it is allergy, infection or other factors (Persson et al., Editorial, Thorax, 1992, 47:993-1000). Plasma exudation is thus a feature of many diseases including asthma, rhinitis, common cold, and inflammatory bowel diseases. Particularly in allergy mast cell tryptase will be released (Salomonsson et al., Am. Rev. Respir. Dis., 1992, 146:1535-1542) to contribute to kinin formation and other pathogenic events in asthma, rhinitis, and intestinal diseases.

15  
20  
25  
30

The kinins are biologically highly active substances with smooth muscle effects, secretory effects, neurogenic effects, and actions that may perpetuate inflammatory processes including activation of phospholipase A<sub>2</sub> and increasing vascular permeability. The latter action potentially induces a vicious circle

35

with kinins providing for the generation of more kinins etc.

Tissue kallikrein cleaves primarily low molecular weight kininogen to produce kallidin and plasma kallikrein preferably releases bradykinin from high molecular weight kininogen.

#### PRIOR ART

10

Inhibitors of thrombin based on the amino acid sequence around the cleavage site for the fibrinogen A $\alpha$  chain were first reported by Blombäck et al. in J. Clin. Lab. Invest. 24, suppl 107, 59, (1969), who suggested the sequence Phe-Val-Arg (P9-P2-P1, herein referred to as the P3-P2-P1 sequence) to be the best inhibitor.

15

20

In US 4,346,078 has S. Bajusz et al. described the thrombin inhibitor H-DPhe-Pro-Agm, a dipeptidyl derivative with an aminoalkyl guanidine in the P1-position.

25

Inhibitors of thrombin based on peptide derivatives with a cyclic aminoalkyl guanidine, e.g. 3-aminomethyl-1-amidinopiperidine, in the P1-position have been disclosed in EP-A2-0,468,231.

30

In EP-A2-0,185,390 has S. Bajusz et. al. disclosed that replacing the agmatine with an arginine aldehyde gave a thrombin inhibitor which had much higher potency.

35

Inhibitors of kallikrein based on the amino acid sequence around the cleavage site Arg-Ser have been reported earlier.

The arginine ch1 romethyl k ton s H-DPr -Phe-Arg-CH<sub>2</sub>Cl

and H-D Phe-Phe-Arg-CH<sub>2</sub>Cl were reported as plasma kallikrein inhibitors by Kettner and Shaw in Bi chemistry 1978, 17:4778-4784 and Meth. Enzym. 1981, 80:826-842.

5

Likewise, esters and amides containing the H-DPro-Phe-Arg sequence were reported by Fareed et al. in Ann. N.Y. Acad. Sci. 1981, 370:765-784 to be plasma kallikrein inhibitors.

10

Inhibitors of serine proteases that are based on electrophilic ketones instead of aldehydes in the P1-position are described in the following patent documents:

15

EP-A2-0,195,212 describing peptidyl  $\alpha$ -keto esters and amides, EP-A1-0,362,002 describing fluoroalkylamide ketones and EP-A2-0,364,344 describing  $\alpha,\beta,\delta$ -triketo compounds possessing different peptidase inhibiting properties.

20

Inhibitors of trypsin-like serine proteases, such as thrombin and kallikrein, based on C-terminal boronic acid derivatives of arginine and isothiuronium analogues thereof have been revealed in EP-A2-0,293,881.

25

WO 92/04371 describing kininogenase inhibitors, e.g. kallikrein inhibitors based on derivatives of arginine.

30

EP-A1-0,530,167 describing  $\alpha$ -alkoxy ketone derivatives of arginine as thrombin inhibitors.

35


#### DISCLOSURE OF THE INVENTION

An object of the present invention is to provide novel

and potent trypsin-like serine protease inhibitors, especially antithrombotic and antiinflammatory compounds with competitive inhibitory activity towards their enzyme i.e. causing reversible inhibition. More specifically anticoagulants for prophylaxis and treatment of thromboembolic diseases such as venous thrombosis, pulmonary embolism, arterial thrombosis, in particular myocardial infarction and cerebral thrombosis, general hypercoagulable states and local hypercoagulable states, e.g. following angioplasty and coronary bypass operations, and other situations where thrombin is believed to play a role, e.g. Alzheimer's disease, as well as inhibition of kininogenases for treatment of inflammatory disorders e.g. asthma, rhinitis, urticaria, inflammatory bowel disease, and arthritis. A further object is to obtain thrombin inhibitors which are orally bioavailable and selective in inhibiting thrombin over other serine proteases. A further object of the invention is to obtain kininogenase inhibitors which can be given orally, rectally, topically e.g. dermally, or via the inhalation route.

#### Compounds

According to the invention it has been found that compounds of the general Formula I, either as such or in the form of physiologically acceptable salts, and including stereoisomers, are potent inhibitors of serine proteases, especially thrombin and kininogenases such as kallikrein:




Formula I

wherein:


A<sup>1</sup> represents a structural fragment of Formula  
IIa, IIb, IIc, IIId or IIe;

5


10



IIa




IIb




IIc

15



IIId

20



IIe

25

wherein:

k is an integer 0, 1, 2, 3 or 4;

m is an integer 1, 2, 3 or 4;

30

q is an integer 0, 1, 2 or 3;

35

R<sup>1</sup> represents H, an alkyl group having 1 to 4 carbon atoms, or R<sup>11</sup>OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group R<sup>17</sup>-(CH<sub>2</sub>)<sub>p</sub>-, where p is



0,1 or 2 and  $R^{17}$  is methyl, phenyl, OH,  $\text{COOR}^{12}$ ,  $\text{CONHR}^{12}$ , where  $R^{12}$  is H or an alkyl group having 1 to 4 carbon atoms, and  $R^{11}$  is H or an alkyl group having 1 to 6 carbon atoms, or

5

$R^1$  represents  $\text{Ph}(4\text{-COOR}^{12})\text{-CH}_2\text{-}$ , where  $R^{12}$  is as defined above, or

10

$R^1$  represents  $R^{13}\text{-NH-CO-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{13}$  is H or an alkyl group having 1 to 4 carbon atoms or  $\text{-CH}_2\text{COOR}^{12}$ , where  $R^{12}$  is as defined above, or

15

$R^1$  represents  $R^{12}\text{OOC-CH}_2\text{-OOC-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{12}$  is as defined above, or

20

$R^1$  represents  $R^{14}\text{SO}_2\text{-}$ ,  $\text{Ph}(4\text{-COOR}^{12})\text{-SO}_2\text{-}$ ,  $\text{Ph}(3\text{-COOR}^{12})\text{-SO}_2\text{-}$ ,  $\text{Ph}(2\text{-COOR}^{12})\text{-SO}_2\text{-}$ , where  $R^{12}$  is as defined above and  $R^{14}$  is an alkyl group having 1-4 carbon atoms, or

25

$R^1$  represents  $\text{-CO-R}^{15}$ , wherein  $R^{15}$  is an alkyl group having 1-4 carbon atoms, or

30

$R^1$  represents  $\text{-CO-OR}^{15}$ , where  $R^{15}$  is as defined above, or

$R^1$  represent  $\text{-CO-(CH}_2)_p\text{-COOR}^{12}$ , where  $R^{12}$  is as defined above and  $p$  is an integer 0, 1 or 2, or

35

$R^1$  represents  $\text{-CH}_2\text{PO(OR}^{16})_2$ ,  $\text{-CH}_2\text{SO}_3\text{H}$  or  $\text{-CH}_2\text{-(5-(1H)-tetrazyl)}$ , where  $R^{16}$  is, individually at each occurrence, H, methyl or ethyl;

$R^2$  represents H or an alkyl group having 1 to 4 carbon atoms or  $R^{21}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and, where  $R^{21}$  is H or an alkyl group having 1 to 4 carbon atoms;

5

$R^3$  represents an alkyl group having 1-4 carbon atoms, and the alkyl group may or may not carry one or more fluorine atoms, or

10  $R^3$  represents a cyclopentyl, cyclohexyl- or a phenyl group which may or may not be substituted with an alkyl group having 1 to 4 carbon atoms, or

15  $R^3$  represents a phenyl group substituted with a  $OR^{31}$  group, where  $R^{31}$  is H or an alkyl group having 1 to 4 carbon atoms and k is 0, 1, or

$R^3$  represents a 1-naphthyl or 2-naphthyl group and k is 0, 1, or

20

$R^3$  represent a cis- or trans-decalin group and k is 0, 1, or


25  $R^3$  represents 4-pyridyl, 3-pyrrolidyl or 3-indolyl which may or may not be substituted with a  $OR^{31}$  group, where  $R^{31}$  is as defined above and k is 0, 1, or

$R^3$  represents  $Si(Me)_3$  or  $CH(R^{32})_2$ , wherein  $R^{32}$  is a cyclohexyl- or a phenyl group;

30

$R^4$  represents H, an alkyl group having 1 to 4 carbon atoms, a cyclohexyl- or a phenyl group;

35  $A^2$  represents a structural fragment of Formula IIIa, IIIb or IIIc



10      wherein:

$p$  is an interger 0, 1 or 2;

$m$  is an integer 1, 2, 3 or 4;

15

$Y$  represents a methylene group, or

$Y$  represents an ethylene group and the resulting 5-membered ring may or may not carry one or two fluorine atoms, a hydroxy group or an oxo group in position 4, or may or may not be unsaturated, or

20

$Y$  represents  $-\text{CH}_2-\text{O}-$ ,  $-\text{CH}_2-\text{S}-$ ,  $-\text{CH}_2-\text{SO}-$ , with the heteroatom functionality in position 4, or

25

$Y$  represents a  $n$ -propylene group and the resulting 6-membered ring may or may not carry in position 5 one fluorine atom, a hydroxy group or an oxo group, carry two fluorine atoms in one of positions 4 or 5 or be unsaturated in position 4 and 5, or carry in position 4 an alkyl group with 1 to 4 carbon atoms, or

30

$Y$  represents  $-\text{CH}_2-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{S}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{SO}-\text{CH}_2-$ , or

35       $Y$  represent  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ;


$R^3$  is as defined ab ve;

$R^5$  represents H or an alkyl group having 1 to 4 carbon at ms, or

$R^5$  represents  $-(CH_2)_p-COOR^{51}$ , where p is 0, 1 or 2 and  
 5  $R^{51}$  is H or an alkyl group having 1 to 4 carbon atoms;

n is an integer 0, 1, 2, 3 or 4;

10 B represents a structural fragment of Formula IVa, IVb, IVc or IVd



20

wherein:

r is an interger 0 or 1;

25  $X^1$  represent  $CH_2$ , NH or is absent;

$X^2$  represents  $CH_2$ , NH or  $C=NH$ ;

30  $X^3$  represents NH,  $C=NH$ ,  $N-C(NH)-NH_2$ ,  $CH-C(NH)-NH_2$ ,  $CH-NH-C(NH)-NH_2$  or  $CH-CH_2-C(NH)-NH_2$ ;

$X^4$  represents  $CH_2$  or NH;

Preferred combinations of  $X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$  and r are

35

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and r is 0, 1, or,

- $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$  is 0, 1,  
 or  
 $X^1$  and  $X^3$  are  $NH$ ,  $X^2$  is  $C=NH$ ,  $X^4$  is  $CH_2$  and  $r$  is 0, 1,  
 or  
 5  $X^1$  and  $X^4$  are  $CH_2$ ,  $X^2$  is  $C=NH$ ,  $X^3$  is  $NH$  and  $r$  is 0, 1,  
 or  
 $X^1$  is  $CH_2$ ,  $X^2$  and  $X^4$  are  $NH$ ,  $X^3$  is  $C=NH$  and  $r$  is 1, or  
 $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-NH-C(NH)-NH_2$   
 and  $r$  is 0, 1, or  
 10  $X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and  
 $r$  is 0, or  
 $X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$   
 is 0;
- 15 Particularly preferred combinations of  $X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$   
 and  $r$  are  
 $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and  $r$  is 1;  
 $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$  is 0 or  
 1;  
 20  $X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$   
 is 0;  
 $X^1$  and  $X^3$  are  $NH$ ,  $X^2$  is  $C=NH$ ,  $X^4$  is  $CH_2$  and  $r$  is 1;
- $X^5$  represents  $C(NH)-NH_2$  or  $NH-C(NH)-NH_2$ ;  
 25  $R^6$  is  $H$  or an alkyl group having 1-4 carbon atoms;  
 $X^6$  represents  $CH$  or  $N$ ;
- 30 Compounds of Formula I having  $S$ -configuration on the  $A^2$   
 amino acid are preferred ones, of those compounds also  
 having  $R$ -configuration on the  $A^1$  amino acid are  
 particularly preferred ones.
- 35 In the present context the term "an alkyl group having  
 1 to 4 carbon atoms" may be straight or branched  
 unless specified otherwise. An alkyl group having 1 to


4 carbon atoms may be methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl and t-butyl.

In the present context the term "an alkyl group having 1 to 6 carbon atoms" may be straight or branched unless specified otherwise. An alkyl group having 1 to 6 carbon atoms may be methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, t-pentyl, neo-pentyl, n-hexyl or i-hexyl. When unsaturation is referred to, a carbon-carbon double bond is intended.

The wavy lines on the carbon atom in the carbonyl group in formulas IIa, IIb, IIc, IId, IIe, IIIa, IIIb, IIIc, on the nitrogen atom in formulas IIIa, IIIb, IIIc and on the carbon atom in the ring system in formulas IVa, IVb, IVc, IVd signify the bond position of the fragment.

Abbreviations are listed at the end of this specification.

According to the invention it has been found that compounds of the general Formula Ia, either as such or in the form of physiologically acceptable salts, and including stereoisomers, are potent inhibitors of thrombin:



30

Ia

wherein:

A<sup>1</sup> represents a structural fragment of Formula IIa, IIb, IIc or IId, preferably IIa or IIb;

wherein:

k is an integer 0, 1, 2, 3 or 4, preferably 0, 1;

5 q is an integer 0, 1, 2 or 3, preferably 1;

R<sup>1</sup> represents H, an alkyl group having 1 to 4 carbon atoms, R<sup>11</sup>OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the  
10 position which is alpha to the carbonyl group, and the alpha substituent is a group R<sup>17</sup>-(CH<sub>2</sub>)<sub>p</sub>-, wherein p is 0, 1 or 2 and R<sup>17</sup> is methyl, phenyl, OH, COOR<sup>12</sup>, CONHR<sup>12</sup>, where R<sup>12</sup> is H or an alkyl group having 1 to 4 carbon atoms, and R<sup>11</sup> is H or an alkyl group having 1 to 6  
15 carbon atoms, or

R<sup>1</sup> represents Ph(4-COOR<sup>12</sup>)-CH<sub>2</sub>-, where R<sup>12</sup> is as defined above, or

20 R<sup>1</sup> represents R<sup>13</sup>-NH-CO-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where R<sup>13</sup> is H or an alkyl group having 1 to 4 carbon atoms or -CH<sub>2</sub>COOR<sup>12</sup> where R<sup>12</sup> is as  
25 defined above, or

R<sup>1</sup> represents R<sup>12</sup>OOC-CH<sub>2</sub>-OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group  
30 having 1 to 4 carbon atoms and where R<sup>12</sup> is as defined above, or

R<sup>1</sup> represents R<sup>14</sup>SO<sub>2</sub>-, Ph(4-COOR<sup>12</sup>)-SO<sub>2</sub>-, Ph(3-COOR<sup>12</sup>)-SO<sub>2</sub>-, Ph(2-COOR<sup>12</sup>)-SO<sub>2</sub>- where R<sup>12</sup> is as defined above  
35 and R<sup>14</sup> is an alkyl group having 1-4 carbon atoms, or

R<sup>1</sup> represents -CO-R<sup>15</sup>, wherein R<sup>15</sup> is an alkyl group

having 1-4 carbon atoms,  $r$

$R^1$  represents  $-\text{CO}-\text{OR}^{15}$ , where  $R^{15}$  is as defined above,  
or

5

$R^1$  represent  $-\text{CO}-(\text{CH}_2)_p-\text{COOR}^{12}$ , where  $R^{12}$  is as defined  
above and  $p$  is an interger 0, 1 or 2, or

10

$R^1$  represents  $-\text{CH}_2\text{PO}(\text{OR}^{16})_2$ ,  $-\text{CH}_2\text{SO}_3\text{H}$  or  
 $-\text{CH}_2-(5-(1\text{H})\text{-tetrazolyl})$ , where  $R^{16}$  is, individually at  
each occurrence, H, methyl or ethyl;

Preferably  $R^1$  represents  $R^{11}\text{OOC-alkyl-}$ , where the alkyl  
group has 1 to 4 carbon atoms and  $R^{11}$  is H.

15

$R^2$  represents H or an alkyl group having 1 to 4 carbon  
atoms, or  $R^{21}\text{OOC-alkyl-}$ , where the alkyl group has 1 to  
4 carbon atoms and  $R^{21}$  is H or an alkyl group having 1  
to 4 carbon atoms;

20

$R^3$  represents an alkyl group having 1-4 carbon atoms,  
and the alkyl group may or may not carry one or more  
fluorine atoms, or

25

$R^3$  represents a cyclopentyl, cyclohexyl- or a phenyl  
group which may or may not be substituted with an alkyl  
group having 1 to 4 carbon atoms, or

30

$R^3$  represents a 1-naphthyl or 2-naphthyl group and  $k$   
is 0, 1, or

$R^3$  represent a cis- or trans-decalin group and  $k$  is 0,  
1, or

35

$R^3$  represents  $\text{Si}(\text{Me})_3$  or  $\text{CH}(\text{R}^{32})_2$ , wherein  $R^{32}$  is a  
cyclohexyl- or phenyl group;



$R^4$  represents an alkyl group having 1 to 4 carbon atoms, a cyclohexyl or a phenyl group, preferably a cyclohexyl or a phenyl group;

- 5  $A^2$  represents a structural fragment of Formula IIIa, IIIb or IIIc, preferably IIIa;

wherein:

- 10  $p$  is an interger 0, 1 or 2;

$m$  is an integer 1, 2, 3 or 4, preferably 2, 3;

$Y$  represents a methylene group, or

15

$Y$  represents an ethylene group and the resulting 5-membered ring may or may not carry one or two fluorine atoms, a hydroxy group or an oxo group in position 4, or may or may not be unsaturated, or

20

$Y$  represents  $-\text{CH}_2-\text{O}-$ ,  $-\text{CH}_2-\text{S}-$ ,  $-\text{CH}_2-\text{SO}-$ , with the heteroatom functionality in position 4, or

- 25  $Y$  represents a n-propylene group and the resulting 6-membered ring may or may not carry in position 5 one fluorine atom, a hydroxy group or an oxo group, carry two fluorine atoms in one of positions 4 or 5 or be unsaturated in position 4 and 5, or carry in position 4 an alkyl group with 1 to 4 carbon atoms, or

30

$Y$  represents  $-\text{CH}_2-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{S}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{SO}-\text{CH}_2-$ , or

$Y$  represent  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ;

- 35  $R^3$  represents an alkyl group having 1-4 carbon atoms, or

$R^3$  represents a  $Si(Me)_3$  group;

$R^5$  represents H or an alkyl group having 1 to 4 carbon atoms, preferably H or a methylgroup, or

5

$R^5$  represents  $-(CH_2)_p-COOR^{51}$ , where p is 0, 1 or 2 and  $R^{51}$  is H or an alkyl group having 1 to 4 carbon atoms, preferably p is 0 and  $R^{51}$  is H;

10 n is an integer 0, 1, 2, 3 or 4, preferably 1, 2, 3;

B represents a structural fragment of Formula IVa, IVb, IVc or IVd, preferably IVa or IVb

15 wherein:

$X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$ ,  $X^5$  and  $X^6$  are as defined above;

r is an integer 0 or 1;

20

$R^6$  is H or an alkyl group having 1-4 carbon atoms, preferably H;

preferred combinations of  $X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$  and r are

25

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and r is 0 or 1, or

30

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and r is 0 or 1, or

$X^1$  and  $X^3$  are NH,  $X^2$  is  $C=NH$ ,  $X^4$  is  $CH_2$  and r is 0 or 1, or

35  $X^1$  and  $X^4$  are  $CH_2$ ,  $X^2$  is  $C=NH$ ,  $X^3$  is NH and r is 0 or 1, or

$X^1$  is  $CH_2$ ,  $X^2$  and  $X^4$  are  $NH$ ,  $X^3$  is  $C=NH$  and  $r$  is 1, or

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-NH-C(NH)-NH_2$  and  $r = 0$  or 1, or

5

or  $X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and  $r$  is 0,

10

or  $X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$  is 0;

Particularly preferred combinations of  $X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$  and  $r$  are

15

$X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$  is 0, or

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and  $r = 1$ , or

20

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r = 0$  or 1, or

$X^1$  and  $X^3$  are  $NH$ ,  $X^2$  is  $C=NH$ ,  $X^4$  is  $CH_2$   $r$  is 1;

25

$X^5$  represents  $C(NH)-NH_2$  or  $NH-C(NH)-NH_2$ , preferably  $C(NH)-NH_2$ ;

$X^6$  represents  $CH$  or  $N$ ;

30

According to a preferred embodiment the invention relates to compounds of Formula Ia,

wherein:

35

$A^1$  represents a structural fragment of Formula IIa,

wherein:

k is 0 or 1;

5  $R^1$  represents  $R^{11}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms, particularly methylene, ethylene and  $R^{11}$  is H;

$R^2$  represents H;

10

$R^3$  represents a cyclohexyl group;

$A^2$  represents a structural fragment of Formula IIIa, wherein:

15

Y represents a methylene group, an ethylene group, or a n-propylene group and the resulting 6-membered ring may or may not carry in position 4 an alkyl group with 1 to 4 carbon atoms, preferably Y represents methylene, ethylene;

20

$R^5$  represents H;

B represents a structural fragment of formula IVa

25

wherein:

$X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$ , r is 0 and n is 1 or 2;

30

$X^1$ , and  $X^3$  are NH,  $X^2$  is  $C=NH$ ,  $X^4$  is  $CH_2$ , r is 1 and n is 2, or

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$ , r is 1 and n is 1, or

35

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$ , r is 0 or 1

and n is 1 or 2, or

More particularly preferred are compounds wherein B represents a structural fragment of formula IVb wherein:

$X^5$  represents  $C(NH)-NH_2$ ,  $R^6$  is H, and  $n = 1$

Preferred compounds of the invention are:

- 10  $HOOC-CH_2-(R)Cgl-Aze-Pab$   
 $HOOC-CH_2-CH_2-(R)Cgl-Aze-Pab$   
 $HOOC-CH_2-(R)Cgl-Pro-Pab$   
 $HOOC-CH_2-CH_2-(R)Cgl-Pro-Pab$   
 15  $(HOOC-CH_2)_2-(R)Cgl-Pro-Pab$   
 $H-(R)Cgl-Pic-Pab$   
 $HOOC-CH_2-(R,S)CH(COOH)-(R)Cgl-Pic-Pab$   
 $H-(R)Cha-Aze-Pab$   
 $HOOC-CH_2-(R)Cha-Aze-Pab$   
 20  $HOOC-CH_2-(R,S)CH(COOH)-(R)Cha-Aze-Pab$   
 $HOOC-CH_2-(RorS)CH(COOH)-(R)Cha-Aze-Pab/a$   
 $HOOC-CH_2-(RorS)CH(COOH)-(R)Cha-Aze-Pab/b$   
 $HOOC-CH_2-CH_2-(R)Cha-Aze-Pab$   
 $HOOC-CH_2-NH-CO-CH_2-(R)Cha-Aze-Pab$   
 25  $H-(R)Cha-Pro-Pab$   
 $HOOC-CH_2-(R)Cha-Pro-Pab$   
 $HOOC-CH_2-(Me)(R)Cha-Pro-Pab$   
 $HOOC-CH_2-CH_2-(R)Cha-Pro-Pab$   
 $HOOC-CH_2-CH_2-(Me)(R)Cha-Pro-Pab$   
 30  $HOOC-CH_2-(RorS)CH(COOH)-(R)Cha-Pro-Pab/a$   
 $HOOC-CH_2-(RorS)CH(COOH)-(R)Cha-Pro-Pab/b$   
 $HOOC-CH_2-NH-CO-CH_2-(R)Cha-Pro-Pab$   
 $EtOOC-CH_2-CH_2-CH_2-(R)Cha-Pro-Pab$   
 $Ph(4-COOH)-SO_2-(R)Cha-Pro-Pab$   
 35  $H-(R)Cha-Pic-Pab$   
 $HOOC-CH_2-(R)Cha-Pic-Pab$   
 $HOOC-CH_2-(R or S)CH(COOH)-(R)Cha-Pic-Pab/a$

- HOOC-CH<sub>2</sub>-(RorS)CH(COOH)-(R)Cha-Pic-Pab/b  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pic-Pab  
 HOOC-CO-(R)Cha-Pic-Pab  
 HOOC-CH<sub>2</sub>-CO-(R)Cha-Pic-Pab  
 5 Me-OOC-CH<sub>2</sub>-CO-(R)Cha-Pic-Pab  
 H<sub>2</sub>N-CO-CH<sub>2</sub>-(R)Cha-Pic-Pab  
 Boc-(R)Cha-Pic-Pab  
 Ac-(R)Cha-Pic-Pab  
 Me-SO<sub>2</sub>-(R)Cha-Pic-Pab  
 10 H-(R)Cha-(R,S)betaPic-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)betaPic-Pab  
 HOOC-CH<sub>2</sub>-(R)Cha-Val-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Val-Pab  
 H-(R)Hoc-Aze-Pab  
 15 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Hoc-Aze-Pab  
 HOOC-CH<sub>2</sub>-(R,S)CH(COOH)-(R)Hoc-Pro-Pab  
 HOOC-CH<sub>2</sub>-(R)Hoc-Pic-Pab  
 (HOOC-CH<sub>2</sub>)<sub>2</sub>-(R)Hoc-Pic-Pab  
 HOOC-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab  
 20 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Tic-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pig  
 HOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pig  
 H-(R)Cha-Aze-Pig  
 25 HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pac  
 H-(R)Cha-Pro-Pac  
 H-(R)Cgl-Ile-Pab  
 H-(R)Cgl-Aze-Pab  
 HOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab  
 30 MeOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab  
 EtOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab  
<sup>n</sup>BuOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab  
<sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab  
 H-(R)Cgl-Pro-Pac  
 35 HOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pr -Pac  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Aze-Pac

- HOOC-CH<sub>2</sub>-(R) Cha-Aze-Pig  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pig  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pig  
 (HOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Cgl-Pro-Pig  
 5 HOOC-CH<sub>2</sub>-CH<sub>2</sub>(HOOC-CH<sub>2</sub>)-(R) Cha-Pro-Pig  
 HOOC-CH<sub>2</sub>-(R) Cgl-Aze-(R,S) Itp  
 HOOC-CH<sub>2</sub>-(R) Cha-Aze-(R,S) Itp  
 H-(R) Cha-Pic-(R,S) Itp  
 HOOC-CH<sub>2</sub>-(R) Cha-Pic-(R,S) Itp  
 10 H-(R) Cgl-Pro-(R,S) Hig  
 HOOC-CH<sub>2</sub>-(R) Cgl-Pro-(R,S) Hig  
 H-(R) Cha-Pro-(R,S) Hig  
 H-(R) Cgl-Aze-Rig  
 HOOC-CH<sub>2</sub>-(R) Cgl-Aze-Rig  
 15 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Rig  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Rig  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-(S) Itp  
 H-(R) Cha-Pro-(R,S) Nig  
 H-(R) Cha-Pro-Mig  
 20 H-(R) Cha-Pro-Dig  
 H-(R) Cha-Aze-Dig

At present the particularly preferred compounds of formula Ia is

- 25 HOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Pab  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pab  
 30 HOOC-CH<sub>2</sub>-(R) Cha-Pic-Pab  
 HOOC-CH<sub>2</sub>-(R) Cgl-Pro-Pig  
 EtOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pac  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pig  
 35


In the above tables of compounds, the letters /a and /b refer to a substantially pure stereoisomer at the

carbon atom noted "RorS". The stereoisomer can be identified for each compound with reference to the experimental part herein. "R,S" refers to a mixture of stereoisomers.

5

According to the invention it has been found that compounds of the general Formula Ib, either as such or in the form of physiologically acceptable salts, and including stereoisomers, are potent inhibitors of kininogenases:

10



Ib

15 wherein:

A<sup>1</sup> represents a structural fragment of formula IIa, IIb or IIe, preferably IIa or IIb;

wherein:

20

k is an integer 0, 1, 2, 3 or 4, preferably 0, 1;

q is an integer 0, 1, 2, or 3, preferably 1;

25

R<sup>1</sup> represents H, an alkyl group having 1 to 4 carbon atoms, or R<sup>11</sup>OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group R<sup>17</sup>-(CH<sub>2</sub>)<sub>p</sub>-, wherein p is 0, 1 or 2 and R<sup>17</sup> is methyl, phenyl, OH, COOR<sup>12</sup>, CONHR<sup>12</sup>, where R<sup>12</sup> is H or an alkyl group having 1 to 4 carbon atoms, and R<sup>11</sup> is H or an alkyl group having 1 to 6 carbon atoms, or

30

35 R<sup>1</sup> represents Ph(4-COOR<sup>12</sup>)-CH<sub>2</sub>-, where R<sup>12</sup> is H or an alkyl group having 1 to 4 carbon atoms, or



5  $R^1$  represents  $R^{13}$ -NH-CO-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{13}$  is H or an alkyl group having 1 to 4 carbon atoms or  $-CH_2COOR^{12}$  where  $R^{12}$  is as defined above, or

10  $R^1$  represents  $R^{12}OOC-CH_2-OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{12}$  is as defined above, or

15  $R^1$  represents  $R^{14}SO_2$ -,  $Ph(4-COOR^{12})-SO_2$ -,  $Ph(3-COOR^{12})-SO_2$ -,  $Ph(2-COOR^{12})-SO_2$ -, where  $R^{12}$  is as defined above and  $R^{14}$  is an alkyl group having 1-4 carbon atoms, or

20  $R^1$  represents  $-CO-R^{15}$ , wherein  $R^{15}$  is an alkyl group having 1-4 carbon atoms, or

$R^1$  represents  $-CO-OR^{15}$ , where  $R^{15}$  is as defined above, or

25  $R^1$  represents  $-CO-(CH_2)_p-COOR^{12}$ , where  $R^{12}$  is as defined above and p is 0, 1 or 2, or

$R^1$  represents  $-CH_2PO(OR^{16})_2$ ,  $-CH_2SO_3H$  or  $-CH_2-(5-(1H)-tetrazolyl)$ , where  $R^{16}$  is, individually at each occurrence, H, methyl or ethyl;

30  $R^2$  represents H or an alkyl group having 1 to 4 carbon atoms or  $R^{21}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and  $R^{21}$  is H or an alkyl group having 1 to 4 carbon atoms;

35  $R^3$  represents an alkyl group having 1-4 carbon atoms, and the alkyl group may or may not carry one or more

fluorin atoms, or

5  $R^3$  represents a cyclopentyl, cyclohexyl- or a phenyl group which may or may not be substituted with an alkyl group having 1 to 4 carbon atoms, or

10  $R^3$  represents a phenyl group substituted with a  $OR^{31}$  group, where  $R^{31}$  is H or an alkyl group having 1 to 4 carbon atoms and k is 0, 1, or

$R^3$  represents a 1-naphthyl or 1-naphthyl group and k is 0, 1, or

15  $R^3$  represent a cis- or trans-decalin group and k is 0,1, or

20  $R^3$  represents 4-pyridyl, 3-pyrrolidyl or 3-indolyl which may or may not be substituted with a  $OR^{31}$  group, where  $R^{31}$  is as defined above and k is 0, 1, or

$R^3$  represents  $Si(Me)_3$  or  $CH(R^{32})_2$ , wherein  $R^{32}$  is a cyclohexyl- or phenyl group;

25  $R^4$  represents H, an alkyl group having 1 to carbon atoms, a cyclohexyl or a phenyl group, preferably H;

$A^2$  represents a structural fragment of formula IIIb or IIIc, preferably IIIb

30 wherein:

p is an integer 0, 1 or 2;

m is an integer 1, 2, 3, or 4, preferably 2, 3;

$R^3$  is as defined above;

35 n is an integer 0, 1, 2, 3 or 4, preferably 1,2,3;

B represents a structural fragment of Formula IVa, IVb, IVc or IVd, preferably IVa or IVb;

wherein:

5

$X^1$ ,  $X^2$ ,  $X^3$ ,  $X^4$  are as defined above;

$R^6$  is H or an alkyl group having 1-4 carbon atoms, preferably H or a methyl group;

10

r is an integer 0 or 1;

preferred combinations of  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  are

15

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and r is 0 or 1, or

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and r is 0 or 1, or

20

$X^1$  and  $X^3$  are NH,  $X^2$  is  $C=NH$ ,  $X^4$  is  $CH_2$  and r is 0 or 1, or

25

$X^1$  and  $X^4$  are  $CH_2$ ,  $X^2$  is  $C=NH$ ,  $X^3$  is NH and r is 0 or 1, or

$X^1$  is  $CH_2$ ,  $X^2$  and  $X^4$  are NH,  $X^3$  is  $C=NH$  and r is 1, or

30

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-NH-C(NH)-NH_2$  and r is 0 or 1,

or  $X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and r is 0, or

35

$X^1$  is absent,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and r is 0;

particularly preferred combinations of  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  are

5  $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$  and  $r$  is 1  
or,

$X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$  and  $r$  is 1;

10  $X^5$  represents  $C(NH)-NH_2$  or  $NH-C(NH)-NH_2$ , preferably  
 $C(NH)-NH_2$ ;

$X^6$  represents  $CH$  or  $N$ .

Preferred compound of the invention are:

15

$H-(R)Pro-Phe-Pab$

$HOOC-CH_2-(R)Pro-Phe-Pab$

$H-(R)Phe-Phe-Pab$

$HOOC-CO-(R)Phe-Phe-Pab$

20

$HOOC-CH_2-(R)Phe-Phe-Pab$

$H-(R)Cha-Phe-Pab$

$HOOC-CH_2-(R)Cha-Phe-Pab$

$H-(R)Phe-Cha-Pab$

$HOOC-CH_2-(R)Phe-Cha-Pab$


25

$H-(R)Cha-Cha-Pab$

$HOOC-CH_2-(R)Cha-Cha-Pab$

Furthermore, it has been found that compounds of the  
general Formula V, either as such or in the form of  
30 physiologically acceptable salts, and including  
stereoisomers, are potent inhibitors of serine  
proteases, especially thrombin and kininogenases such  
as kallikrein after oral or parenteral administration:

35




Formula V

wherein:

A<sup>1</sup> represents a structural fragment of Formula  
IIa, IIb, IIc, IIId or IIe;

5



wherein:

k is an integer 0, 1, 2, 3 or 4;

25

m is an integer 1, 2, 3 or 4;

q is an integer 0, 1, 2 or 3;

30 R<sup>1</sup> represents R<sup>11</sup>OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group R<sup>17</sup>-(CH<sub>2</sub>)<sub>p</sub>-, wherein p is 0, 1 or 2 and R<sup>17</sup> is COOR<sup>12</sup>, CONHR<sup>12</sup>, where R<sup>12</sup> is H or an

35 alkyl group having 1 to 4 carbon atoms or a benzyl group, and R<sup>11</sup> is H or an alkyl group having 1 to 6 carbon atoms, or a benzyl group, or

$R^1$  represents  $\text{Ph}(4\text{-COOR}^{12})\text{-CH}_2\text{-}$ , where  $R^{12}$  is as defined above, or

5  $R^1$  represents  $R^{13}\text{-NH-CO-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{13}$  is H or an alkyl group having 1 to 4 carbon atoms or  $\text{-CH}_2\text{COOR}^{12}$ , where  $R^{12}$  is as defined above, or

10  $R^1$  represents  $R^{12}\text{OOC-CH}_2\text{-OOC-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{12}$  is as defined above, or

15  $R^1$  represents  $R^{14}\text{SO}_2\text{-}$ ,  $\text{Ph}(4\text{-COOR}^{12})\text{-SO}_2\text{-}$ ,  $\text{Ph}(3\text{-COOR}^{12})\text{-SO}_2\text{-}$ ,  $\text{Ph}(2\text{-COOR}^{12})\text{-SO}_2\text{-}$ , where  $R^{12}$  is as defined above and  $R^{14}$  is an alkyl group having 1-4 carbon atoms, or

20  $R^1$  represents  $\text{-CO-R}^{15}$ , wherein  $R^{15}$  is an alkyl group having 1-4 carbon atoms, or

25  $R^1$  represents  $\text{-CO-OR}^{15}$ , where  $R^{15}$  is as defined above, or

$R^1$  represent  $\text{-CO-(CH}_2)_p\text{-COOR}^{12}$ , where  $R^{12}$  is as defined above and p is an interger 0, 1 or 2, or

30  $R^2$  represents H or an alkyl group having 1 to 4 carbon atoms or  $R^{21}\text{OOC-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms and, where  $R^{21}$  is H, an alkyl group having 1 to 4 carbon atoms or a benzyl group;

35  $R^3$  represents an alkyl group having 1-4 carbon atoms, and the alkyl gr up may r may not carry ne r more flourin atoms, or

$R^3$  represents a cycl pentyl, cyclohexyl- or a phenyl group which may or may not be substituted with an alkyl group having 1 to 4 carbon atoms, or

5  $R^3$  represents a phenyl group substituted with a  $OR^{31}$  group, where  $R^{31}$  is H or an alkyl group having 1 to 4 carbon atoms and k is 0, 1, or

10  $R^3$  represents a 1-naphthyl or 2-naphthyl group and k is 0, 1, or

$R^3$  represent a cis- or trans-decalin group and k is 0, 1, or

15  $R^3$  represents 4-pyridyl, 3-pyrrolidyl or 3-indolyl which may or may not be substituted with a  $OR^{31}$  group, where  $R^{31}$  is as defined above and k is 0, 1, or

20  $R^3$  represents  $Si(Me)_3$  or  $CH(R^{32})_2$ , wherein  $R^{32}$  is a cyclohexyl- or a phenyl group;

$R^4$  represents H, an alkyl group having 1 to 4 carbon atoms, a cyclohexyl- or a phenyl group;

25  $A^2$ , B and n are defined as described under Formula I above;

D is Z or  $(Z)_2$ , wherein Z represents a benzyloxycarbonyl group.


30

The benzyloxycarbonyl group (Z or  $(Z)_2$ ) will bind to the amidino- or guanidino nitrogens present in B.

35 Preferred and particularly preferred combinations are the same as described for Formula I above.

Furtherm re, it has been found that compounds of the

general Formula Va, either as such or in the form of physiologically acceptable salts, and including stereoisomers, are potent inhibitors of thrombin after oral or parenteral administration:



Formula Va

wherein:

$A^1$  represents a structural fragment of Formula IIa, IIb, IIc or IId, preferably IIa or IIb;

wherein:

$k$  is an integer 0, 1, 2, 3 or 4, preferably 0, 1;

$q$  is an integer 0, 1, 2 or 3, preferably 1;

$R^1$  represents  $R^{11}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group  $R^{17}-(CH_2)_p$ -, wherein  $p$  is 0, 1 or 2 and  $R^{17}$  is  $COOR^{12}$ ,  $CONHR^{12}$ , where  $R^{12}$  is H, an alkyl group having 1 to 4 carbon atoms or a benzyl group, and  $R^{11}$  is H or an alkyl group having 1 to 6 carbon atoms, or a benzyl group, or

$R^1$  represents  $Ph(4-COOR^{12})-CH_2$ -, where  $R^{12}$  is as defined above, or

$R^1$  represents  $R^{13}-NH-CO$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{13}$  is H or an alkyl group having



1 to 4 carbon atoms  $r$   $-\text{CH}_2\text{COOR}^{12}$  where  $\text{R}^{12}$  is as defined above, or

5  $\text{R}^1$  represents  $\text{R}^{12}\text{OOC}-\text{CH}_2-\text{OOC}-\text{alkyl}-$ , where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $\text{R}^{12}$  is as defined above, or

10  $\text{R}^1$  represents  $\text{R}^{14}\text{SO}_2-$ ,  $\text{Ph}(4-\text{COOR}^{12})-\text{SO}_2-$ ,  $\text{Ph}(3-\text{COOR}^{12})-\text{SO}_2-$ ,  $\text{Ph}(2-\text{COOR}^{12})-\text{SO}_2-$  where  $\text{R}^{12}$  is as defined above and  $\text{R}^{14}$  is an alkyl group having 1-4 carbon atoms, or

15  $\text{R}^1$  represents  $-\text{CO}-\text{R}^{15}$ , wherein  $\text{R}^{15}$  is an alkyl group having 1-4 carbon atoms, or

$\text{R}^1$  represents  $-\text{CO}-\text{OR}^{15}$ , where  $\text{R}^{15}$  is as defined above, or

20  $\text{R}^1$  represent  $-\text{CO}-(\text{CH}_2)_p-\text{COOR}^{12}$ , where  $\text{R}^{12}$  is as defined above and  $p$  is an interger 0, 1 or 2, or

25 Preferably  $\text{R}^1$  represents  $\text{R}^{11}\text{OOC}-\text{alkyl}-$ , where the alkyl group has 1 to 4 carbon atoms and  $\text{R}^{11}$  is as defined above.

30  $\text{R}^2$  represents H or an alkyl group having 1 to 4 carbon atoms, or  $\text{R}^{21}\text{OOC}-\text{alkyl}-$ , where the alkyl group has 1 to 4 carbon atoms and  $\text{R}^{21}$  is H or an alkyl group having 1 to 4 carbon atoms or a benzyl group;

$\text{R}^3$  represents an alkyl group having 1-4 carbon atoms, and the alkyl group may or may not carry one or more fluorine atoms, or

35  $\text{R}^3$  represents a cycl pentyl, cyclohexyl-  $r$  a phenyl group which may  $r$  may not be substituted with an alkyl

group having 1 to 4 carbon atoms, or

$R^3$  represents a 1-naphthyl or 2-naphthyl group and  $k$  is 0, 1, or

5

$R^3$  represent a cis- or trans-decalin group and  $k$  is 0, 1, or

10

$R^3$  represents  $Si(Me)_3$  or  $CH(R^{32})_2$ , wherein  $R^{32}$  is a cyclohexyl- or phenyl group;

15

$R^4$  represents an alkyl group having 1 to 4 carbon atoms, a cyclohexyl or a phenyl group, preferably a cyclohexyl or a phenyl group;

$A^2$ ,  $B$  and  $n$  are defined as described under Formula Ia above;

20

$D$  is  $Z$  or  $(Z)_2$ ;

$Z$  represents a benzyloxycarbonyl group.

25

Preferred integers, groups or combinations and particularly preferred combinations are the same as described for Formula Ia above but  $R^{11}$  is H, an alkyl group having 1 to 6 carbon atoms or a benzyl group.

30

Preferred compounds having Formula Va are:

35

$BnOOC-CH_2-(R)Cgl-Aze-Pab(Z)$

$BnOOC-CH_2-CH_2-(R)Cgl-Aze-Pab(Z)$

$BnOOC-CH_2-(R)Cgl-Pro-Pab(Z)$

$BnOOC-CH_2-CH_2-(R)Cgl-Pro-Pab(Z)$

$(BnOOC-CH_2)_2-(R)Cgl-Pro-Pab(Z)$

$BnOOC-CH_2-(R,S)CH(COOBn)-(R)Cgl-Pic-Pab(Z)$

$BnOOC-CH_2-(R)Cha-Aze-Pab(Z)$

- BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(RorS)CH(COOBn)-(R)Cha-Az -Pab(Z)/a  
 BnOOC-CH<sub>2</sub>-(RorS)CH(COOBn)-(R)Cha-Az -Pab(Z)/b  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Aze-Pab(Z)  
 5 BnOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R)Cha-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(Me)(R)Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(Me)(R)Cha-Pro-Pab(Z)  
 10 BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z)  
 Ph(4-COOH)-SO<sub>2</sub>-(R)Cha-Pro-Pab(Z)  
 Boc-(R)Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pic-Pab(Z)  
 15 BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pic-Pab(Z)  
 EtOOC-CO-(R)Cha-Pic-Pab(Z)  
 MeOOC-CH<sub>2</sub>-CO-(R)Cha-Pic-Pab(Z)  
 H<sub>2</sub>N-CO-CH<sub>2</sub>-(R)Cha-Pic-Pab(Z)  
 20 Ac-(R)Cha-Pic-Pab(Z)  
 Me-SO<sub>2</sub>-(R)Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Val-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)Val-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Hoc-Aze-Pab(Z)  
 25 BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Hoc-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Hoc-Pic-Pab(Z)  
 (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R)Hoc-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab(Z)  
 30 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Tic-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pig(Z)<sub>2</sub>  
 BnOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pig(Z)<sub>2</sub>  
 BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pac(Z)  
 BnOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab(Z)  
 35 MeOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
 EtOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
<sup>n</sup>BuOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)

- <sup>n</sup>HexOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Pro-Pac(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Pro-Pac(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Pac(Z)  
 5 BnOOC-CH<sub>2</sub>-(R) Cha-Aze-Pig(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Pro-Pig(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pig(Z)  
 (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Cgl-Pro-Pig(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>(BnOOC-CH<sub>2</sub>)-(R) Cha-Pro-Pig(Z)  
 10 BnOOC-CH<sub>2</sub>-(R) Cha-Pic-(R,S) Itp(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cgl-Pro-(R,S) Hig(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cgl-Aze-Rig(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Pro-Rig(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Rig(Z)


15

Particularly preferred compounds are:

- BnOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Pro-Pab(Z)  
 20 BnOOC-CH<sub>2</sub>-(R) Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cgl-Pro-Pig(Z)<sub>2</sub>  
 EtOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Pro-Pac(Z)  
 Bn OOC-CH<sub>2</sub>-(R) Cha-Pro-Pig(Z)

25

- Furthermore, it has been found that compounds of the  
 general Formula Vb, either as such or in the form of  
 physiologically acceptable salts, and including  
 30 stereoisomers, are potent inhibitors of kallikrein  
 after oral or parenteral administration:



35

Formula Vb

wh rein:

A<sup>1</sup> represents a structural fragment of formula IIa, IIb or IIe, preferably IIa or IIb;

5 wherein:

k is an integer 0, 1, 2, 3 or 4, preferably 0, 1;

q is an integer 0, 1, 2, or 3, preferably 1;

10

R<sup>1</sup> represents R<sup>11</sup>OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group R<sup>17</sup>-(CH<sub>2</sub>)<sub>p</sub>-, wherein p is 0, 1 or 2 and R<sup>17</sup> is COOR<sup>12</sup>, CONHR<sup>12</sup>, where R<sup>11</sup> is H or an alkyl group having 1 to 4 carbon atoms, and R<sup>11</sup> is H or an alkyl group having 1 to 6 carbon atoms, or a benzyl group, or

15

20 R<sup>1</sup> represents Ph(4-COOR<sup>12</sup>)-CH<sub>2</sub>-, where R<sup>12</sup> is as defined above, or

R<sup>1</sup> represents R<sup>13</sup>-NH-CO-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where R<sup>13</sup> is H or an alkyl group having 1 to 4 carbon atoms or -CH<sub>2</sub>COOR<sup>12</sup> where R<sup>12</sup> is as defined above, or

25

30 R<sup>1</sup> represents R<sup>12</sup>OOC-CH<sub>2</sub>-OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where R<sup>12</sup> is as defined above, or

35

R<sup>1</sup> represents R<sup>14</sup>SO<sub>2</sub>-, Ph(4-COOR<sup>12</sup>)-SO<sub>2</sub>-, Ph(3-COOR<sup>12</sup>)-SO<sub>2</sub>-, Ph(2-COOR<sup>12</sup>)-SO<sub>2</sub>-, wh r R<sup>12</sup> is as defined above

and  $R^{14}$  is an alkylgroup having 1-4 carbon atoms, or

$R^1$  represents  $-\text{CO}-R^{15}$ , wherein  $R^{15}$  is an alkyl group having 1-4 carbon atoms, or

5

$R^1$  represents  $-\text{CO}-\text{OR}^{15}$ , where  $R^{15}$  is as defined above, or

10

$R^1$  represent  $-\text{CO}-(\text{CH}_2)_p-\text{COOR}^{12}$ , where  $R^{12}$  is as defined above and p is 0, 1 or 2, or

15

$R^2$  represents H or an alkyl group having 1 to 4 carbon atoms or  $R^{21}\text{OOC-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms and  $R^{21}$  is H, an alkyl group having 1 to 4 carbon atoms or a benzyl group;

20

$R^3$  represents an alkyl group having 1-4 carbon atoms, and the alkyl group may or may not carry one or more fluorine atoms, or

25

$R^3$  represents a cyclopentyl, cyclohexyl- or a phenyl group which may or may not be substituted with an alkyl group having 1 to 4 carbon atoms, or

30

$R^3$  represents a phenyl group substituted with a  $\text{OR}^{31}$  group, where  $R^{31}$  is H or an alkyl group having 1 to 4 carbon atoms and k is 0, 1, or

35

$R^3$  represents a 1-naphthyl or 1-naphthyl group and k is 0, 1, or

$R^3$  represent a cis- or trans-decalin group and k is 0,1, or

40

$R^3$  represents 4-pyridyl, 3-pyrrolidyl or 3-indolyl which may or may not be substitut d with a  $\text{OR}^{31}$  group, where  $R^{31}$  is as defin d above and k is 0, 1, or

$R^3$  represents  $Si(M)_3$  or  $CH(R^{32})_2$ , wherein  $R^{32}$  is a cyclohexyl- or phenyl group;

$R^4$  represents H, an alkyl group having 1 to 6 carbon atoms, a cyclohexyl or a phenyl group, preferably H;

$A^2$ , B and n are defined as described under Formula Ib above;

D represents Z or  $(Z)_2$ .

Preferred integers, groups or combinations and particularly preferred combinations are the same as described in Formula Ib above but  $R^{11}$  is H, an alkyl group having 1 to 6 carbon atoms or a benzyl group.


Preferred compounds having Formula Vb are:

Boc-(R)Pro-Phe-Pab(Z)  
 20 BnOOC-CH<sub>2</sub>-(R)Pro-Phe-Pab(Z)  
 Boc-(R)Phe-Phe-Pab(Z)  
 MeOOC-CO-(R)Phe-Phe-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Phe-Phe-Pab(Z)

25

In a further embodiment the invention relates to novel use of a compound of the formula:

30




35

as a starting material in synthesis of a peptidic serine protease inhibitor, and in particular in synthesis of peptidic thrombin inhibitors or kininogenase inhibitors. It can be used as such or


having the amidino group either mono- or diprotected at the nitrogens with a protective group such as benzyloxy carbonyl. Protection of the amidino derivatives is carried out by methods known in the art for amidino compounds. This compound is named "1-amidino-4-aminomethylbenzene" or "H-Pab" herein. The compound has been previously disclosed in inter alia Biochem. Pharm. vol 23, p. 2247-2256.

The structural fragment of the formula



has however not been previously disclosed as a structural element in a pharmaceutically active compound, especially a peptic compound. The fragment renders a serine protease inhibitor, and in particular a thrombin inhibitor or kininogenases inhibitor valuable.

In a further embodiment the invention relates to novel use of a compound of the formula:



as a starting material in synthesis of a thrombin inhibitor. The compound may have the amidino group either mono- or diprotected at the nitrogens with a protective group such as benzyloxy carbonyl. Protection of the amidino derivatives is carried out by methods known in the art for amidino compounds. This compound




is named "1-amidino-4-aminomethyl cyclohexane" or "H-Pac" herein.

The compound has been previously disclosed in DE 2748295.

5

The structural fragment of the formula

10




has however not been previously disclosed as a structural element in a thrombin inhibitor valuable.

15

In a further embodiment the invention relates to a novel compound of the formula:

20




and the use of said compound as a starting material in synthesis of a serine protease inhibitor, especially a thrombin inhibitor or kininogenase inhibitor. The compound may have the amidino group either mono- or diprotected at the nitrogens with a protective group such as benzyloxy carbonyl. Protection of the amidino derivatives is carried out by methods known in the art for amidno compounds. This compound is named "4-aminoethyl-1-amidino piperidine" or "H-Rig" herein.

30


The structural fragment of the formula

35




has however not been previously disclosed as a structural element in a pharmaceutically active compound, especially a peptic compound. The fragment renders a serine protease inhibitor, and in particular a thrombin inhibitor or kininogenases inhibitor variable.

In a further embodiment the invention relates to a novel compound of the formula:



and the use of said compound as a starting material in synthesis of a serine protease inhibitor especially a thrombin inhibitor or kininogenase inhibitor. The compound may have the amidino group either mono- or diprotected at the nitrogens with a protective group such as benzyloxy carbonyl. Protection of the amidino derivatives is carried out by methods known in the art for amidino compounds. This compound is named "1,3-diaza-2-imino-4-aminoethyl cyclohexane" or "H-Itp" herein.


The structural fragment of the formula



has however not been previously disclosed as a structural element in a pharmaceutically active

compound, especially a peptic compound. The fragment renders a serine protease inhibitor, and in particular a thrombin inhibitor or kininogenases inhibitor variable.

In a further embodiment the invention relates to novel compounds of the formula:



where  $n$  is 1 or 2

$s$  is 0 or 1,

and the use of said compounds as a starting material in synthesis of serine protease inhibitors, especially thrombin inhibitors or kininogenases inhibitors. The compound may have the amidino group either mono- or diprotected at the nitrogens with a protective group such as benzyloxy carbonyl. Protection of the amidino derivatives is carried out by methods known in the art for amidino compounds. These compounds are named:


1-amidino-3-aminomethyl pyrrolidine or "H-Nig" when  $n$  is 1 and  $s$  is 1

1-amidino-3-aminoethyl pyrrolidine or "H-Hig" when  $n$  is 2 and  $s$  is 1

3-aminomethyl-1-amidino azetidine or "H-Mig" when  $n$  is 1 and  $s$  is 0

3-aminoethyl-1-amidino azetidine or "H-Dig" when  $n$  is 2 and  $s$  is 0

The structural fragment of the formula



has however not been previously disclosed as a structural element in a pharmaceutically active compound, especially a peptic compound. The fragment renders a serine protease inhibitor, and in particular a thrombin inhibitor or kininogenases inhibitor valuable.

A further embodiment of the invention are the novel compounds having the amidino group mono- or di-protected at the nitrogens with a benzyloxy carbonyl group, examples of such compounds are

4-aminomethyl-1-(N-benzyloxycarbonylamidino) benzene (H-Pab(Z)),  
 4-aminomethyl-1-(N,N'-di(benzyloxycarbonyl)amidino) benzene (H-Pab(Z)<sub>2</sub>),  
 4-aminomethyl-1-(N-benzyloxycarbonylamidino) cyclohexane (H-Pac(Z)),  
 4-aminomethyl-1-(N,N'-di(benzyloxycarbonyl)amidino) cyclohexane (H-Pac(Z)<sub>2</sub>),  
 4-aminoethyl-1-(N-benzyloxy-carbonylamidino) piperidine (H-Rig(Z)),  
 4-aminoethyl-1-N,N'-di(benzyloxycarbonyl)amidino piperidine (H-Rig(Z)<sub>2</sub>),  
 (3RS)-1-(N-benzyloxycarbonylamidino)-3-aminomethyl pyrrolidine (H-Nig(Z)),

(3RS)-1-(N,N'-di(benzyloxycarbonyl)amidino)-3-aminomethyl pyrrolidine (H-Nig(Z)<sub>2</sub>),  
(3RS)-1-(N-benzyloxycarbonylamidino)-3-aminoethyl pyrrolidine (H-Hig(Z)),  
5 (3RS)-1-(N,N'-di(benzyloxycarbonyl)amidino)-3-aminoethyl pyrrolidine (H-Hig(Z)<sub>2</sub>),  
3-aminomethyl-1-(N-benzyloxycarbonylamidino) azetidine (H-Mig(Z)),  
3-aminomethyl-1-(N,N'-di(benzyloxycarbonyl)amidino) azetidine (H-Mig(Z)<sub>2</sub>),  
10 3-aminoethyl-1-(N-benzyloxycarbonylamidino) azetidine (H-Dig(Z)),  
3-aminoethyl-1-(N,N'-di(benzyloxycarbonyl)amidino) azetidine (H-Dig(Z)<sub>2</sub>),  
15

Said compounds are used as starting materials in the preparation of the claimed peptide derivatives of formulas I, Ia, Ib, V, Va and Vb.

## 20 Medical and pharmaceutical use

The invention also provides compositions and methods for the treatment, in a human or animal organism, of conditions where inhibition of thrombin is required and  
25 of physiologically disorders especially inflammatory diseases.

The thrombin inhibiting compounds of the invention are expected to be useful in particular in animals  
30 including man in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. They are furthermore expected to be useful in situations where there is an undesirable excess of the thrombin without signs of hypercoagulability, for example as in  
35 Alzheimers disease and pancreatitis. Disease states in which these compounds have a potential utility, in treatment and/or prophylaxis, include venous thrombosis

and pulmonary embolism, arterial thrombosis, such as in myocardial infarction, unstable angina, thrombosis-based stroke and peripheral arterial thrombosis and systemic embolism usually from the atrium during

5 arterial fibrillation or from the left ventricle after transmural myocardial infarction. Further, these compounds have expected utility in prophylaxis of atherosclerotic diseases such as coronary arterial disease, cerebral arterial disease and peripheral

10 arterial disease. Further, these compounds are expected to have synergistic antithrombotic effects when combined with any antithrombotic agent with a different mechanism of action, such as the antiplatelet agent acetylsalicylic acid. Further, these compounds are

15 expected to be useful together with thrombolytics in thrombotic diseases, in particular myocardial infarction. Further, these compounds have expected utility in prophylaxis for re-occlusion after thrombolysis, percutaneous trans-luminal angioplasty

20 (PTCA) and coronary bypass operations. Further, these compounds have expected utility in prevention of re-thrombosis after microsurgery and vascular surgery in general. Further, these compounds have expected utility in treatment and prophylaxis of disseminated

25 intravascular coagulation caused by bacteria, multiple trauma, intoxication or any other mechanism. Further, these compounds are expected to be useful in anticoagulant treatment when blood is in contact with foreign surfaces in the body such as vascular grafts,

30 vasculars stemts, vascular catheters, mechanical and biological prosthetic or any other medical device. Further, these compounds have expected utility in anticoagulant treatment when blood is in contact with

35 medical devices outside the body such as during cardiovascular surgery using or heart-lung machine or in haemodialysis.

A further expected utility of the anticoagulant compounds of the invention are in rinsing of catheters and mechanical devices used in patients in vivo, and as anticoagulants for preservation of blood, plasma and other blood products in vitro.

The antiinflammatory inhibiting compounds of the invention are expected to be useful in particular in animals including man in treatment or prophylaxis of inflammatory diseases such as asthma, rhinitis, pancreatitis, urticaria, inflammatory bowel diseases, and arthritis. An effective amount of kininogenase inhibiting compounds with or without a physiologically acceptable carrier or diluent can be used solely or in combination with other therapeutic agents.

The compounds inhibit the activity of kallikreins assessed with chromogenic substrates according to known procedures. The anti-inflammatory actions of the present compounds can for example be studied by their inhibition of allergen-induced exudative inflammatory processes in airway mucosa or gut mucosa.

#### Pharmaceutical preparations

The compounds of the invention will normally be administered orally, rectally, dermally, nasally, tracheally, bronchially, parenterally or via inhalation route, in the form of pharmaceutical preparations comprising the active ingredient either as a free base or a pharmaceutical acceptable non-toxic organic or inorganic acid addition salt, e.g. the hydrochloride, hydrobromide, sulphate, hydrosulphate, nitrate, lactate, acetate, citrate, benzoate, succinate, tartrate, trifluoroacetate and the like in a pharmaceutically acceptable dosage form. Depending upon

the disorder and patient to be treated and the route of administration, the compositions may be administered at varying dos s.

5 The dosage form may be a solid, semisolid or liquid preparation prepared by per se known techniques. Usually the active substance will constitute between 0.1 and 99 % by weight of the preparation, more specifically between 0.1 and 50 % by weight for  
10 preparations intended for parenteral administration and between 0.2 and 75 % by weight for preparations suitable for oral administration.

Suitable daily doses of the compounds of the invention  
15 in therapeutical treatment of humans are about 0.001-100 mg/kg body weight at peroral administration and 0.001-50 mg/kg body weight at parenteral administration.

20

#### Preparation

A further objective of the invention is the mode of preparation of the compounds. The compounds of  
25 Formula I and V may be prepared by processes comprise coupling of an N-terminally protected dipeptide or aminoacid, when a N-terminally amino acid is used a second aminoacid is added afterwards using standard methods to a compound

30



35 wherein n is an integer 0, 1, 2, 3 or 4, X is B or B-D where B is as defin d in formula I and D is as defin d in f rmula V as such or having the guanidino r amidino




nitrogens either mono or diprotected with an amine protecting group such as a benzyloxycarbonyl-, tert-butyloxycarbonyl- or p-toluenesulphonyl- group or X is a group transferable into B followed by removal of the  
 5 protectory group(s) or deprotection of the N-terminal nitrogen followed by alkylation of the N-terminal nitrogen and if desired deprotection by known methods and if desired forming a physiologically acceptable salt, and in those cases where the reaction results in  
 10 a mixture of stereoisomers, these are optionally separated by standard chromatographic or re-crystallisation techniques, and if desired a single stereoisomer is isolated.


15 In more detail the compounds of Formula I or V may be prepared by either of the following methods:

Method Ia

20 Coupling of an N-terminally protected dipeptide, selected from A<sup>1</sup> and A<sup>2</sup> in Formulas I or V and prepared by standard peptide coupling, with a compound



using standard peptide coupling, shown in the formula




wherein n is as defined in Formula I  $W^1$  is an N-terminal amino protecting group such as tert-butyloxy carbonyl and benzyloxy carbonyl and  $Q^1$  is  $-C(NH)-NH_2$ ,  $-C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH_2$ ,  $-NH-C(NH)-NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$ , where  $W^2$  is an amine protecting group such as tert-butyloxy carbonyl or benzyloxy carbonyl, or  $Q^1$  is  $-CN$ ,  $-CO-NH_2$  or  $-CS-NH_2$ , where the group is subsequently transferred into a amidino group (e.g giving  $Q^1 = -C(NH)-NH_2$ ) by methods known in the art or  $Q^1$  is  $NH_2$  or  $NH-W^2$ , where  $W^2$  is as defined above, where the amino group is subsequently transferred into a guanidino group (giving  $Q^1 = -NH-C(NH)-NH_2$ ), after deprotection of the  $W^2$ -group when  $Q^1$  is  $-NH-W^2$  ( $W^2$  in this case must be orthogonal to  $W^1$ ), by methods known in the art.


The final compounds can be made in any of the following ways, depending on the nature of the  $Q^1$ - group used: Removal of the protecting group(s) (when  $Q^1 = -C(NH)-NH_2$ ,  $-C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH_2$ ,  $-NH-C(NH)-NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$ ), or a selective deprotection of the  $W^1$ - group (e.g when  $Q^1 = -C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$  ( $W^2$  in this case must be orthogonal to  $W^1$ ) followed by alkylation of the N-terminal nitrogen by methods known in the art and if desired deprotection by known methods.

#### Method Ib

Coupling of an N-terminally protected amino acid, selected from  $A^2$  in Formulas I or V and prepared by standard methods, with a compound of formula



using standard peptide coupling, shown in the formula




20 wherein  $n$ ,  $W^1$ , and  $Q^1$  are as defined above followed by deprotection of the  $W^1$ -group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method Ia. The synthesis to the final peptides is then continued according to Method Ia.

25

#### Method IIa

30 Coupling of an N-terminally protected dipeptide, selected from  $A^1$  and  $A^2$  in Formulas I or V and prepared by standard peptide coupling, with a compound




using standard peptide coupling, shown in the formula



5

10



15


wherein  $n$  is as defined in Formula I,  $W^1$  is an N-  
 teminal amino protecting group such as tertbutyloxy  
 carbonyl and benzyloxy carbonyl and and  $Q^1$  is  $-C(NH)-$   
 $NH_2$ ,  $-C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH_2$ ,  $-NH-$   
 $C(NH)-NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$ ,  
 20 where  $W^2$  is an amine protecting group such as tert-  
 butyloxy carbonyl or benzyloxy carbonyl, or  $Q^1$  is -  
 $CN$ ,  $-CO-NH_2$  or  $-CS-NH_2$ , where the group is subsequently  
 transferred into a amidino group (e.g giving  $Q^1 = -$   
 $C(NH)-NH_2$ ) by methods known in the art or  $Q^1$  is  $NH_2$  or  
 $NH-W^2$ , where  $W^2$  is as defined above, where the amino  
 group is subsequently transferred into a guanidino  
 group (giving  $Q^1 = -NH-C(NH)-NH_2$ ), after deprotection of  
 the  $W^2$ -group when  $Q^1$  is  $-NH-W^2$  ( $W^2$  in this case must be  
 30 orthogonal to  $W^1$ ), by methods known in the art.

The final compounds can be made in any of the following  
 ways, depending on the nature of the  $Q^1$ - group used:  
 Removal of the protecting group(s) (when  $Q^1 = -C(NH)-NH_2$ ,  
 $-C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH_2$ ,  $-NH-C(NH)-$   
 $NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$ ), or a  
 selective deprotection of the  $W^1$ - group (e.g when  $Q^1 = -$   
 35



C(NW<sup>2</sup>)-NH-W<sup>2</sup>, -C(NH)-NH-W<sup>2</sup>, -NH-C(NH)-NH-W<sup>2</sup>, -N(W<sup>2</sup>)-C(NH)-NH-W<sup>2</sup> or -NH-C(NW<sup>2</sup>)-NH-W<sup>2</sup> (W<sup>2</sup> in this case must be orthogonal to W<sup>1</sup>) followed by alkylation of the N-terminal nitrogen by methods known in the art and if  
 5 desired deprotection by known methods.

#### Method IIb

Coupling of an N-terminally protected amino acid,  
 10 selected from A<sup>2</sup> in Formulas I or V and prepared by standard methods, with a compound of formula




using standard peptide coupling, shown in the formula



wherein n, W<sup>1</sup> and Q<sup>1</sup> are as defined above followed by deprotection of the W<sup>1</sup>-group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method IIa. The  
 35 synthesis to the final peptides is then continued according to Method IIa.

Method IIIa

Coupling of an N-terminally protected dipeptide,  
 selected from A<sup>1</sup> and A<sup>2</sup> in Formulas I or V and prepared  
 5 by standard peptide coupling, with a compound




10


using standard peptide coupling, shown in the formula



15



20



25


wherein n is as defined in Formula I and r is 0.1 when  
 X<sup>1</sup>, X<sup>2</sup> and X<sup>4</sup> are CH<sub>2</sub> or r is 0 when X<sup>2</sup> and X<sup>4</sup> are CH<sub>2</sub>  
 and X<sup>1</sup> is absent, W<sup>1</sup> is an N-terminal amino protecting  
 group such as tert-butyloxy carbonyl and benzyloxy  
 30 carbonyl and Q<sup>2</sup> is -C(NH)-NH<sub>2</sub>, -C(NW<sup>2</sup>)-NH-W<sup>2</sup>, or -  
 C(NH)-NH-W<sup>2</sup>, where W<sup>2</sup> is an amine protecting group such  
 as tert-butyloxy carbonyl or benzyloxy carbonyl, or Q<sup>2</sup>  
 is equal to W<sup>2</sup> where the amino group, after  
 deprotection of the W<sup>2</sup> group (W<sup>2</sup> in this case must be  
 35 orthogonal to W<sup>1</sup>), is subsequently transferred into a  
 guanidin group using an unprotected, N-protected or  
 N,N'-diprotected guanidation reagent by methods known

in the art (giving  $Q^2 = -C(NH)-NH_2$ ,  $-C(NW^2)-NH-W^2$  or  $-C(NH)-NH-W^2$ ).



The final compounds can be made in any of the following ways, depending on the nature of the  $Q^2$ - group used:  
 Removal of the protecting group(s) (when  $Q^2 = -C(NH)-NH_2$ ,  $-C(NW^2)-NH-W^2$  or  $-C(NH)-NH-W^2$ ), or a selective deprotection of the  $W^1$ - group (e.g when  $Q^2 = -C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $W^2$  in this case must be orthogonal to  $W^1$ ) followed by alkylation of the N-terminal nitrogen by methods known in the art and if desired deprotection known methods.

#### Method IIIB

Coupling of an N-terminally protected amino acid, selected from  $A^2$  in Formulas I or V and prepared by standard methods, with a compound of formula



using standard peptide coupling, shown in the formula




wherein  $n$ ,  $r$ ,  $X^1$ ,  $X^2$  and  $X^4$ ,  $W^1$ , and  $Q^2$  are as defined



above followed by deprotection of the  $W^1$ -group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method IIIa. The synthesis to the final peptides is then continued according to Method IIIa.

#### Method IVa

Coupling of an N-terminally protected dipeptide, selected from  $A^1$  and  $A^2$  in Formulas I or V and prepared by standard peptide coupling, with a compound



using standard peptide coupling, shown in the formula




wherein  $n$  is as defined in Formula I,  $W^1$  is an N-terminal amino protecting group such as tert-butyloxy carbonyl or benzyloxy carbonyl and  $W^3$  is H or an amino protecting group such as aryl sulfonyl, benzyl xy carbonyl or tert-butyloxy carbonyl. The final compounds





can be made in any of the following ways: Removal of the protecting group(s), or a selective deprotection of the  $W^1$ -group ( $W^1$  must be orthogonal to  $W^3$ ) followed by alkylation of the N-terminal nitrogen and if desired deprotection.

#### Method IVb

Coupling of an N-terminally protected amino acid, selected from  $A^2$  in Formulas I or V and prepared by standard methods, with a compound of formula



using standard peptide coupling, shown in the formula



wherein  $n$ ,  $W^1$ , and  $W^3$  are as defined above followed by deprotection of the  $W^1$ -group ( $W^1$  must be orthogonal to  $W^3$ ) and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method IVa. The synthesis to the final peptides is then continued according to Method IVa.

**DETAILED DESCRIPTION OF THE INVENTION**

The following description is illustrative of aspects of the invention.

5

**EXPERIMENTAL PART****General experimental Procedures.**

- 10 Mass spectra were recorded on a Finnigan MAT TSQ 700 triple quadrupole mass spectrometer equipped with an electrospray interface.

- 15 The  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR measurements were performed on BRUKER AC-P 300 and BRUKER AM 500 spectrometers, the former operating at a  $^1\text{H}$  frequency of 500.14 MHz and a  $^{13}\text{C}$  frequency of 125.76 MHz and the latter at  $^1\text{H}$  and  $^{13}\text{C}$  frequency of 300.13 MHz and 75.46 MHz respectively.

- 20 The samples were about 10-50 mg dissolved in 0.6 ml of either of the following solvents ;  $\text{CDCl}_3$  (isotopic purity > 99.8%),  $\text{CD}_3\text{OD}$  (isotopic purity > 99.95%),  $\text{D}_2\text{O}$  (isotopic purity > 99.98%) or  $\text{DMSO}-d_6$  (isotopic purity > 99.8%). All solvents were purchased from Dr. Glaser  
25 AG, Basel.

- The  $^1\text{H}$  and  $^{13}\text{C}$  chemical shift values in  $\text{CDCl}_3$  and  $\text{CD}_3\text{OD}$  are relative to tetramethylsilane as an external standard. The  $^1\text{H}$  chemical shifts in  $\text{D}_2\text{O}$  are relative to  
30 the sodium salt of 3-(trimethylsilyl)- $d_4$ -propanoic acid and the  $^{13}\text{C}$  chemical shifts in  $\text{D}_2\text{O}$  are referenced relative to 1,4-dioxane (67.3 ppm), both as external standard. Calibrating with an external standard may in some cases cause minor shift differences compared to an  
35 internal standard, however, the difference in  $^1\text{H}$  chemical shift is less than 0.02 ppm and in  $^{13}\text{C}$  less than 0.1 ppm.

The  $^1\text{H}$  NMR spectrum of peptide sequences containing a prolin or a "proline like" residue frequently exhibits two sets of resonances. This corresponds to the existence of two contributing conformers with respect to the rotation around the amide bond, where proline is the N-part of the amide bond. The conformers are named *cis* and *trans*. In our compounds the sequences (R)Cha-Aze-, (R)Cha-Pro- and (R)Cha-Pic- often give rise to a *cis-trans* equilibrium with one conformer as the preponderant conformer (>90%). In those cases only the  $^1\text{H}$  chemical shifts of the major rotamer is reported. Only in the cases where the signals of the minor rotamer are clearly resolved they are reported in the NMR documentation. The same criterium is valid for the NH-signals in  $\text{CDCl}_3$ , only in the cases where the signals are clearly resolved they are reported in the NMR-documentation. This implies that the number of protons reported for some of the intermediates are less than the number of protons expected from the chemical formula.

Thin-Layer Chromatography was carried out on commercial Merck Silicagel 60F<sub>254</sub> coated glass or aluminium plates. Visualisation was by a combination of UV-light, followed by spraying with a solution prepared by mixing 372 ml of EtOH(95%), 13.8 ml of concentrated  $\text{H}_2\text{SO}_4$ , 4.2 ml of concentrated acetic acid and 10.2 ml of p-methoxy benzaldehyde or phosphomolybdic acid reagent (5-10 w.t % in EtOH(95%)) and heating.

Flash chromatography was carried out on Merck Silica gel 60 (40-63 mm, 230-400 mesh) under pressure of air.

Reversed phase high-performance liquid chromatography (in the Examples referred to as RPLC) was performed on a Waters M-590 instrument quipped with three reverse phase Kromasil 100,C8 columns (Eka-Nobel) having

different dimensions for analytical (4.6 mm x 250 mm), semipreparative (1" x 250 mm) and preparative (2" x 500 mm) chromatography detecting at 226 nm.

- 5 Freeze-drying was done on a Leybold-Heraeus, model Lyovac GT 2, apparatus.

Preparation of starting materials

10

**Boc-(R)Pgl-OH**

Prepared in the same way as described for Boc-(R)Cha-OH (vide infra) from H-(R)Pgl-OH.

15

**Boc-(R)Cha-OH**

- 20 To a solution of H-(R)Cha-OH, 21.55 g (125.8 mmol), in 130 ml 1 M NaOH and 65 ml THF was added 30 g (137.5 mmol) of (Boc)<sub>2</sub>O and the mixture was stirred for 4.5 h at room temperature. The THF was evaporated and an additional 150 ml of water was added. The alkaline aqueous phase was washed twice with EtOAc, then acidified with 2 M KHSO<sub>4</sub> and extracted with 3 x 150 ml of EtOAc. The combined organic phase was washed with water, brine and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent afforded 30.9 g (90.5 %) of the title compound as a white solid.

30

**Boc-(R)Hop-OH**

Prepared by the same procedure as described for Boc-(R)Cha-OH starting from H-(R)Hop-OH.

35

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 1.45 (s, 9H), 2.00 (m, 1H), 2.22 (m, 1H), 2.75 (bt, 2H), 4.36 (bs, 1H), 5.05 (bs, 1H), 7.15-7.33 (m, 5H).

**4-(tert-butyloxycarbonylaminomethyl) pyridine**

To a solution of 10.81 g (100 mmol) 4-aminomethyl pyridine in 100 ml THF was added 24 g (110 mmol) Boc<sub>2</sub>O dissolved in 70 ml THF at 10°C for 20 minutes. The solution was allowed to reach room temperature and stirred for 4 h (a precipitate was formed during the reaction and the slurry became red). The solvent was removed and the residue was dissolved in EtOAc and filtered through silica gel. Evaporation of the solvent gave the title compound as a red oil which crystallized on standing. The crude product was used without further purification.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 1.45 (s, 9H), 4.32 (d, 2H), 5.05 (bs, 1H (NH)), 7.2 (d, 2H), 8.55 (d, 2H).

**4-aminomethyl-1-(N-benzyloxycarbonylamidino)-benzene (H-Pab(Z))****(i) 4-cyanobenzyl azide**

A solution of 20.23 g (0.31 mol) sodium azide in 50 ml water was added to 49.15 g (251 mmol) 4-cyanobenzyl bromide in 200 ml DMF at ambient temperature. An exothermic reaction took place and after 1.5 h the reaction mixture was diluted with 200 ml toluene (caution: In order to avoid separation of potentially explosive azide compounds it is advisable to add the toluene to the reaction mixture before addition of the water) and 500 ml water. The aqueous phase was extracted with an additional 2x50 ml toluene. The combined organic extracts were washed with 2x50 ml water and brine and finally dried (MgSO<sub>4</sub>) and filtered. The solution was used as such in the next step.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>); δ 4.4 (s, 2H), 7.4 (d, 2H), 7.7

(d, 2H).

(ii) 4-amidino benzyl azide

5 Hydrogen chloride was bubbled into a mixture of 250 ml absolute ethanol and the solution from step (i) (approximatly 200 ml) above at - 5°C until saturation. Storage at 8°C for 24 h and evaporation of most of the solvent followed by precipitation by addition of  
10 anhydrous ether gave white crystals which were isolated by filtration and dissolved in 1.8 l of alcoholic ammonia. After 48 h most of the solvent was removed and 200 ml 3.75 M NaOH solution was added whereupon 4-amidino benzyl azide precipitated as colourless  
15 crystals. The crystals were isolated by filtration. At this point the yield of 4-amidino benzyl azide was 22.5 g (total 51%).

Ethylimidatobenzyl azide hydrochloride:

20

<sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD); δ 1.6 (t, 3H), 4.5 (s, 2H), 4.65 (q, 2H), 4.8 (br s, 2H), 7.6 (d, 2H), 8.1 (d, 2H)

4-amidino benzyl azide:

25

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>); δ 4.3 (s, 2H), 5.7 (br s, 3H), 7.3 (d, 2H), 7.6 (d, 2H).

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>): amidine carbon: δ 165.5.

30

(iii) 4-(benzyloxycarbonylamidino) benzyl azide

The crystals from (ii) above were dissolved in 500 ml methylene chloride and the resulting solution was dried  
35 (K<sub>2</sub>CO<sub>3</sub>), filtered and 27 ml (194 mmol) triethyl amine was added. 25 ml Benzyl chloroformate was slowly added to the stirred solution while the reaction mixture was

cooled in an ice bath. After 30 minutes an additional 2 ml benzyl chloroformate was added and stirring was continued for another 30 minutes. Subsequently, water was added and the aqueous phase was adjusted to pH 7 with 2M HCl. The organic phase was dried (MgSO<sub>4</sub>) and the solvent was removed in vacuo. 4-(benzyloxycarbonylamidino) benzyl azide was finally isolated as colorless crystals from ether/methylene chloride/hexane.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>); δ 4.4 (s, 2H), 5.3 (s, 2H), 6.3-7.0 (br s, 1H), 7.3-7.4 (m, 5H), 7.5 (d, 2H), 7.9 (d, 2H), 9.3-9.6 (br s, 1H).

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>): amidine carbon: δ 167.5.

(iv) 4-aminomethyl-1-(N-benzyloxycarbonylamidino)-benzene (H-Pab(Z))

26.3 g (100 mmol) triphenylphosphine was added at room temperature to the 4-(benzyloxycarbonylamidino) benzyl azide from (iii) above dissolved in 160 ml THF. After 16 h an additional 6.6 g (25 mmol) triphenylphosphine was added and the solution was allowed to stand for 4 h before removal of the solvent in vacuo. The residue was dissolved in methylene chloride and extracted with 2M HCl. The aqueous phase was washed with methylene chloride and ether and was subsequently made alkaline with 3.75M sodium hydroxide solution. Extraction with methylene chloride followed by drying (K<sub>2</sub>CO<sub>3</sub>) and removal of the solvent in vacuo gave 20 g (The total yield starting from cyanobenzyl bromide is 28%) of a yellow oil which solidified on standing.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>); δ 1.2-2.2 (br s, 2H), 3.8 (s, 2H), 5.2 (s, 2H), 7.2-7.35 (m, 5H), 7.4 (d, 2H), 7.8

(d, 2H), 9.1-9.6 (br s, 1H).

$^{13}\text{C}$ -NMR (125 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  
 $\delta$  164.6 and 168.17.

5

**H-Pig(3)<sub>2</sub>**

(i) 4-(tert-butyloxycarbonyl-aminomethyl) piperidine

10 To a solution of 17.7 g 4-tert-  
butyloxycarbonylaminomethyl pyridine in 125 ml MeOH was  
added 2 g of 5 % Rh/ $\text{Al}_2\text{O}_3$  and the mixture was  
hydrogenated at 0.34 MPa over night.  $^1\text{H}$ -NMR showed that  
the hydrogenation was incomplete. Therefore, the  
15 catalyst was filtered off and the solvent removed in  
vacuo and the residue was dissolved in 100 ml acetic  
acid, 2 g of 5 % Rh/ $\text{Al}_2\text{O}_3$  was added and the mixture was  
hydrogenated for 4 days at 0.34 MPa. The catalyst was  
filtered off and most of the acetic acid was removed in  
20 vacuo. After addition of 50 ml water to the residue  
the mixture was made alkaline with 5 M NaOH and the  
water phase was extracted with 1 x 200 + 1 x 100 ml  
 $\text{CH}_2\text{Cl}_2$ . The combined organic phase was washed with 25 ml  
water and dried ( $\text{MgSO}_4$ ). Evaporation of the solvent  
25 gave a 17.2 g of a brownish oil which was dissolved in  
50 ml of diethyl ether. Addition of 200 ml pentane gave  
a precipitate which was filtered off to give 7.7 g of a  
brown powder. Evaporation of the mother liquor gave 7 g  
of a white oil. The brown powder was dissolved in 100  
30 ml EtOAc and the organic phase was washed with 1 x 50  
ml + 1 x 25 ml 1 M  $\text{KHSO}_4$ . The combined acidic phase was  
made alkaline with 2 M NaOH and extracted with 1 x 200  
+ 1 x 75 ml EtOAc. The combined organic phase was dried  
and evaporated to give 5.2 g of the title compound as a  
35 white powder.

Treatment of the white oil obtained from the mother



liquor above in the same way afforded an additional 3.4 g of the product. Total yield 40 %.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>, mixture of two rotamers, 3:1):  
5 major rotamer: δ 1.11 (dq, 2H), 1.44 (s, 9H), 1.49-1.60 (m, 1H), 1.63-1.70 (m, 2H), 2.58 (dt, 2H), 2.93-3.03 (m, 2H), 3.07 (m, 2H), 4.75 (bs, 1H (NH)).

10 Resolved signals arising from the minor rotamer appear at δ 1.21 (dq) and 1.91 (dt).

(ii) Boc-Pig(Z)<sub>2</sub>

To a solution of 2 g (9.33 mmol) 4-(tert-  
15 butyloxycarbonyl-aminomethyl) piperidine in 60 ml CH<sub>3</sub>CN was added 3.34 g (9.33 mmol) of N,N'-(dibenzoyloxy-carbonyl)methylisothiurea and the mixture was stirred at 60°C for 22 h. The solvent was evaporated and the  
20 residue was dissolved in EtOAc. The organic phase was washed with 2 x 20 ml 1 M KHSO<sub>4</sub>, 1 x 20 ml water, 1 x 20 ml brine and dried (MgSO<sub>4</sub>). Evaporation of the solvent followed by flash chromatography using  
petroleum ether/EtOAc (1/1) as eluent afforded 2.43 g  
(50%) of the desired product.

25 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): Some signals, especially in the piperidine ring, are selectively broadend due to an intramolecular exchange process. This is especially pronounced for the 2- and 6-CH<sub>2</sub> groups of the  
30 piperidine ring, which exhibit a broad peak ranging from 3.7 to 4.5 ppm.

δ 1.19-1.31 (m, 2H), 1.43 (s, 9H), 1.63-1.80 (m, 3H),  
2.66-3.05 (m, 4H), 3.7-4.5 (bs, 2H), 4.65 (bt, 1H(NH)),  
5.13 (s, 4H), 7.2-7.4 (m, 10H), 10.5 (bs, 1H(NH)).

35

(iii) H-Pig(Z)<sub>2</sub>

A solution of 163 mg (0.31 mmol) Boc-Pig(Z)<sub>2</sub> in 5 ml EtOAc saturated with HCl(g) was stirred at ambient temperature for 3 h and 20 minutes. The solvent was evaporated and the residue was dissolved in 30 ml CH<sub>2</sub>Cl<sub>2</sub>. The organic phase was washed with 5 ml 2 M NaOH, 1 x 5 ml water, 1 x 5 ml brine and dried (MgSO<sub>4</sub>). Evaporation of the solvent afforded 100 mg (76 %) of the title compound.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): Some signals, especially in the piperidin ring, are selectively broadened due to an intramolecular exchange process. This is especially pronounced for the 2- and 6-CH<sub>2</sub> groups of the piperidine ring, which exhibit a broad peak ranging from 3.7 to 4.5 ppm.  
δ 1.18-1.37 (m, 2H), 1.46-1.63 (m, 1H), 1.68-1.83 (m, 2H), 2.57 (d, 2H), 2.86-3.03 (m, 2H), 3.7-4.5 (bs, 2H), 5.13 (s, 4H), 7.2-7.4 (m, 10H).

4-aminomethyl-1-(N-benzyloxy carbonylamidino)-cyclohexane (H-Pac(Z) x 2HCl).

(i) N-[N-4-(benzyloxycarbonyl)amidino benzyl] tert-butyl carbamate

1.466 g (6.7 mmol) (Boc)<sub>2</sub>O was added to a stirred ice cold solution of 1.81 g (6.4 mmol) 4-(benzyloxycarbonyl)amidino benzyl amine and 1 ml (7.1 mmol) triethyl amine in 25 ml methylene chloride. After 20 minutes more methylene chloride was added and the mixture was washed with 5% acetic acid and 10% sodium carbonate solution. Drying (magnesium sulphate) and removal of the solvent in vacuo left a residue which could be crystallised from methylene chloride/hexane. The yield was 1.66 g (68%).

(ii) N-[N-4-amidin benzyl]tert-butyl carbamate

A mixture of 1.60 g (4.2 mmol) N-[4-(benzyloxycarbonyl)amidino benzyl] tert-butyl carbamate, 5 ml acetic acid, and 160 mg 10% palladium on charcoal in 50 ml ethanol was stirred in an atmosphere of hydrogen for 2 h. The catalyst was removed by filtration through celite and the solvent was removed in vacuo to give the acetate of the title compound in quantitative yield.

(iii) N-[4-amidino cyclohexyl methyl]tert-butyl carbamate

17 mmol of the acetate of N-[4-amidino benzyl]tert-butyl carbamate was hydrogenated in 100 ml methanol in the presence of 863 mg 5% rhodium on alumina at 3.4 MPa for 20 h. The catalyst was removed by filtration and the solvent was removed in vacuo. The residue was dissolved in water and the solution was made basic with sodium hydroxide. Subsequent extraction with methylene chloride, drying of the combined organic phases (potassium carbonate) and removal of the solvent in vacuo gave 3.8 g (87%) of the title compound.

(iv) N-[N-4-(benzyloxycarbonyl)amidino cyclohexyl methyl] tert-butyl carbamate

1.25 ml (8.8 mmol) benzyl chloroformate was added at 0°C to a stirred solution of 2.04 g (8 mmol) N-[4-amidino cyclohexyl]tert-butyl carbamate, 1.23 ml (8.8 mmol) triethyl amine, and 197 mg DMAP in 40 ml methylene chloride. After 10 minutes the reaction mixture was diluted with methylene chloride and extracted with water, dilute acetic acid, and sodium hydrogen carbonate solution. The organic phase was applied on a column of silica and subsequent elution with methylene chloride containing increasing amounts of ethyl acetate yielded 2.49 g (80%) of the title

compound.

(v) 4-aminomethyl-1-(N-benzyloxy carbonylamidino)-  
cyclohexane (H-Pac(Z) x 2HCl).

5

Hydrogen chloride was passed through a solution of 2 g  
(5.1 mmol) N-[4-(benzyloxycarbonyl)amidino cyclohexyl  
methyl]tert-butyl carbamate in 40 ml ethyl acetate.  
After 10 minutes methanol was added and upon removal of  
10 some of the solvent in vacuo the dihydrochloride of  
title compound crystallised.

4-aminomethyl-1-(N-benzyloxy carbonylamidino)  
piperidine (H-Pig(Z) x HCl)

15

(i) 4-(N-tert-butyloxycarbonylaminomethyl)-1-(N-  
benzyloxycarbon ylamidino) piperidine (Boc-Pig(Z))

7.8 g (36.4 mmole) of 4-(N-tert-  
20 butyloxycarbonylaminomethyl) piperidine and 8.98 g (40  
mmole) of N-benzyloxycarbonyl-S-methylisothiourea was  
mixed in 25 mL ethanol. The mixture was heated at  
60-70°C for six hours and left at roomtemperature for  
two days. The solvent was evaporated and the residue  
25 was dissolved in CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed  
twice with 0.3 M KHSO<sub>4</sub> and once with brine. The  
combined organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and  
evaporated. The crude product was purified by flash  
chromatography using a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH  
30 (100/0, 97/3, 95/5, 90/10) as eluent to yield 5.22 g  
(37%) of the title product.

(ii) H-Pig(Z) x HCl (4-aminomethyl-1-(N-benzyloxy  
carbonylamidino) piperidine

35

5.22 g (13.5 mmole) f Boc-Pig(Z) was dissolv d in 100  
mL ethyl acetate saturated with HCl(g). The mixture was

allowed to stand for one hour and then evaporated. The residue was dissolved in water and washed with a mixture of diethylether and ethyl acetate. The water layer was freeze-dried to yield 4.0 g (91%) of the title compound.

$^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ , 300 MHz):  $\delta$  1.40-1.60 (m, 2H), 2.05 (bd, 2H), 2.19 (m, 1H), 3.07(d, 2H), 3.34(bt, 2H), 4.08 (bd, 2H), 5.40 (s, 2H), 7.5-7.63 (m, 5H)

10

MS  $m/z$  291 ( $\text{M}^+ + 1$ )

4-Aminoethyl-1-benzyloxycarbonylamidino piperidine (H-Rig(Z))

15

(i) 1-Benzyloxycarbonylamidino-4-hydroxyethyl piperidine

A mixture of 6.2 g (0.028 mol) of 4-hydroxyethyl piperidine and 3.6 g (0.028 mol) of N-benzyloxycarbonyl-S-methyl isothiurea in 50 ml of acetonitrile was refluxed overnight. Evaporation and flash chromatography on silica gel with ethyl acetate gave 3.5 g (41%) of the title compound.

25

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.1-1.85 (m, 7 H), 2.83 (bt, 2 H), 4.70 (bt, 2 H), 4.18 (bd, 2 H), 5.12 (s, 2 H), 6.9- 7.2 (m, 2 H), 7.2-7.5 (m, 5 H).

(ii) 1-Benzyloxycarbonylamidino-4-mesyloxyethyl piperidine

To an ice cooled solution of 3.50 g (0.0115 mol) of 1-benzyloxy-carbonylamidino-4-hydroxyethyl piperidine, 1.15 g (0.0115 mol) of triethyl amine in 40 ml of methylene chloride and 10 ml of tetrahydrofuran was added dropwise 1.30 g (0.115 mol) of mesyl chloride.

35

The reaction mixture was allowed to stir for 1 h. The mixture was poured into water and the organic layer was kept. The aqueous layer was extracted with methylene chloride and the combined organic layers were washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. The product was used without further purification in the next step. Yield: 4.4 g (100%).

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.15-1.3 (m, 2 H), 1.65-1.8 (m, 5 H), 2.84 (bt, 2 H), 3.01 (s, 3 H), 4.20 (bd, 2 H), 4.27 (t, 2 H), 5.12 (s, 2 H), 7.1-7.5 (m, 7 H).

(iii) 4-Azidoethyl-1-benzyloxycarbonylamidino piperidine

In 100 ml of dimethylformamide was dissolved 4.4 g (0.0115 mol) of crude 1-benzyloxycarbonylamidino-4-mesyloxyethyl piperidine and 4.5 g (0.069 mol) of sodium azide was added. The mixture was heated at  $100^\circ\text{C}$  for 2.5 h. It was then poured into water and extracted with ethyl acetate three times. The combined organic phase was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. The residue was flash chromatographed on silica gel using ethyl acetate/heptane 1/1 as eluent. Yield: 3.0 g (79%).

$^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.20 (dq, 2H), 1.5-1.8 (m, 5 H), 2.85 (dt, 2 H), 3.35 (t, 2 H), 4.22 (bd, 2 H), 5.13 (s, 2 H), 6.9-7.2 (b, 2 H), 7.2-7.45 (m, 5 H).

(iv) 4-Aminoethyl-1-benzyloxycarbonylamidino piperidine (H-Rig(Z))

To 30 ml of water was added 0.40 g of 10% Pd/C. Sodium borohydride, 1.0 g (0.031 mol), was dissolved in 30 ml of water and was added carefully to the stirred and ice-cooled slurry of Pd/C and water. 4-Azidoethyl-1-

benzyl xycarbonylamidino piperidine, 2.9 g (8.8 mmol), was dissolved in 80 ml of tetrahydrofuran and this solution was added dropwise to the ice-cooled aqueous slurry above. After 4 h of stirring at room temperature the mixture was ice-cooled again and 30 ml of 2 M HCl was added. The mixture was filtered through celite and the celite was rinsed with additional water. The tetrahydrofuran was evaporated and the aqueous phase was washed with ethyl acetate. The aqueous phase was made alkaline with 2 M NaOH and extracted with methylene chloride three times. The combined organic phase was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. The product was used in the following step without further purification.

$^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.1-1.5 (m, 6 H), 1.55-1.65 (m, 1H), 1.73 (bd, 2 H), 2.72 (b, 2 H), 2.81 (dt, 2 H), 4.20 (bd, 2 H), 5.12 (s, 2 H), 6.9-7.2 (b, 2 H), 7.2-7.5 (m, 5 H).

(3RS)-1-(N-benzyloxycarbonylamidino)-3-aminomethyl pyrrolidine (H-(R,S)Nig(Z))

(i) (3RS)-3-hydroxymethyl pyrrolidine

16.4 g (0.0857 mole) (3RS)-1-benzyl-3-hydroxymethyl pyrrolidine (See H-(R,S)Hig(Z) (i) vide supra) was mixed with 1.6 g Pd/C (10%), 5 ml water and 150 ml ethanol and the mixture was hydrogenated at 0.26 MPa over night. After filtration through hyflo and evaporation of the solvent the  $^1\text{H-NMR}$  showed that the reaction was not completed. Continued hydrogenation at 0.26 MPa over 1.6 g Pd/C (10%) in 5 ml water/150 ml ethanol for three days completed the reduction. Filtration through hyflo and evaporation of the solvent gave the product in a quantitative yield.

(ii) (3RS)-1-(N-benzyloxycarbonylamidino)-3-hydroxymethyl pyrrolidine

1.01 g (0.01 mole) (3RS)-3-hydroxymethyl pyrrolidine  
5 and 2.29 g (0.011 mole) N-benzyloxycarbonyl-O-methylisourea was dissolved (the amine not very soluble) in toluene and heated to 60°C for three hours followed by stirring at room temperature over night. The mixture was evaporated and the <sup>1</sup>H-NMR showed that  
10 the reaction was not completed. The mixture was therefore dissolved in 15 ml acetonitrile and heated to 60°C for three hours followed by stirring at room temperature over night. The solvent was evaporated and the mixture was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, washed once with  
15 water, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH 95/5 as eluent to yield 0.70 g (25%) of the product.

20 MS m/z 278 (M<sup>+</sup>+1)

(iii) (3RS)-1-(N-benzyloxycarbonylamidino)-3-mesyloxymethyl pyrrolidine

25 0.7 g (2.53 mmole) (3RS)-1-(N-benzyloxycarbonylamidino)-3-hydroxymethyl pyrrolidine and 0.70 ml (5.05 mmole) triethylamine was dissolved in 15 ml diethyl ether/CH<sub>2</sub>Cl<sub>2</sub> 1/1 and the mixture was cooled to 0°C. 0.25 ml (3.29 mmole) methanesulphonyl  
30 chloride in 3 ml diethyl ether was slowly added and the reaction mixture was stirred at 0°C for three hours. The solvent was evaporated and the residue was dissolved in ethyl acetate and extracted with a 0.3 M KHSO<sub>4</sub>-  
35 solution. The water layer was washed once with CH<sub>2</sub>Cl<sub>2</sub>. The water layer was made neutral with 10 M NaOH-solution and extracted twice with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and



evaporated to yield 0.450 g (50%) of the title compound.

(iv) (3RS)-1-(N-benzyloxycarbonylamidino)-3-azidomethyl pyrrolidine

0.450 g (1.27 mmole) (3RS)-1-(N-benzyloxycarbonylamidino)-3-mesyloxymethyl pyrrolidine and 0.124 g (1.9 mmole) of sodium azide were dissolved in 10 ml dimethylformamide and heated to 60°C for four hours followed by stirring at room temperature over night. Water was added and the mixture was extracted twice with toluene/ethyl acetate 2/1. The combined organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. The crude product was purified by flash chromatography using  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  95/5 as eluent to yield 0.262 g (68%) of the product.

MS m/z 303 ( $\text{M}^+ + 1$ )

(v) (3RS)-1-(N-benzyloxycarbonylamidino)-3-aminomethyl pyrrolidine (H-(R,S)Nig(Z))

32 mg Pd/C (10%) and 2.6 ml  $\text{H}_2\text{O}$  was mixed and a gentle stream of nitrogen was passed. 98 mg  $\text{NaBH}_4$  in 2.6 ml  $\text{H}_2\text{O}$  was added followed by a slow addition of 262 mg (0.87 mmole) (3RS)-1-(N-benzyloxycarbonylamidino)-3-mesyloxymethyl pyrrolidine dissolved in 7 ml MeOH. The mixture was allowed to stand for one hour. 5 ml 1M HCl was added and the mixture was filtered through hyflo. The organic solvent was evaporated under reduced pressure and the remaining water layer was washed once with ethyl acetate, made alkaline with NaOH-solution and extracted several times with  $\text{CH}_2\text{Cl}_2$ . The combined organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to yield 130 mg (54%) of the product.

MS m/z 277 ( $\text{M}^+ + 1$ )

(3RS)-1-(N-benzyloxycarbonylamidin)-3-aminoethyl  
pyrrolidine (E-(R,S)Hig(2))

5 (i) (3RS)-1-benzyl-3-hydroxymethyl pyrrolidine

25.2 g (0.1063 mole) (3RS)-1-benzyl-2-oxo-4-methoxycarbonyl pyrrolidine was slowly added to a  
10 slurry of 6.22 g lithium aluminium hydride in 160 ml  
diethyl ether under an argon-atmosphere. The mixture  
was stirred over night and then heated to reflux for  
one hour. The reaction mixture was cooled to room  
temperature and 0.2 g of  $\text{Na}_2\text{SO}_4 \times 10 \text{ H}_2\text{O}$  was added  
15 followed by a slow addition of, in that order, 6 ml  
water, 18 ml 3.75 M NaOH-solution and 6 ml water. The  
slurry was dried from excess of water with  
 $\text{Na}_2\text{SO}_4$ /cellite, filtered and evaporated to give (20.3 g)  
of the product.

20  $^1\text{H}$ -NMR ( $\text{CDCl}_3$ , 300 MHz):  $\delta$  1.64-1.77 (m, 1H), 1.93-2.07  
(m, 1H), 2.27-2.40 (m, 2H), 2.51 (dd, 1H), 2.62 (dd,  
1H), 2.82 (m, 1H), 3.52 (dd, 1H), 3.59 (s, 2H), 3.67  
(dd, 1H), 7.15-7.40 (m, 5H)

25 (ii) (3RS)-1-benzyl-3-chloromethyl pyrrolidine

To a refluxing solution of 15.3 g (0.08 mole) (3RS)-1-benzyl-3-hydroxymethyl pyrrolidine in 220 ml  $\text{CHCl}_3$  was  
30 slowly added a solution of 330 ml thionyl chloride in  
60 ml  $\text{CHCl}_3$ , and the reflux was continued for one hour.  
The mixture was evaporated and the residue was  
dissolved in water.

The water layer was washed with ethyl acetate and then  
made alkaline with 0.2 M NaOH-solution. The water layer  
35 was extracted three times with ethyl acetate and the  
combined organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and  
evaporated to give the product in a quantitative yield

(16.8 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 1.55 (m, 1H), 2.05 (m, 1H),  
2.38 (dd, 1H), 2.48-2.64 (m, 3H; thereof 2.58 (t,  
5 2H))), 2.73 (dd, 1H), 3.51 (d, 2 H), 3.60 (s, 2H), 7.2-  
7.4 (m, 5H)

(iii) (3RS)-1-benzyl-3-cyanomethyl pyrrolidine

10 16.8 g (0.08 mole) (3RS)-1-benzyl-3-chloromethyl  
pyrrolidine and 5.88 g (0.12 mole) of sodium cyanide  
was dissolved in 250 ml dimethyl sulfoxide. The  
mixture was stirred at 60°C for two days and at room  
temperature for one week. Water was added and the  
15 mixture was extracted three times with ethyl acetate.  
The combined organic layer was washed with brine, dried  
(Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to yield 14.7 g (92%)  
of the product.

20 <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz): δ 1.55 (m, 1H), 2.13 (m, 1H),  
2.35 (dd, 1H), 2.42 (t, 2H), 2.44-2.59 (m, 2H), 2.65  
(m, 1H), 2.73 (dd, 1H), 3.61 (s, 2H), 7.2-7.4 (m, 5H)

(iv) (3RS)-1-benzyl-3-aminoethyl pyrrolidine

25 14.7 g (0.0734 mole) (3RS)-1-benzyl-3-cyanomethyl  
pyrrolidine dissolved in 220 ml diethyl ether was  
slowly added to a slurry of 2.94 g of lithium aluminium  
hydride in 74 ml diethyl ether under an argon  
30 atmosphere. The mixture was stirred over night, and 6 ml  
water, 18 ml 3.75 M NaOH-solution and 6 ml water was  
added to the mixture. The slurry was dried from excess  
of water with Na<sub>2</sub>SO<sub>4</sub>/cellite, filtered by suction and  
evaporated to yield 14.84 g (99%) of the product.

35 <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 1.41 (m, 1H), 1.51 (q, 2H),  
1.90-2.10 (m, 2H; thereof 2.05 (dd, 1H)), 2.18 (m,

1H), 2.43 (m, 1H), 2.55-2.73 (m, 3H), 2.80 (apparent t, 1H), 3.58 (apparent d, 2H), 7.15-7.4 (m, 5H)

(v) (3RS)-1-benzyl-3-(N-tert-butylloxycarbonylaminoethyl) pyrrolidine

To a mixture of 14.84 g (0.0726 mole) (3RS)-1-benzyl-3-aminoethyl pyrrolidine, 72.6 ml 1M NaOH-solution, 76 ml water and 145 ml THF was added 17.44 g (0.08 mole) di-tert-butyl dicarbonate and the mixture was stirred over night. The solution was concentrated and extracted three times with ethyl acetate. The combined organic layer was washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated. The crude product was purified by flash chromatography using a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95/5, 90/10) as eluent to yield 14.69 g (80%) of the product.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 1.25-1.65 (m, 12H; thereof 1.40 (s, 9H)), 1.90-2.25 (m, 3H), 2.46 (m, 1H), 2.67 (m, 1H), 2.80 (apparent t, 1H), 3.09 (m, 2H), 3.59 (s, 2H), 4.60 (bs, NH), 7.15-7.35 (m, 5H)

(vi) (3RS)-3-(N-tert-butylloxycarbonylaminoethyl) pyrrolidine

3.1 g (0.01 mol) (3RS)-1-benzyl-3-(N-tert-butylloxycarbonylaminoethyl) pyrrolidine was hydrogenated at 0.28 MPa over 0.6 g of Pearlman's catalyst (Pd(OH)<sub>2</sub>) in 40 ml ethanol (95%) over night. After filtration of the catalyst through cellite and evaporation of the solvent <sup>1</sup>H-NMR showed that the reaction was not completed. Therefore 0.6 g of Pearlman's catalyst was added in 40 ml ethanol (95%) once more and the mixture was treated under H<sub>2</sub>-atmosphere at 0.28 MPa over night. Filtration through cellite and evaporation of the solvent gave the product

in a quantitativ yield (2.18 g).

MS m/z 214 ( $M^+$ )

- 5 (vii) (3RS)-1-(N-benzyloxycarbonylamidino)-3-aminoethyl  
pyrrolidine (H-(R,S)Hig(Z))

2.18 g (0.0102 mmole) (3RS)-3-(N-tert-  
butyloxycarbonylaminoethyl) pyrrolidine and 2.81 g  
10 (0.0125 mole) N-benzyloxycarbonyl-S-methylisothioure  
was dissolved in 30 ml toluene and heated to 60-70°C  
for eight hours followed by stirring at room  
temperature for one day. 0.3 M  $KHSO_4$ -solution was added  
and the water layer was washed with a mixture of the  
15 toluene and ethyl acetate and left for 2 days under  
which time the Boc group was removed. The acidic water  
phase was made alkaline and extracted four times with  
 $CH_2Cl_2$ . The combined organic layer was dried ( $Na_2SO_4$ ),  
filtered and evaporated to yield 2.0 g (51%) of the  
20 title compound.

$^1H$ -NMR ( $CDCl_3$ , 330 K, 300 MHz):  $\delta$  1.45-1.7 (m, 3H), 2.07  
(m, 1H), 2.26 (m, 1H), 2.74 (t, 2H), 3.00 (apparent t,  
1H), 3.33 (apparent q, 1H), 3.45-3.80 (m, 2H), 5.12 (s,  
25 2H), 6.72 (bs, 2 NH), 7.15-7.45 (m, 5H)

(4RS)-1,3-diaza-2-tosylimino-4-aminoethyl cyclohexane  
(H-(R,S)Itp(Ts))

- 30 (i) (4RS)-1,3-diaza-2-tosylimino-4-carboxy cyclohexane

Prepared using the same method as described in Journal  
of Org. Chem., p. 46, 1971.

- 35 (ii) (4RS)-1,3-diaza-2-tosylimino-4-hydroxymethyl  
cycl hexane

12.9 g (345 mmol)  $\text{LiAlH}_4$  was carefully added to a cold slurry (ice bath temperature) of 9.9 g (33 mmol) of (4RS)-1,3-diaza-2-tosylimino-4-carboxy cyclohexane in 330 mL dry THF. The reaction was stirred at room temperature over night. The reaction mixture was worked up according to Fieser & Fieser, e.g by adding 12.9 g water, 38.7 g 3.75 M NaOH, 12.9 g water,  $\text{Na}_2\text{SO}_4$ ,  $\text{CH}_2\text{Cl}_2$  and celite to the mixture, and filtered. Evaporation of the solvent gave 7.0 g (75%) of the desired product.

MS m/z 284 ( $\text{M}^+ + 1$ )

(iii) (4RS)-1,3-diaza-2-tosylimino-4-mesyloxymethyl cyclohexane

2.9 mL  $\text{MsCl}$  (37.1 mmol) was added carefully to a cold (ice bath temperature) slurry of 7.0 g (24.7 mmol) of (4RS)-1,3-diaza-2-tosylimino-4-hydroxymethyl cyclohexane in 6.9 mL (49.4 mmol) triethylamine and 125 mL  $\text{CH}_2\text{Cl}_2$ . Water was added after 1h 15 min and the organic phase was separated, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to give the title compound in quantitative yield.

MS m/z 362 ( $\text{M}^+ + 1$ )

(iv) (4RS)-1,3-diaza-2-tosylimino-4-cyanomethyl cyclohexane

8.9 g (24.7 mmol) of (4RS)-1,3-diaza-2-tosylimino-4-mesyloxymethyl cyclohexane and 1.3 g (27.2 mmol) NaCN was dissolved in 75 mL DMSO. After stirring at 40°C for 60 hours an additional amount of 0.31 g (6 mmol) NaCN was added and the solution was stirred at 65°C for three hours. 150 mL water was added and crystals precipitated out of the solution. They were filtered off and dried to give 5.4 g (75%) of the desired

product.

MS m/z 293 ( $M^+ + 1$ )

5 (4RS)-1,3-diaza-2-tosylimino-4-aminoethyl cyclohexane  
(H-(R,S)Itp(Ts))

935 mg  $LiAlH_4$  was added carefully to a cooled (ice bath  
temperature) slurry of 2.4 g (8.2 mmol) of (4RS)-1,3-  
10 diaza-2-tosylimino-4-cyanomethyl cyclohexane in 90 mL  
THF. After stirring for 2 hours 1 g  $H_2O$ , 3 g 3.75M  
NaOH, 1 g  $H_2O$ ,  $Na_2SO_4$ , celite and  $CH_2Cl_2$  was added. The  
mixture was filtered and the solvent removed in vacuo  
to give 2.2 g (90%) of the desired product.

15

$^1H$  NMR (500 MHz, MeOD);  $\delta$  1.52-1.71 (m, 3H), 1.88-1.96  
(m, 1H), 2.37 (s, 3H), 2.64-2.73 (m, 2H), 3.2-3.4 (m,  
2H, partially overlapping with the solvent signal),  
3.44-3.53 (m, 1H), 7.28 (d, 2H), 7.71 (d, 2H)

20

(4S)-1,3-diaza-2-tosylimino-4-aminoethyl cyclohexane  
(H-(S)Itp(Ts))

Prepared in the same way as described for H-  
25 (R,S)Itp(Ts) starting from optically pure 2,4-  
diaminobutyric acid.

$^1H$  NMR (300.13 MHz,  $CDCl_3$ );  $\delta$  0.97-1.15 (s broad, 1H),  
1.48-1.69 (m, 3H), 1.84-1.95 (m, 1H), 2.37 (s, 3H),  
30 2.68-2.82 (m, 1H), 2.86-2.98 (m, 1H), 3.22-3.44 (m,  
2H), 3.45-3.58 (m, 1H), 7.19 (d, 2H), 7.72 (d, 2H)

$^{13}C$  NMR (300.13 MHz,  $CDCl_3$ );  $\delta$  guanidinecarbon 154.05

35 H-Aze-OEt x HCl

Prepared in the same way as described for H-Pic-OEt x

HCl from H-Aze-OH (vide infra).

**H-Aze-OMe x HCl**

- 5     Prepared according to the procedure described by Seebach D. et. al. in Liebigs Ann. Chem., p. 687, 1990.

**H-Pab(Z) x HCl**

- 10    Prepared by adding 1 mole equivalent of 5 M HCl in isopropanol to a solution of crude H-Pab(Z) in EtOH (about 1 g/10 ml) where upon H-Pab(Z) x HCl immediately precipitates out of the solution. After filtration the precipitate was washed 2 times with cold EtOH and dried  
15    to give the title compound in almost quantitative yield.

**H-Pic-OEt x HCl**

- 20    L-Pipecolinic acid, 4.0 g (0.031 mol), was slurried in 100 ml of abs. ethanol and HCl (g) was carefully bubbled through until a clear solution was obtained. It was cooled in an ice bath and 17 ml of thionyl chloride was added dropwise over 15 min. The ice bath was  
25    removed and the mixture was refluxed for 2.5 h. The solvent was evaporated and the product was obtained as its hydrochloride salt in a quantitative yield.

- 30    <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 1.33 (t, 3H), 1.8-2.1 (m, 5H), 2.3-2.5 (m, 1H), 3.1-3.3 (m, 1H), 3.5-3.7 (m, 1H), 4.14 (dd, 1H), 4.44 (q, 2H).

**H-(R,S)betaPic-OMe x HCl**

35

A mixture of 2.0 g (15.5 mmol) nipecotic acid in 8 ml methanol was cooled in an ice-bath and 2.76 g (23.2



mmol) thionyl chloride was added. The mixture was stirred at room temperature for 20 hours. The solvent was evaporated and the residue was dissolved in a small amount of methanol, diethylether was added and H-(R,S)betaPic-OMe x HCl precipitated as white crystals. The crystals 2.57 g (92%) were isolated by filtration.

#### Boc-(R)Cgl-OH

Boc-(R)-Pgl-OH, 32.6 g (0.13 mol), was dissolved in 300 ml of methanol and 5 g of Rh/Al<sub>2</sub>O<sub>3</sub> was added. The solution was hydrogenated at 5.2 to 2.8 MPa for 3 days. After filtration and evaporation of the solvent NMR showed the presence of about 25 % of the methyl ester of the title compound. The crude material was dissolved in 500 ml of THF and 300 ml of water and 20 g of LiOH were added. The mixture was stirred overnight and the THF was evaporated. The remaining water phase was acidified with KHSO<sub>4</sub> and extracted three times with ethyl acetate. The combined organic layer was washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give 28.3 g (83 %) of the desired product.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 0.9-1.7 (m, 20H), 4.0-4.2 (m, 1H), 5.2 (d, 1H).

#### Boc-(R)Cgl-OSu

To an ice-cold solution of 2.01 g (7.81 mmol) of Boc-(R)Cgl-OH and 1.83 g (15.6 mmol) of HOSu in 25 ml of CH<sub>3</sub>CN was added 1.69 g (8.2 mmol) of DCC and the reaction was allowed to reach room temperature. After stirring for 3 days the precipitated DCU was filtered off and the solvent evaporated. The residue was dissolved in EtOAc and the organic phase was washed with H<sub>2</sub>O, KHSO<sub>4</sub>, NaHCO<sub>3</sub>, brine and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent gave the title compound in

quantitative yield.

**Boc-(R)Cha-OSu**

5    Boc-(R)Cha-OH (1 eq.), HOSu (1.1 eq) and DCC or CME-CDI  
     (1.1 eq) were dissolved in acetonitrile (about 2.5  
     ml/mmol acid) and stirred at room temperature over  
     night. The precipitate formed during the reaction was  
10    filtered off, the solvent evaporated and the product  
     dried in vacuo. (When CME-CDI was used in the reaction  
     the residue, after evaporation of the CH<sub>3</sub>CN, was  
     dissolved in EtOAc and the organic phase washed with  
     water and dried). Evaporation of the solvent gave the  
     title compound.

15

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>, 2 rotamers ca: 1:1 ratio) \_  
0.85-1.1 (m, 2H), 1.1-1.48 (m, 4H), 1.5-1.98 (m, 16H;  
thereof 1.55 (bs, 9H)), 2.82 (bs, 4H), 4.72 (bs, 1H,  
major rotamer), 4.85 (bs, 1H, minor).

20

**Boc-(R)Hoc-OH**

Boc-(R)Hoc-OH (See above), 3.2 g (11.46 mmol) was  
dissolved in methanol (75 ml). Rhodium on activated  
25    aluminium oxide (Rh/Al<sub>2</sub>O<sub>3</sub>), 0,5 g was added and the  
     mixture was stirred under a hydrogen atmosphere at 0.41  
     MPa for 18 h. The catalyst was filtered off through  
     hyflo and the solvent evaporated giving the product in  
     almost quantitative yield.

30

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.90 (m, 2H), 1.08-1.33 (m,  
6H), 1.43 (s, 9H), 1.60-1.74 (m, 6H), 1.88 (bs, 1H),  
4.27 (bs, 1H).

35

**Boc-(R)Hoc-OSu**

Prepared in the same way as described for Boc-(R)Cha-

OSu from Boc-(R)Hoc-OH.

**Boc-(R)Pro(3-(S)Ph)-OH**

- 5     Prepared according to the method described by J.Y.L  
Chung et al in Journal of Organic Chemistry, No 1, pp.  
270-275, 1990.

**Boc-(R)Cgl-Aze-OH**

10

(i) Boc-(R)Cgl-Aze-OMe

- To a stirred mixture of 3.86 g (15 mmol) Boc-(R)Cgl-OH,  
2.27 g (15 mmol) H-Aze-OMe x HCl and 2.75 g (22.5 mmol)  
15     DMAP in 40 mL CH<sub>3</sub>CN at 5°C was added 3.16 g (16.5  
mmol) EDC. The reaction mixture was stirred at room  
temperature for 48h. The solvent was evaporated and the  
residue was dissolved in 150 ml EtOAc and 20 ml H<sub>2</sub>O.  
The separated organic layer was washed with 2 x 20 ml  
20     0.5 M KHSO<sub>4</sub>, 2 x 10 ml NaHCO<sub>3</sub>(saturated), 1 x 10 ml H<sub>2</sub>O,  
1 x 10 ml brine and dried (MgSO<sub>4</sub>). Evaporation of the  
solvent gave 4.91 g (92 %) of the title compound which  
was used without further purification in the next step.

- 25     <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, 0.1 g/ml): major rotamer, 0.83-  
1.35 (m, 5H), 1.38 (s, 9H), 1.47-1.84 (m, 6H), 2.18-  
2.27 (m, 1H), 2.50-2.62 (m, 1H), 3.72 (s, 3H), 3.94-  
4.06 (m, 1H), 4.07-4.15 (m, 1H), 4.39-4.47 (m, 1H),  
4.68 (dd, J=9.1, J=5.1, 1H), 5.09 (d, J=9.2, 1H).  
30     Resolved peaks from minor rotamer, 2.27-2.35 (m, 1H),  
3.77 (s, 3H), 3.80-3.87 (m, 1H), 3.88-3.95 (m, 1H),  
4.92 (d, J=9.2, 1H), 5.21 (dd, J=9.1, J=5, 1H).

(ii) Boc-(R)Cgl-Aze-OH

35

The hydrolysis of Boc-(R)Cgl-Aze-OMe was carried out  
according to the procedure described for Boc-(R)Cha-

Pic-OEt (vide infra). The product was crystallized from EtOH/acetone/water (1/1/3.95) yield 80 %.

5  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.85-1.3 (m, 5H), 1.40 (s, 9H), 1.5-1.9 (m, 6H), 1.95-2.2 (m, 2H), 3.92 (m, 1H), 4.09 (m, 1H), 4.35 (m, 1H), 4.95 (m, 1H), 5.16 (bd, 1H).

10 **Boc-(R)Cgl-Pic-OH**

(i) Boc-(R)Cgl-Pic-OMe

15 Pivaloyl chloride (1.0 ml, 8.1 mmol) was added to a solution of Boc-(R)Cgl-OH (2.086 g, 8.1 mmol) and triethyl amine (1.13 ml, 8.1 mmol) in toluene (25 ml) and DMF (5 ml). A mixture of H-Pic-OMe x HCl (1.46 g, 8.1 mmol) and triethyl amine (1.13 ml, 8.1 mmol) in DMF (20 ml) was subsequently added at ice bath temperature. The reaction mixture was slowly allowed to warm up to  
20 room temperature and after 24 h it was diluted with water and extracted with toluene. After washing with 0.3 M  $\text{KHSO}_4$ , 10%  $\text{Na}_2\text{CO}_3$  and brine the solvent was removed in vacuo to give 2.52 g (81%) of colorless oil which was used without further purification.

25  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ , 2 rotamers, 5:1 ratio)  $\delta$  0.8-1.8 (m, 25H), 2.25 (d, 1H), 2.75 (t, 1H, minor rotamer), 3.3 (t, 1H), 3.7 (s, 3H), 3.85 (d, 1H), 4.3 (t, 1H, minor rotamer), 4.5-4.6 (m, 1H), 5.25 (d, 1H), 5.30 (d, 1H).  
30

(ii) Boc-(R)Cgl-Pic-OH

35 Prepared according to the procedure for hydrolysis of Boc-(R)Cha-Pic-OEt (vide infra) using the product from (i) above. The product was crystallized from diisopropyl ether and hexane.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>, 2 rotamers, 5:1 ratio) δ 0.8-1.8 (m, 25H), 2.3 (d, 1H), 2.8 (t, 1H, minor rotamer), 3.3 (t, 1H), 3.9 (d, 1H), 4.4 (t, 1H, minor), 4.5-4.6 (m, 1H), 5.1 (s, 1H, minor rotamer), 5.3 (d, 1H), 5.40 (d, 1H).

#### Boc-(R)Cgl-Pro-OH

3.59 g (31.24 mmol) of L-proline was mixed with 20 ml water and 1.18 g (29.7 mmol) of sodium hydroxide. 2.8 g (7.8 mmol) of Boc-(R)Cgl-OSu in 10 ml DMF was added to the mixture. Because of solubility problem an additional 30 ml of DMF was added and the reaction mixture was stirred for three days. The solvent was evaporated and water was added. The water phase was washed with ethyl acetate, acidified with 0.3 M KHSO<sub>4</sub>-solution and extracted three times with ethyl acetate. The organic phase was washed once with water and once with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to yield 2.3 g (83 %) of the product.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 0.89-2.17 (m, 23H), 2.37 (m, 1H), 3.55 (q, 1H), 3.90 (bs, 1H), 4.28 (t, 1H), 4.52 (bs, 1H), 5.22 (bs, 1H (NH)).

#### Boc-(R)Cha-Aze-OH

Prepared in the same way as described for Boc-(R)Cha-Pic-OH starting from Boc-(R)Cha-OH and H-Aze-OEt x HCl (vide infra).

#### Boc-(R)Cha-Pro-OH

H-(S)Pro-OH (680 mmol) was dissolved in 0.87 M sodium hydroxide (750 ml). Boc-(R)Cha-OSu (170 mmol) dissolved in DMF (375 ml) was added dropwise during 20 min. The reaction mixture was stirred at room temperature for

20 h. The mixture was acidified (2M KHSO<sub>4</sub>) and extracted three times with ethyl acetate. The organic layers were combined and washed three times with water and once with brine. After drying over sodium sulphate and evaporation of the solvent, the syrupy oil was dissolved in diethyl ether, the solvent evaporated and finally the product dried in vacuo to yield Boc-(R)Cha-Pro-OH as a white powder in almost quantitative yield.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>, mixture of two rotamers 9:1) δ 0.8-1.05 (m, 2H), 1.05-1.55 (m, 15H; thereof 1.5 (bs, 9H)), 1.55-1.8 (m, 5H), 1.8-2.15 (m, 3H), 2.47 (m, 1H), 3.48 (m, 1H), 3.89 (m, 1H), 4.55 (m, 2H), 5.06 (m, 1H); Resolved signals from the minor rotamer appears at δ 2.27 (m), 3.58 (m), 4.33 (m), 5.0 (m).

#### Boc-(Me) (R)Cha-Pro-OSu

##### (i) Boc-(Me) (R)Cha-Pro-OH

Prepared in the same way as described above for Boc-(R)Cha-Pro-OH starting from Boc-(Me) (R)Cha-OSu and H-Pro-OH.

##### (ii) Boc-(Me) (R)Cha-Pro-OSu

Prepared in the same way as described for Boc-(R)Cha-OSu starting from Boc-(Me) (R)Cha-Pro-OH.

#### Boc-(R)Cha-Pic-OH

##### (ia) Boc-(R)Cha-Pic-OEt

Boc-(R)Cha-OH, 6.3 g (0.023 mol), was dissolved in 150 ml of CH<sub>2</sub>Cl<sub>2</sub>. The solution was cooled in an ice bath and 6.3 g (0.047 mol) of N-hydroxybenzotriazole and 11.2 g (0.0265 mol) of CME-CDI were added. The ice bath

was removed after 15 min and the reaction mixture was stirred for 4 h at room temperature. The solvent was evaporated and the residue dissolved in 150 ml of DMF and cooled in an ice bath. H-Pic-OEt $\cdot$ HCl, 4.1 g (0.021 mol) was added and the pH adjusted to approximately 9 by addition of N-methylmorpholine. The ice bath was removed after 15 min and the reaction mixture was stirred for 3 days. The solvent was evaporated and the residue was dissolved in ethyl acetate and washed with dilute KHSO<sub>4</sub> (aq), NaHCO<sub>3</sub> (aq) and water. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give 7.7 g (89 %) of Boc-(R)Cha-Pic-OEt which was used without further purification.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>, 2 rotamers, 3:1 ratio)  $\delta$  0.7-1.0 (m, 2H), 1.1-1.9 (m, 29H; thereof 1.28 (t, 3H)), 1.45 (bs, 9H), 2.01 (bd, 1H, major rotamer), 2.31 (bd, 1H), 2.88 (bt, 1H, minor), 3.30 (bt, 1H, major), 3.80 (bd, 1H, major), 4.15-4.3 (m, 2H), 4.5-4.7 (m, 2H, minor), 4.77 (bq, 1H, major), 4.90 (bd, 1H, minor), 5.28 (bd, 1H, major), 5.33 (bd, 1H, major).

(1b) Boc-(R)Cha-Pic-OMe

400  $\mu$ l (3.23 mmol) of pivaloyl chloride was added to a stirred mixture of 875 mg (3.22 mmol) Boc-(R)Cha-OH and 450  $\mu$ l (3.23 mmol) triethyl amine in 10 ml toluene and 2 ml DMF. A mixture of 596 mg (3.32 mmol) methyl (S)-pipercolate hydrochloride and 463  $\mu$ l (3.32 mmol) triethyl amine in 5 ml DMF was added to the resulting slurry after 45 minutes. After 2 h 100  $\mu$ l (0.72 mmol) triethyl amine was added and stirring was continued for another 18 h. Water and toluene was added to the reaction mixture and the organic phase was washed with 0.3 M KHSO<sub>4</sub>, 10% Na<sub>2</sub>CO<sub>3</sub> and brine. Drying (MgSO<sub>4</sub>) and removal of the solvent in vacuo gave 1.16 g of the title compound.

(ii) Boc-(R)Cha-Pic-OH

Boc-(R)Cha-Pic-OEt, 5.6 g (0.014 mol), was mixed with 100 ml of THF, 100 ml of water and 7 g of LiOH. The mixture was stirred at room temperature overnight. The THF was evaporated and the aqueous solution was acidified with KHSO<sub>4</sub> (aq) and extracted three times with ethyl acetate. The combined organic phase was washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to give 4.9 g (94 %) of Boc-(R)Cha-Pic-OH which was used without further purification. The compound can be crystallized from diisopropyl ether/hexane.

The methyl ester formed in procedure (ib) above can be hydrolysed using the same procedure as described for the ethyl ester in (ii).

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>, 2 rotamers, 3.5:1 ratio) δ 0.8-1.1 (m, 2H), 1.1-2.1 (m, 27H; thereof 1.43 (s, 9H, major rotamer), 1.46 (s, 9H, minor)), 2.33 (bd, 1H), 2.80 (bt, 1H, minor), 3.33 (bt, 1H, major), 3.85 (bd, 1H, major), 4.57 (bd, 1H, minor), 4.68 (m, 1H, minor), 4.77 (bq, 1H, major), 5.03 (bs, 1H, minor), 5.33 (bd, 1H, major), 5.56 (m, 1H, major).

25

Boc-(R)Cha-(R,S)betaPic-OH

(i) Boc-(R)Cha-(R,S)betaPic-OMe

Pivaloyl chloride 0.9 ml (7.3 mmol) was added to a solution of 2.0 g (7.3 mmol) Boc-(R)Cha-OH and 0.81 ml (7.3 mmol) 4-N-methyl morpholin in 20 ml acetonitrile. After stirring for 1 h and 30 minutes 1.3 g (7.3 mmol) H-(R,S)betaPic-OMe x HCl and 1.62 ml (14.6 mmol) 4-N-methyl morpholine was added and the reaction mixture was stirred for 24 h. The solvent was evaporated and the residu was dissolved in toluene and some

35



diethyleter. After washing with 0.3 M  $\text{KHSO}_4$  and  $\text{KHCO}_3$ -  
solutions, and drying with  $\text{Na}_2\text{SO}_4$  the solvent was removed  
in vacuo. Flash chromatography using heptane/ethyl  
acetate (7/3) as eluent gave 2.4 g (83%) of the desired  
5 product.

(ii) Boc-(R)Cha-(R,S)betaPic-OH

At room temperature 2.35 g (5.9 mmol) of Boc-(R)Cha-  
10 (R,S)betaPic-OMe was dissolved in 35 ml THF and 2.1 g  
of LiOH in 35 ml water was added. After stirring for 5  
h the THF was removed in vacuo. The aqueous phase was  
acidified with 2M  $\text{KHSO}_4$  and extracted with ethyl  
acetate, dried over  $\text{Na}_2\text{SO}_4$  and evaporated to give 2.0 g  
15 (89%) of the product.

Boc-(R)Cha-Val-OH

(i) Boc-(R)Cha-Val-OMe

20

3.1 ml (25 mmol) pivaloyl chloride was added at ambient  
temperature to a stirred mixture of 6.75 g (25 mmol)  
Boc-(R)Cha-OH and 3.5 ml (25 mmol) triethyl amine in 50  
ml DMF. After 3 hours 4.16 g (25 mmol) valine methyl  
25 ester hydrochloride in 50 ml DMF and 3.5 ml triethyl  
amine was added. After stirring over night, a few  
crystals of DMAP were added and the reaction mixture  
was heated to 50°C for 5 minutes. The solvent was  
removed in vacuo and ether and toluene was added to the  
30 residue. Washing with 0.3 M  $\text{KHSO}_4$  and 10%  $\text{Na}_2\text{CO}_3$   
followed by drying ( $\text{MgSO}_4$ ) and removal of the solvent  
in vacuo gave a residue which was subjected to flash  
chromatography using toluene/ethyl acetate as eluent.  
The yield of the title compound was 6.99 g (73%).

35

(ii) Boc-(R)Cha-Val-OH

A mixture of 8.73 g (23 mmol) Boc-(R)Cha-Val-OMe and 5.6 g (230 mmol) lithium hydroxide in 75 ml THF and 75 ml of water was stirred for 4 hours. The THF was removed in vacuo and the remaining solution was diluted with water and extracted with ether. Acidification with 2 M KHSO<sub>4</sub> and extraction with ethyl acetate followed by drying (MgSO<sub>4</sub>) and removal of the solvent in vacuo gave 8.15 g (96%) of the title compound.

10 Boc-(R)Hoc-Aze-OH

(i) Boc-(R)Hoc-Aze-OEt

At room temperature 1.0 g (3.5 mmol) Boc-(R)Hoc-OH and 0.95 g (7.0 mmol) HOBt was dissolved in 15 ml CH<sub>2</sub>Cl<sub>2</sub>. The solution was cooled in an ice bath and 0.77 g (4.0 mmol) of EDC was added. The ice bath was removed and the reaction mixture was stirred for 3 h at room temperature. The solvent was evaporated and the residue dissolved in 20 ml DMF. 0.58 g (3.5 mmol) H-(R)Aze-OH was added and the pH adjusted to approximately 9 by addition of N-methyl morpholin. The reaction mixture was stirred for one day. The reaction mixture was partitioned between water and toluene. The organic phase was separated and washed with 0.3 M KHSO<sub>4</sub>, diluted KHCO<sub>3</sub>, brine, dried with NaSO<sub>4</sub> and evaporated. Flash chromatography (1% EtOH in CH<sub>2</sub>Cl<sub>2</sub> and heptane: EtOAc) gave 0.35 g (25%) of the desired product.

30 (ii) Boc-(R)Hoc-Aze-OH

At room temperature 0.65 g (1.6 mmol) Boc-(R)Hoc-Aze-OEt was dissolved in 10 ml THF and 0.59 g of LiOH in 10 ml water was added. After stirring for 24 hours 2 M KHSO<sub>4</sub> was added and the THF was removed in vacuo. The aqueous phase was then made acidic with more 2M KHSO<sub>4</sub> and extracted with ethyl acetate, dried over Na<sub>2</sub>SO<sub>4</sub> and

evaporated to give 0.5 g (85%) of the title compound.

**Boc-(R)Hoc-Pro-OH**

- 5 Prepared in the same way as described for Boc-(R)Cha-Pro-OH from Boc-(R)Hoc-OSu.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.80-0.94 (m, 2H), 1.05-1.36 (m, 7H), 1.36-1.48 (bs, 9H), 1.48-1.78 (m, 7H), 1.98-  
10 2.14 (m, 2H), 2.34 (m, 1H), 3.48 (m, 1H), 3.85 (m, 1H), 4.43 (m, 1H), 4.52 (bd, 1H), 5.26 (bd, 1H), signals of a minor rotamer appears at: δ 1.92, 2.25, 3.58, 4.20 and 4.93.

15 **Boc-(R)Hoc-Pic-OH**

(i) Boc-(R)Hoc-Pic-OMe

- 20 Prepared the same way as described for Boc-(R)Cha-Pic-OEt (vide supra) from Boc-(R)Hoc-OH and H-Pic-OMe x HCl.

(ii) Boc-(R)Hoc-Pic-OH

- 25 Prepared in the same way as described for Boc-(R)Cha-Pic-OH (vide supra) from Boc-(R)Hoc-Pic-OMe.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.82-0.97 (m, 2H), 1.10-1.36 (m, 7H), 1.36-1.50 (bs, 9H), 1.50-1.82 (m, 11H), 2.35 (bd, 1H) 3.28 (bt. 1H), 3.85 (bd, 1H) 4.63 (m, 1H),  
30 5.33 (bs, 1H), 5.44 (bd, 1H), signals of a minor rotamer appear at: δ 1.88, 2.80, 4.25, 4.55 and 4.97.

**Boc-(R)Pro-Phe-OH**

35

(i) Boc-(R)Pro-Phe-OM

To a solution of 2.0 g (9.29 mmol) Boc-(R)Pro-OH and 0.94 g (9.29 mmol) triethyl amine in 70 ml toluene/DMF (5/2) was added 1.12 g (9.29 mmol) pivaloylchloride and the reaction was stirred for 30 minutes at room temperature. The reaction was cooled to 0°C and a mixture of 2.0 g (9.29 mmol) H-Phe-OMe and 0.94 g triethyl amine in 40 ml DMF was added and the reaction was stirred over night at room temperature. The reaction mixture was diluted with toluene and the organic phase was washed with 3 x 50 ml 0.3 M KHSO<sub>4</sub>, 1 x 50 ml water and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent gave the title compound in quantitative yield which was used in the next step without further purification.

(ii) Boc-(R)Pro-Phe-OH

A mixture of 4.0 g (10.6 mmol) Boc-(R)Pro-Phe-OMe and 8.93 g (21.3 mmol) LiOH x H<sub>2</sub>O in 140 ml water/THF (1/1) was stirred vigorously over night at room temperature. The THF was evaporated and the water phase was made acidic with 1 M KHSO<sub>4</sub> and extracted with 3 x 75 ml EtOAc. The combined organic phase was washed with water and dried (Na<sub>2</sub>SO<sub>4</sub>). Filtration and evaporation of the solvent gave a residue which was purified by crystallization from diisopropyl ether to give 2.329 g (60 %) of the title compound as a white crystalline solid.

Boc-(R)Pro(3-(S)Ph)-Pro-OH

(i) Boc-(R)Pro(3-(S)Ph)-Pro-OBn

To a mixture of 1.61 g Boc-(R)Pro(3-(S)Ph)-OH, 1.65 g H-Pro-OBn x HCl and 0.75 g HOBt in 11 mL DMF was added 0.84 mL NMM and 2.92 g CME-CDI at room temperature and the reaction mixture was stirred for three days. The

solvent was vap rat d and th residue was dissolved in 300 mL EtOAc. The organic phase was washed with 2 x 100 mL H<sub>2</sub>O, 2 x 100 mL 1 M KHSO<sub>4</sub>, 3 x 100 mL 1 M NaOH, 3 x 100 ml H<sub>2</sub>O and dried (MgSO<sub>4</sub>). Evaporation of the solvent gave 2.53 g of an oil which was purified by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (97/3) as eluent to give 2.11 g (88%) of the title compound.

(ii) Boc-(R)Pro(3-(S)Ph)-Pro-OH

10

0.94 g of Boc-(R)Pro(3-(S)Ph)-Pro-OBn was dissolved in 70 ml EtOH and hydrogenated over 0.42 g 5 % Pd/C for 3.5 hours. Filtration of the catalyst and evaporation of the solvent gave the title compound as white crystals in a quantitative yield.

15

Boc-(R)Tic-Pro-OH

Prepared according to the procedure described by P.D. Gesellchen and R.T. Shuman in EP-0,479,489-A2.

20

BnOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-Br

To a solution of p-TsOH x H-Gly-OBn (5 mmol) and triethyl amine (5 mmol) in 10 ml of CH<sub>2</sub>Cl<sub>2</sub> was added 2-bromoacetic acid (5 mmol) dissolved in 10 ml of CH<sub>2</sub>Cl<sub>2</sub> and dicyclohexyl carbodiimide (5 mmol). The mixture was stirred at room temperature over night and filtered. The organic phase was washed twice with 0.2 M KHSO<sub>4</sub>, 0.2 M NaOH, brine and dried. Evaporation and flash chromatography (CH<sub>2</sub>Cl<sub>2</sub>/MeOH, 95/5) gave a quantitative yield of the desired compound.

25

30

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 3.89 (s, 2H), 4.05-4.11 (d, 2H), 5.19 (s, 2H), 7.06 (bs, 1H), 7.3-7.4 (m, 5H).

35

**B c-(R)Cgl-I1 -OH**

Prepared in the same way as described for Boc-(R)Cgl-Pro-OH using H-Ile-OH, instead of H-Pro-OH, in 91 %  
5 yield.

**Boc-(R)Phe-Phe-OH****(i) Boc-(R)Phe-Phe-OMe**

10

Boc-(R)Phe-OH (18.8 mmol; purchased from Bachem  
Feinchemicalien AG), Phe-OMe (20.7 mmol) and  
4-dimethylaminopyridine (37.7 mmol) were dissolved in  
30 mL of acetonitrile. The solution was cooled to ice-  
15 water temperature and 1-(3-dimethylaminopropyl)-3-  
-ethylcarbodiimide hydrochloride (24.5 mmol) was added.  
The cooling bath was removed and the reaction mixture  
was stirred over night. The solvent was then removed  
under reduced pressure and the residue was dissolved in  
20 50 mL of ethylacetate. Extraction of the organic phase  
with 50 mL aliquats of 0.5 M potassiumhydrogensulfate,  
1 M sodiumbicarbonate and finally water followed by  
evaporation of the solvent yielded 7.5 g of Boc-(R)Phe-  
Phe-OMe (94%) which was used in the next step without  
25 further purification.

**(ii) Boc-(R)Phe-Phe-OH**

Boc-(R)Phe-Phe-OMe (16.4 mmol) was dissolved in 40 mL  
30 of tetrahydrofuran and lithiumhydroxide (32.8 mmol)  
dissolved in 20 mL of water was added rapidly. The  
reaction mixture was stirred for 3.5 h after which the  
solvent was removed under reduced pressure. The residue  
was dissolved in 50 mL of ethylacetate and extracted  
35 with 50 mL of 0.5 M potassiumsulfate followed by 50 mL  
of water. The solvent was removed under reduced  
pressure yielding 8.0 g of Boc-(R)Phe-Phe-OH (quant) as

an amorphous solid.  $^1\text{H}$  NMR (200 MHz,  $\text{d-CHCl}_3$ );  $\delta$  7.4-6.7 (m, 10H), 5.7-4.2 (m, 6H), 1.34 (s, 9H).

**HO-CH<sub>2</sub>-COOBn**

5

Prepared according to the procedure described by Lattes A. et al in Bull. Soc. Chim. France., Noll, pp 4018-23, 1971.

10 **Benzyl-2-(ortho-nitrobenzenesulfonyloxy)acetate (2-NO<sub>2</sub>)Ph-SO<sub>2</sub>-OCH<sub>2</sub>-COOBn**

1.66 g (10 mmol) benzylglykolate was dissolved in 25 ml CH<sub>2</sub>Cl<sub>2</sub> and 25 ml diethylether. The mixture was cooled to 0°C and 2.8 ml (20 mmol) triethylamin was added. While keeping the temperature at 0°C 2.44 g (11 mmol) ortho-nitrobenzenesulfonylchloride was added in small portions during 15 minutes. The slurry was stirred at 0°C for 50 minutes and then 20 ml water and 30 ml CH<sub>2</sub>Cl<sub>2</sub> were added. The phases were separated and the organic phase was washed with 20 ml 1 M HCl and 20 ml H<sub>2</sub>O, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated in vacuo to give 3.34 g of a residue that was subjected to flash chromatography, using heptan:EtOAc 2:1 as eluent to give 1.18 g (34 %) of the title compound.

$^1\text{H}$ -NMR (300MHz, CDCl<sub>3</sub>):  $\delta$  4.92 (s, 2H), 5.17 (s, 2H), 7.83 (m, 5H), 7.76 (m, 3H), 8.16(dd, 1H).

30 **Benzyl-2-(para-nitrobenzenesulfonyloxy)acetate (4-NO<sub>2</sub>)Ph-SO<sub>2</sub>-OCH<sub>2</sub>-COOBn**

Prepared according to the same procedure as described for Benzyl-2-(ortho-nitrobenzenesulfonyloxy)acetate above. The final compound was obtained in a crystalline form after evaporation of the solvent and pur enough to use without further purification (64 % yield).

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  4.79 (s, 2H), 5.13 (s, 2H), 7.2-7.4 (m, 5H), 8.10 (d, 2H), 8.30 (d, 2H).

**TfO-CH<sub>2</sub>COOMe**

5  
10.09 ml (60 mmol) trifluoromethanesulfonic anhydride dissolved in  $\text{CH}_2\text{Cl}_2$  was added dropwise to a mixture of 4.05 ml (50 mmol) methylglycolate and 4.04 ml (50 mmol) pyridine in  $\text{CH}_2\text{Cl}_2$  (totally 62.5 ml) at 0°C during 25  
10 minutes, and thereafter stirred at 0°C for 1 h. After washing with 0.3 M  $\text{KHSO}_4$  and saturated  $\text{Na}_2\text{CO}_3$ , drying ( $\text{Na}_2\text{SO}_4$ ) and filtration, evaporation of the solvent in vacuo gave 9.94 g (90 %) of the title compound.

15 **TfO-CH<sub>2</sub>COOEt**

Prepared in the same way as described for TfO-CH<sub>2</sub>COOMe starting with ethylglycolate.

20 **TfO-CH<sub>2</sub>COO<sup>n</sup>Bu**

Prepared in the same way as described for TfO-CH<sub>2</sub>COOMe starting with butylglycolate.

25 **TfO-CH<sub>2</sub>COOBn**

Prepared in the same way as described for TfO-CH<sub>2</sub>COOMe starting with HO-CH<sub>2</sub>COOBn

30 **TfO-CH<sub>2</sub>COO<sup>n</sup>Hex**

(i) HO-CH<sub>2</sub>COO<sup>n</sup>Hex

35 To 215 mg (2.82 mmol) glycolic acid in 12.8 ml  $\text{CH}_3\text{CN}$  was added 719 mg (3.39 mmol) 1-hexyl iodide and 429 mg (2.82 mmol) DBU. After stirring over night and reflux



for 4 h, the solvent was evaporated, ethylacetate and 1 M  $\text{KHSO}_4$  was added and the phases were separated. The organic layer was washed with brine, dried ( $\text{MgSO}_4$ ), filtered and evaporated in vacuo to give 333 mg (74 %) of the product.

(ii)  $\text{TfO-CH}_2\text{COO}^n\text{Hex}$

Prepared in the same way as described for  $\text{TfO-CH}_2\text{COOMe}$  starting with  $\text{HO-CH}_2\text{COO}^n\text{Hex}$ .

**H-Mig(2) (3-aminomethyl-1-(N-benzyloxycarbonylamidino) azetidine**

(i) 3-aminomethyl-1-benzhydryl azetidine was prepared according to the literature, see A.G. Anderson, Jr., and R. Lok, *J.Org.Chem.*, 37, 3953, 1972.

(ii) 3-(N-tert-butyloxycarbonylaminomethyl)-1-benzhydryl azetidine

To 3.50 g (13.9 mmol) of 3-aminomethyl-1-benzhydryl azetidine dissolved in 45 mL THF was added a solution of 0.56 g (13.9 mmol) NaOH in 45 mL  $\text{H}_2\text{O}$ . The reaction mixture was cooled to  $0^\circ\text{C}$  and 3.03 g (13.9 mmol) of di-tert-butyl dicarbonate was added. The cooling bath was removed after a few minutes and the mixture was stirred at room temperature over night. The THF was evaporated and the residue was extracted with 3x45 mL of diethyl ether. The combined organic layer was washed with brine, dried with  $\text{Na}_2\text{SO}_4$  and filtered. Evaporation of the solvent gave 4.6 g (94 %) of the title compound.

(iii) 3-(N-tert-butyloxycarbonylaminomethyl) azetidine

3.4 g (9.6 mmol) of 3-(N-tert-

butyloxycarbonylaminomethyl)-1-benzhydryl azetidine was dissolved in 170 mL MeOH and hydrogenated over 0.30 g Pd(OH)<sub>2</sub> at 5 MPa over night. The catalyst was filtered off and the solvent evaporated. The crude product was  
5 purified by flash chromatography using MeOH/CH<sub>2</sub>Cl<sub>2</sub>, 1/9, followed by MeOH (saturated with NH<sub>3</sub> (g))/CH<sub>2</sub>Cl<sub>2</sub>, 1/9, as eluent to yield 1.2 g (67 %) of the title compound.

10 (iv) 3-(N-tert-butyloxycarbonylaminomethyl)-1-(N-benzyloxycarbonylamidino) azetidine (Boc-Mig(Z))

0.9 g (4.8 mmol) of 3-(N-tert-butyloxycarbonylaminomethyl) azetidine and 1.3 g (6.3  
15 mmol) of N-benzyloxycarbonyl-O-methylisourea was mixed in 6.5 mL toluene and heated to 70°C for 72 h and then left at roomtemperature for another 72 h. Evaporation followed by flash chromatography using EtOAc followed by MeOH (saturated with NH<sub>3</sub>(g))/CH<sub>2</sub>Cl<sub>2</sub>, 1/9, as eluent  
20 gave 0.67 g (38 %) of the title compound as a white powder.

(v) 3-aminomethyl-1-(N-benzyloxycarbonylamidino) azetidine (H-Mig(Z))  
25

0.67 g (1.85 mmol) of Boc-Mig(Z) was dissolved in 10 mL of EtOAc saturated with HCl(g) and stirred for 10 min. at roomtemperature. 10 mL of a saturated solution of KOH(aq) was added dropwise. The layers were separated  
30 and the aqueous phase was extracted with 3x8 mL EtOAc. The organic layers were combined, washed with brine, dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated to yield 0.43 g (89 %) of the title compound.

35 <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 2.55-2.65 (m, 1H), 2.84 (d, 2H), 3.66 (dd, 2H) 4.03 (dd, 2H) 5.07 (s, 2H), 7.2-7.4 (m, 5H).

MS m/z 263 ( $M^+ + 1$ )

**3-aminoethyl-1-(N-benzyloxycarbonylamidino) azetidine  
(H-Dig(Z))**

5

(i) 3-carboxylic acid-1-benzhydryl azetidine was prepared according to the literature, see A.G. Anderson, Jr., and R. Lok, *J.Org.Chem.*, 37, 3953, 1972.

10

**(ii) 3-hydroxymethyl-1-benzhydryl azetidine**

A solution of 8.7 g (32.5 mmol) 3-carboxylic acid-1-benzhydryl azetidine in 80 mL of dry THF was added slowly to a suspension of 4.9 g (130.2 mmol) of  $LiAlH_4$  in 30 mL THF at roomtemperature. The reaction mixture was refluxed for 3.5 h. Excess hydride reagent was hydrolyzed by careful addition, with cooling, of  $NH_4Cl(aq)$ , the gelatinous mixture was filtered and the filter cake was washed repeatedly with THF. Evaporation of the solvent gave 7.1 g (86 %) of the title compound as pale yellow crystals.

15

20

**(iii) 3-methanesulfonatomethyl-1-benzhydryl azetidine**

25

To a solution of 6.62 g (26.1 mmol) 3-hydroxymethyl-1-benzhydryl azetidine in 50 mL of dry pyridine was added 4.50 g (39.2 mmol) of methanesulfonyl chloride at 0°C. The reaction mixture was stirred for 1 h. and then allowed to stand in a refrigerator over night. The reaction mixture was poured into a mixture of ice and  $H_2O$ . The precipitate was collected, washed with  $H_2O$  and dried in vacuo to yield 7.75 g (89.5 %) of the title compound.

30

35

**(iv) 3-cyanomethyl-1-benzhydryl azetidine**

To a solution of 7.75 g (23.4 mmol) 3-

methanesulfonatomethyl-1-benzhydryl azetidine in 50 mL DMF was added a solution of 3.44 g (70.0 mmol) NaCN in 10 mL H<sub>2</sub>O. The mixture was heated at 65°C for 20 h, cooled, and poured into a mixture of ice and H<sub>2</sub>O. The precipitate was collected, washed with H<sub>2</sub>O and dried in vacuo to yield 5.7 g (93 %) of the title compound.

(v) 3-aminoethyl-1-benzhydryl azetidine

5.7 g (21.7 mmol) of 3-cyanomethyl-1-benzhydryl azetidine was added slowly to a suspension of 2.9 g (76.0 mmol) of LiAlH<sub>4</sub> in 80 mL of dry THF at roomtemperature. The reaction mixture was refluxed for 4 h. Excess hydride reagent was hydrolyzed by careful addition, with cooling, of NH<sub>4</sub>Cl(aq), the gelatinous mixture was filtered and the filter cake was washed repeatedly with THF. The solvent was evaporated, the residue was dissolved in diethyl ether, washed with brine and dried with Na<sub>2</sub>SO<sub>4</sub>. Evaporation of the solvent gave 5.0 g (87 %) of the title compound.

(vi) 3-(N-tert-butyloxycarbonylaminoethyl)-1-benzhydryl azetidine

The title compound was prepared from 3-aminoethyl-1-benzhydryl azetidine according the procedure for 3-(N-tert-butyloxycarbonylaminomethyl)-1-benzhydryl azetidine, in a yield of 6.5 g (95 %).

(vii) 3-(N-tert-butyloxycarbonylaminoethyl) azetidine

The title compound was prepared from 3-(N-tert-butyloxycarbonylaminoethyl)-1-benzhydryl azetidine according the procedure for 3-(N-tert-butyloxycarbonylaminomethyl) azetidine, in a yield of 1.2 g (70 %).

(viii) 3-(N-tert-butyloxycarbonylaminoethyl)-1-(N-benzyloxycarbonylamidino) azetidine (Boc-Dig(Z))

5 The title compound was prepared from 3-(N-tert-butyloxycarbonylaminoethyl) azetidine according the procedure for 3-(N-tert-butyloxycarbonylaminomethyl)-1-(N-benzyloxycarbonylamidino) azetidine, in a yield of 0.090 g (34 %).

10 (ix) 3-aminoethyl-1-(N-benzyloxycarbonylamidino) azetidine (H-Dig(Z))

0.589 g (1.56 mmol) of Boc-Dig(Z) was dissolved in 10 mL of EtOAc saturated with HCl(g) and stirred for 10  
15 min. at roomtemperature. 10 mL of a saturated solution of KOH(aq) was added dropwise. The layers were separated and the aqueous phase was extracted with 3x8 mL EtOAc. The organic layers were combined, washed with brine, dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated to yield 0.415 g  
20 (96 %) of the title compound.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 1.60 (dt, 2H), 2.52-2.54 (m, 3H), 3.53 (bs, 2H), 4.0 (bt, 2H), 5.00 (s, 2H), 7.17-7.31 (m, 5H).

25

#### Working Examples

##### Example 1

30 HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab

(i) Boc-(R)Cgl-Aze-Pab(Z)

To a stirred mixture of 3.40 g (10 mmol) Boc-(R)Cgl-Aze-OH (See Preparation of starting materials) and 5.13  
35 g DMAP (42 mmol) in 120 ml CH<sub>3</sub>CN was added 3.18 g H-Pab(Z) x HCl (See Preparation of starting materials).

After stirring for 2 hours at room temperature the mixture was cooled to  $-8^{\circ}\text{C}$  and 2.01 g (10.5 mmol) EDC was added. The reaction was allowed to reach room temperature and the stirring was continued for an additional 47 hours. The solvent was evaporated and the residue was dissolved in 200 ml EtOAc. The organic phase was washed with 1 x 50 ml water, 1 x 50 + 2 x 25 ml 0.5 M  $\text{KHSO}_4$ , 2 x 25 ml  $\text{NaHCO}_3$  (saturated), 1 x 50 ml water and dried. Evaporation of the solvent gave 5.21 g (86%) of the title compound.

$^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.8-1.9 (m, 20H; thereof 1.30 (s, 9H)), 2.35-2.6 (m, 2H), 3.74 (bt, 1H), 4.10 (m, 1H), 4.25-4.4 (m, 2H), 4.45-4.6 (m, 1H, rotamers), 4.75-5.0 (m, 1H, rotamers), 5.08 (bd, 2H), 5.15 (s, 2H), 7.15-7.35 (m, 5H), 7.41 (d, 2H), 7.77 (d, 2H), 8.21 (m, 1H).

(ii) H-(R)Cgl-Aze-Pab(Z)

To a cold (ice bath temperature) solution of 18.8 g  $\text{HCl}_{(g)}$  in 195 ml EtOAc was added 4.69 g (7.743 mmol) of Boc-(R)Cgl-Aze-Pab(Z) together with 40 ml EtOAc. The reaction mixture was allowed to reach room temperature and stirred for 30 min. 140 ml  $\text{Et}_2\text{O}$  was added to the clear solution where upon a precipitate was formed. The reaction was left at room temperature for an additional 1 h and 40 minutes. The precipitate was filtered off, washed quickly with 150 ml  $\text{Et}_2\text{O}$  and dried in vacuo. The precipitate was dissolved in 50 ml of water and made alkaline with 15 ml 2 M NaOH. The alkaline waterphase was extracted with 1 x 100 + 1 x 50 ml  $\text{CH}_2\text{Cl}_2$ . The combined organic phase was washed with 1 x 20 ml water, 1 x 20 ml Brine and dried ( $\text{MgSO}_4$ ). Evaporation of the solvent gave 3.44 g (88%) of the title compound.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.8-2.0 (m, 11H), 2.51 (m, 1H), 2.67 (m, 1H), 3.07 (d, 1H), 4.11 (m, 1H), 4.18 (m, 1H), 4.43 (dd, 1H), 4.53 (dd, 1H), 4.91 (m, 1H), 5.22 (s, 2H), 7.2-7.4 (m, 7H), 7.45 (d, 2H), 8.51 (d, 2H).

5

(iii) BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)

1.13 g (2.2 mmol) H-(R)Cgl-Aze-Pab(Z), 0.9 g (2.6 mmol) benzyl-2-(orto-nitrobenzenesulfonyloxy)acetate ((2-NO<sub>2</sub>)Ph-SO<sub>2</sub>-OCH<sub>2</sub>-COOBn) (See Preparation of starting materials), 0.99 g (5.6 mmol) K<sub>2</sub>CO<sub>3</sub> and 113 ml CH<sub>3</sub>CN were mixed and heated in a 60°C oilbath for 3 h. The solvent was evaporated in vacuo. EtOAc was added and the mixture was washed with water, the organic phase was extracted with 1 M KHSO<sub>4</sub> and this waterphase was washed with EtOAc. The acidic waterphase was made alkaline with 1 N NaOH to pH>8 and extracted with EtOAc. The organic phase was washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated in vacuo to give 1.17 g of a residue that was twice subjected to flash chromatography using first CH<sub>2</sub>Cl<sub>2</sub>/MeOH(NH<sub>3</sub>-saturated) 95/5 and then diethylether/MeOH(NH<sub>3</sub>-saturated) 9/1 as eluents to give 0.525 g (36 %) of the title compound.

The alkylation was also carried out using Benzyl-2-(para-nitrobenzenesulfonyloxy)acetate ((4-NO<sub>2</sub>)Ph-SO<sub>2</sub>-OCH<sub>2</sub>-COOBn) (See Preparation of starting materials) using the same procedure as above to give the title compound in 52 % yield.

30

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 0.85-2.15 (m, 11H), 2.48 (m, 1H), 2.63 (m, 1H), 2.88 (d, 1H), 3.24 (d, 1H), 3.27 (d, 1H), 3.95 (m, 1H), 4.05 (m, 1H), 4.44 (m, 1H), 4.55 (m, 1H), 4.91 (m, 1H), 5.07 (s, 2H), 5.22 (s, 2H), 7.2-7.4 (m, 10H), 7.45 (d, 2H), 7.79 (d, 2H), 8.42 (m, 1H).

35

(iva) HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab x 2 HCl

BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z), 20 mg (0.031 mmol), was dissolved in 5 ml of methanol. A few drops of chloroform and 5 % Pd/C were added and the mixture was hydrogenated at atmospheric pressure for 1 h. After filtration and evaporation the product was lyophilized from water to give 11 mg (72%) of the title compound.

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O): δ 1.0-2.0 (m, 11H), 2.10 (m, 1H), 2.44 (m, 1H), 2.82 (m, 1H), 3.90 (s, 2H), 4.09 (d, 1H), 4.4-4.55 (m, 2H), 4.66 (s, 2H), 5.08 (m, 1H), 7.65 (d, 2H), 7.89 (d, 2H).

<sup>13</sup>C-NMR (75.5 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 167.3, 167.9, 169.9 and 172.4.

(ivb) HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab

BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z) was dissolved in EtOH (99%) and hydrogenated over 5 % Pd/C at atmospheric pressure for 5 hours. Filtration of the catalyst through cellite and evaporation of the solvent gave the title compound in 97 % yield.

<sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD, mixture of two rotamers): major rotamer: δ 1.00-1.12 (m, 1H), 1.13-1.34 (m, 4H), 1.55-1.70 (m, 3H), 1.73-1.85 (m, 2H), 1.94-2.02 (bd, 1H), 2.32-2.42 (m, 1H), 2.54-2.64 (m, 1H), 2.95-3.10 (AB-system plus d, 3H), 4.18-4.25 (bq, 1H), 4.28-4.32 (bq, 1H), 4.43-4.60 (AB-system, 2H), 4.80-4.85 (dd, 1H), 7.48-7.54 (d, 2H), 7.66-7.71 (d, 2H).

Resolved signals from the minor rotamer appears at δ 0.95 (m), 1.43 (m), 2.24 (m), 2.84 (d), 3.96 (m), 4.03 (m), 7.57 (bd), 7.78 (bd).

<sup>13</sup>C-NMR (125 MHz, CD<sub>3</sub>OD): amidine and carbonyl carbons: δ 168.0, 173.0, 176.3 and 179.0



Example 2

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pab x 2 HCl

5 (i) H-(R)Cgl-Aze-Pab(Z)

Prepared in the same way as described in Example 1 (ii) by treating the formed hydrochloride salt with base to afford the free base.

10

(ii) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)

H-(R)Cgl-Aze-Pab(Z), 0.19 g (0.38 mmol), and 70 mg (0.43 mmol) of benzyl acrylate were dissolved in 2 ml of isopropanol. The mixture was left standing for 6 days. Flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/THF = 8/2 as eluent afforded 0.12 g (48%) of the title compound.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.8-1.9 (m, 10 H), 1.95 (bd, 1 H), 2.4-2.6 (m, 4 H), 2.7-2.8 (m, 3 H; thereof 2.79 (d, 1 H)), 4.13 (m, 1 H), 4.37 (dd, 1 H), 4.60 (dd, 1 H), 4.97 (dd, 1 H), 5.09 (dd, 2 H), 5.22 (s, 2 H), 7.25-7.4 (m, 10 H), 7.47 (d, 2 H), 7.83 (d, 2 H), 8.61 (bt, 1 H).

25 (iii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pab x 2 HCl

BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z), 0.10 g (0.15 mmol), was dissolved in 10 ml of ethanol and hydrogenated over 5% Pd/C at atmospheric pressure for 1 h.

30 The solution was filtered, evaporated and the crude product was purified on RPLC using CH<sub>3</sub>CN/0.1 M NH<sub>4</sub>OAc (1/4). The resulting product was freeze dried, treated with an excess of conc. HCl and freeze dried again to give 31 mg of the dihydrochloride salt.

35

<sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O) δ 0.8-2.1 (m, 11 H), 2.38 (m, 1 H), 2.7-2.9 (m, 3 H), 3.2-3.4 (m, 2 H), 3.98 (d, 1 H), 4.35-

4.55 (m, 2 H), 4.60 (s, 2 H), 5.04 (dd, 1 H), 7.59 (d, 2 H), 7.83 (d, 2 H).

<sup>13</sup>C NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ  
5 167.2, 167.8, 172.3 and 175.5.

Example 3

HOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pab x 2 HCl

10

(i) Boc-(R)Cgl-Pro-Pab(Z)

Boc-(R)Cgl-Pro-OH (See preparation of starting  
materials), 2.3 g (6.49 mmol), DMAP, 2.38 g (19.47  
15 mmol), and H-Pab(Z) (See preparation of starting  
materials), 1.84 g (6.49 mmol) was mixed in 30 ml  
acetonitrile. The mixture was cooled to -15°C and EDC,  
1.31 g (6.81 mmole) was added. The temperature was  
20 allowed to reach room temperature and the mixture was  
stirred over night. After evaporation, the residue was  
dissolved in ethyl acetate and 0.3 M KHSO<sub>4</sub>-solution. The  
acidic water phase was extracted three times with ethyl  
acetate. The organic phase was washed twice with a 0.3 M  
KHSO<sub>4</sub>-solution, twice with a NaHCO<sub>3</sub>-solution and once with  
25 brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated. The crude  
product was purified by flash chromatography on silica  
gel using ethyl acetate as eluent to yield 1.77 g (44%  
) of the product.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.9-1.49 (m, 14H), 1.5-2.1 (m,  
9H), 2.37 (bs, 1H), 3.53 (q, 1H), 3.94 (bs, 1H), 4.02 (m,  
1H), 4.43 (bs, 2H), 4.65 (d, 1H), 5.09 (bs, 1H), 5.20 (s,  
2H), 7.18-7.4 (m, 5H), 7.45 (d, 2H), 7.62 (bs, 1H), 7.81  
(m, 2H),

35

(ii) H-(R)Cgl-Pro-Pab(Z)

1.45 g (2.34 mmol) of Boc-(R)Cgl-Pro-Pab(Z) was dissolved in 75 ml HCl saturated ethyl acetate. The mixture was allowed to stand for 10 min at room temperature. The solvent was evaporated to yield 1.3 g of the dihydrochloride salt of the product.

$^1\text{H-NMR}$  (300 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  1.0-1.45 (m, 5H), 1.58-2.2 (m, 9H), 2.3-2.5 (m, 1H), 3.75-3.90 (m, 2H), 4.25 (d, 2H), 4.5-4.66 (m, 3H), 5.49 (s, 2H), 7.45-7.7 (m, 7H), 7.87 (d, 2H)

The amine was obtained by dissolving the dihydrochloride salt in 0.1 M NaOH-solution and extracting the water phase three times with ethyl acetate. The organic phase was washed once with brine, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to yield 1.19 g (97%) of the title compound.

(iii)  $\text{BnOOC-CH}_2\text{-(R)Cgl-Pro-Pab(Z)}$

0.340 g (0.65 mmole) H-(R)Cgl-Pro-Pab(Z) was mixed with 0.215 g (0.65 mmole)  $\text{BnOOC-CH}_2\text{-OTf}$  (see preparation of starting materials), 0.299 g (2.17 mmole)  $\text{K}_2\text{CO}_3$  in 4 ml dichloromethane and refluxed for half an hour. The reaction mixture was then stirred over night at room temperature. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  and the organic layer was washed once with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. The crude product was purified by flash chromatography using a stepwise gradient of  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (97/3 followed by 95/5) to give 299 mg of a mixture of two products according to TLC. The mixture was therefore purified further by flash chromatography using a stepwise gradient of ethyl acetate/toluene (9/1, 93/7, 95/5, 100/0) to give 46 mg (9%) of  $(\text{BnOOC-CH}_2)_2\text{-(R)Cgl-Pro-Pab(Z)}$  which eluted first from the column followed by 133 mg (31%) of the desired product  $\text{BnOOC-CH}_2\text{-(R)Cgl-Pro-Pab(Z)}$ .

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): BnOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pab(Z): δ 0.9-1.3 (m, 5H), 1.4-2.1 (m, 9H), 2.3-2.4 (m, 1H), 3.05 (d, 1H), 3.20-3.37 (AB-system centered at δ 3.29, 2 H), 3.5-3.6 (m, 2H), 4.29-4.57 (ABX-system centered at d 4.43, 2 H), 4.62 (d, 1H), 4.91 (apparent singlet, 2H), 5.19 (s, 2H), 6.75 (bs, NH), 7.1-7.5 (m, 12H), 8.7-8.8 (m, 2H+NH), 9.45 (bs, NH)

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R)Cgl-Pro-Pab(Z): δ 0.68-0.9 (m, 2H), 1.0-1.3 (m, 3H), 1.43 (bd, 1H), 1.55-2.0 (m, 7H), 2.05 (bd, 1H), 2.3-2.4 (m, 1H), 3.15 (d, 1H), 3.25-3.48 (m, 2H), 3.55-3.79 (AB-system centered at d 3.67, 4 H), 4.38-4.58 (ABX-system centered at d 4.48, 2 H), 4.68 (d, 1H), 4.82-4.98 (AB-system centered at d 4.91, 4 H), 5.19 (s, 2H), 6.66 (bs, NH), 7.1-7.5 (m, 17H), 7.75 (d, 2H), 7.80 (t, NH), 9.37 (bs, NH)

<sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 164.7, 168.1, 171.5, 172.3 and 172.6

(iv) HOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pab x 2 HCl

0.133 g (0.20 mmole) of BnOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pab(Z) was mixed with 0.060 g 5 % Pd/C, 1ml 1M HCl-solution and 10 ml ethanol. The mixture was treated under H<sub>2</sub>-atmosphere for one hour. After filtration through hyflo and evaporation of the solvent the product in 90% yield, 93 mg, was obtained by freeze drying twice from water.

<sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 1.0-1.45 (m, 5H), 1.5-2.1 (m, 9H), 2.2-2.4 (m, 1H), 3.55-3.85 (m, 4H; thereof 3.79 (s, 2H)), 4.23 (d, 1H), 4.33-4.57 (m, 3H), 7.44 (d, 2H), 7.69 (d, 2H)

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 166.9, 167.2, 169.1, 174.5

Example 4

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pab x 2 HCl

5 (i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pab(Z)

0.406 g (0.782 mmole) of H-(R)Cgl-Pro-Pab(Z) (See Example 3) was dissolved in 3 ml ethanol and 132 µl (0.861 mmole) of benzylacrylate was added. The reaction mixture was stirred for three days at room temperature. The mixture was evaporated and the crude product purified by flash chromatography using a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>:MeOH 95/5 and 90/10 as eluent to yield 0.399 g (75%) of the product.

15

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 0.8-1.0 (m, 1H), 1.0-1.3 (m, 4H), 1.35-2.2 (m, 9H), 2.3-2.6 (m, 4H), 2.65-2.78 (m, 1H), 3.05 (d, 1H), 3.4-3.6 (m, 2H), 4.25-4.52 (ABX-system central at d 4.40, 2 H), 4.64 (dd, 1H), 5.05 (s, 2H), 5.20 (s, 2H), 7.2-7.38 (m, 10H), 7.43 (d, 2H), 7.78 (d, 2H)

20

<sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 164.7, 167.9, 171.3, 172.7 and 175.4.

25

(ii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pab x 2 HCl

0.261 g (0.383 mmole) of BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pab(Z) was mixed with 0.075 g 5 % Pd/C, 1ml 1M HCl-solution and 10 ml ethanol. The mixture was hydrogenated at atmospheric pressure for two hours. After filtration through hyflo and evaporation of the solvent the product 0.196 g (96%) was obtained by freeze drying twice from water

35

<sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 1.17-1.40 (m, 5H), 1.60-1.92 (m, 5H), 1.92-2.2 (m, 4H), 2.32-2.48 (m, 1H), 2.81 (t, 2H),

3.11-3.36 (ABX<sub>2</sub>-system centered at  $\delta$  3.24, 2H), 3.63-3.90 (m, 2H), 4.25 (d, 1H), 4.42-4.63 (m, 3H), 7.54 (d, 2H), 7.78 (d, 2H)

- 5 <sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons:  $\delta$  167.0, 167.30, 174.6 and 174.7.

#### Example 5

- 10 (HOOC-CH<sub>2</sub>)<sub>2</sub>-(R)Cgl-Pro-Pab x 2 HCl

- 46 mg (0.056 mmole) of (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R)Cgl-Pro-Pab(Z) (See Example 3) was mixed with 25 mg 5 % Pd/C, 0.7 ml 1M HCl-solution and 7 ml ethanol. The mixture was hydrogenated  
15 at atmospheric pressure for one hour. The catalyst was filtered off through hyflo and the solvent evaporated. The final product 25 mg (77 %) was obtained by freeze drying twice from water.

- 20 <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O):  $\delta$  1.0-1.4 (m, 5H), 1.45-2.2 (m, 9H), 2.25-2.45 (m, 1H), 3.53-3.84 (m, 2H), 3.84-4.22 (AB-system centered at  $\delta$  4.03, 4 H), 4.26 (d, 1H), 4.35-4.6 (m, 3H), 7.53(d, 2H), 7.77 (d, 2H)

- 25 <sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons:  $\delta$  167.1, 167.3, 170.6 and 174.5

#### Example 6

- 30 H-(R)Cgl-Pic-Pab x 2 HCl

(i) Boc-(R)Cgl-Pic-Pab(Z)

- 0.478 g (2.49 mmol) EDC was added at -18°C to a stirred  
35 solution of 0.875 g (2.37 mmol) Boc-(R)Cgl-Pic-OH (See preparation of starting materials), 1.22 g (9.97 mmol) DMAP, and 0.706 g (2.49 mmol) H-Pab(Z) (See preparation

f starting materials) in a mixture of 30 ml acetonitrile and 1 ml DMF. The reaction mixture was allowed to reach room temperature during a couple of hours and stirring was continued for 48 h. The solvent was removed in vacuo and the residue was dissolved in 50 ml ethyl acetate. The solution was washed with 15 ml water, 3x15 ml 0.3 M KHSO<sub>4</sub>, 2x15 ml Na<sub>2</sub>CO<sub>3</sub> solution and water. Removal of the solvent gave a residue which was subjected to flash chromatography using ethyl acetate/heptane 9:1 as eluent. The yield was 0.96 g (64%).

(ii) H-(R)Cgl-Pic-Pab(Z)

Hydrogen chloride was bubbled through a solution of 0.56 g (0.88 mmol) Boc-(R)Cgl-Pic-Pab(Z) in 25 ml ethyl acetate. After a couple of minutes crystals precipitated from the solution. The solvent was removed in vacuo and 50 ml ethyl acetate was added. Washing with 2x15 ml 2 M sodium hydroxide solution and extraction of the aqueous phase with 25 ml ethyl acetate was followed by drying (sodium sulphate) of the combined extracts and removal of the solvent in vacuo to give 0.448 g (95%) of the desired product.

(iii) H-(R)Cgl-Pic-Pab x 2 HCl

A solution of 98 mg (0.18 mmol) H-Cgl-Pic-Pab(Z) in 5 ml 95% ethanol and 1 ml water was stirred in an atmosphere of hydrogen for 4 hours in the presence of 5 % Pd/C. The mixture was filtered and 0.3 ml 1 M hydrochloric acid was added. The ethanol was removed in vacuo and the residue was freeze dried to give 70 mg (81%) of the desired compound.

<sup>1</sup>H-NMR (300 MHz, CD<sub>3</sub>OD): δ 1.00-1.56 (m, 7H), 1.56-1.94 (m, 9H), 2.32 (bd, 1H), 3.32-3.45 (m, 1H), 3.90 (bd, 1H), 4.35 (d, 1H), 4.50 (s, 2H), 5.10-5.20 (m, 1H), 7.55 (d,

2H), 7.76 (d, 2H)

$^{13}\text{C}$ -NMR (75 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.2, 170.5 and 173.4.

5

Example 7

$\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Cgl-Pic-Pab} \times 2 \text{ HCl}$

10 (i)  $\text{BnOOC-CH}_2\text{-(R,S)CH(COObn)-(R)Cgl-Pic-Pab(Z)}$

A mixture of 350 mg (0.66 mmol) H-(R)Cgl-Pic-Pab(Z) (See Example 6) and 233 mg dibenzyl maleate in 2.5 ml ethanol was kept at room temperature for 4 days. The ethanol was removed in vacuo and the residue was subjected to flash chromatography using ethyl acetate/heptane 9:1 as eluent to give 0.108 mg of the product.

15

(ii)  $\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Cgl-Pic-Pab} \times 2 \text{ HCl}$

20

105 mg (0.13 mmol)  $\text{BnOOC-CH}_2\text{-(R,S)CH(COObn)-(R)Cgl-Pic-Pab(Z)}$  dissolved in 5 ml 95% ethanol and 1 ml water was hydrogenated for 5 hours in the presence of 5 % Pd/C. 0.3 ml 1 M hydrochloric acid was added and the mixture was filtered and the solvent was removed in vacuo. The residue was dissolved in water and freeze dried to yield 54 mg (73%) of the desired substance.

25

$^1\text{H}$ -NMR (300 MHz,  $\text{CD}_3\text{OD}$ , mixture of two diastereomers 5/4):  
 $\delta$  1.10-1.60 (m, 7H), 1.60-2.04 (m, 9H), 2.23-2.42 (m, 1H), 2.93-3.15 (m 2H), 3.30-3.42 (m, 1H, partially hidden by the MeOD-peak), 3.71-3.95 (m, 1H), 3.98-4.10 (m, 1H), 4.40-4.60 (m, 3H), 5.10-5.20 (m, 1H), 7.49-7.60 (m, 2H), 7.70-7.81 (m, 2H)

35

$^{13}\text{C}$ -NMR (75 MHz  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.1, 168.95, 169.6 and 173.1.



MS m/z 516 ( $M^+ + 1$ )

Example 8

5 H-(R)Cha-Aze-Pab x 2 HCl

(i) Boc-(R)Cha-Aze-Pab(Z)

409 mg (2.13 mmol) EDC was added at  $-18^{\circ}\text{C}$  to a stirred  
10 mixture of 0.72 g (2.03 mmol) Boc-(R)Cha-Aze-OH (See  
preparation of starting materials), 1.04 g (8.53 mmol)  
DMAP, and 604 mg (2.13 mmol) H-Pab(Z) (See preparation of  
starting materials) in 20 ml acetonitrile. The reaction  
15 mixture was allowed to reach room temperature over night  
and the solvent was subsequently removed in vacuo. The  
residue was dissolved in 40 ml ethyl acetate and the  
organic phase was washed succesively with 10 ml water,  
3x10 ml 0.3M  $\text{KHSO}_4$ , 2x10 ml  $\text{Na}_2\text{CO}_3\text{-NaCl}$  (aq), and finally  
20 10 ml Brine. Drying ( $\text{Na}_2\text{SO}_4$ ) and removal of the solvent in  
vacuo gave a residue which was subjected to flash  
chromatography using ethyl acetate/methanol 9:1 as eluent  
to yield 645 mg (51%) of the title compound.

(ii) H-(R)Cha-Aze-Pab(Z)

25

Hydrogen chloride was bubbled through a solution of 640  
mg (1.03 mmol) Boc-(R)Cha-Aze-Pab(Z) in 25 ml of ethyl  
acetate. After a couple of minutes, TLC analysis  
indicated the completion of the reaction. Vacuum was  
30 applied to remove excess hydrogen chloride and the  
mixture was then diluted to 50 ml with ethyl acetate.  
Washing with 2x15 ml  $\text{Na}_2\text{CO}_3$  (aq) was followed by  
extraction of the aqueous phase with 15 ml ethyl acetate.  
The combined organic extracts were washed with water and  
35 dried ( $\text{Na}_2\text{CO}_3$ ) and the solvent was removed in vacuo to  
give 513 mg (96%) of H-(R)Cha-Aze-Pab(Z).

(iii) H-(R)Cha-Aze-Pab x 2 HCl

76 mg (0.15 mmol) H-(R)Cha-Aze-Pab(Z) dissolved in 5 ml 95% ethanol and 1 ml water was hydrogenated at atmospheric pressure in the presence of 5% Pd/C for 4 h. Removal of the catalyst by filtration, addition of 0.4 ml 1M hydrochloric acid and evaporation of the solvent in vacuo gave a residue which was dissolved in 2 ml water. Freeze drying gave 57 mg (85%) of the product.

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O, 2 rotamers, 3:1 mixture): δ 1.02-1.20 (m, 2H), 1.22-1.92 (m, 11H), 2.40-2.50 (m, 1H), 2.80-2.90 (m, 1H), 4.25 (bt, 1H), 4.40 (dq, 1H), 4.53 (dq, 1H), 4.65 (s, 2H), 5.05-5.10 (m, 1H), 7.65 (d, 2H), 7.88 (d, 2H).

Chemical shifts of resolved signals of the minor rotamer:

δ 0.57 (m), 0.85 (m), 2.95 (m), 4.06 (dq), 4.17 (dq), 4.63 (s), 5.33(m), 7.70(d), 7.93 (d).

<sup>13</sup>C-NMR (125 MHz D<sub>2</sub>O): amidine and carbonyl carbons: δ 167.2, 170.4 and 172.8.

#### Example 9

HOOC-CH<sub>2</sub>-(R)Cha-Aze-Pab x 2 HCl

(i) BnOOC-CH<sub>2</sub>-(R)Cha-Aze-Pab(Z)

0.119 g (0.52 mmol) benzyl bromoacetate was added to a mixture of 0.27 g (0.52 mmol) H-(R)Cha-Aze-Pab(Z) (See Example 8) and 0.158 g (1.14 mmol) K<sub>2</sub>CO<sub>3</sub> in 5.2 ml acetonitrile and heated to 60°C in an oilbath for 1 h. The solvent was removed and ethyl acetate and water was added. The phases were separated and the organic phase was washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation in vacuo gave 0.344 g of a residue which was subjected to

flash chromatography using thyl acetate as eluent, and then another time using ethylacetate:tetrahydrofuran:  $\text{NH}_3$ -saturated methanol (60:5:2) to give 0.163 g of the desired product.

5

$^1\text{H}$ -NMR (300 MHz,  $\text{CDCl}_3$ );  $\delta$  0.7-1.0 (m, 2H), 1.05-2.05 (m, 11H), 2.35-2.55 (m, 1H), 2.55-2.75 (m, 1H), 3.15-3.32 (m, 3H), 3.95-4.05 (t, 2H), 4.4 and 4.5 (ABX-system, 2H), 4.8-4.95 (m, 1H), 5.05 (s, 2H), 5.2 (s, 2H), 7.2-7.5 (m, 12H), 7.7-7.85 (d, 2H), 8.3-8.45 (t, 1H).

10

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.5, 167.8, 170.7, 171.9 and 175.9.

15

(ii)  $\text{HOOC-CH}_2\text{-(R)Cha-Aze-Pab} \times 2 \text{ HCl}$

0.163 g (0.243 mmol)  $\text{BnOOC-CH}_2\text{-(R)Cha-Aze-Pab(Z)}$  dissolved in 5.5 ml ethanol (99.5 %) and 0.7 ml hydrogen chloride (1 N) was hydrogenated in the presence of 0.17 g 5 % Pd/C for 4 h. Removal of the catalyst by filtration and evaporation of the solvent followed by dissolving in water and freeze drying gave 107 mg (85 %) of the title compound.

20

$^1\text{H}$ -NMR (500 MHz,  $\text{CD}_3\text{OD}$ , mixture of two rotamers): major rotamer:  $\delta$  0.95-1.95 (m, 13H), 2.3-2.4 (m, 1H), 2.6-2.75 (m, 1H), 3.5-3.75 (m, 2H), 4.05-4.15 (m, 1H), 4.15-4.23 (m, 1H), 4.36-4.43 (m, 1H), 4.43-4.5 (m, 1H), 4.58-4.65 (m, 1H), 4.83-4.88 (m, 1H), 7.5-7.6 (m, 2H), 7.73-7.82 (m, 2H).

30

Resolved signals from the minor rotamer appears at  $\delta$  2.2-2.3 (m), 3.95-4.05 (m), 5.1-5.17 (m), 7.6-7.67 (m).

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CD}_3\text{OD}$ ): amidine and carbonyl carbons;  $\delta$  168.2, 169.8, 168.9 and 172.3.

35

Example 10

HOOC-CH<sub>2</sub>-(R,S)CH(COOH)-(R)Cha-Aze-Pab x 2 HCl

5 (i) BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Aze-Pab(Z)

10 A mixture of 230 mg (0.443 mmol) H-(R)Cha-Aze-Pab(Z) (See Example 8) and 144 mg (0.487 mmol) dibenzyl maleate in 1.5 ml 95% ethanol was stirred at ambient temperature for 5 days. After removal of the ethanol in vacuo, the residue was subjected to flash chromatography using ethyl acetate/methanol 95/5 as eluent to give 54 mg (15%) of the product.

15 (ii) HOOC-CH<sub>2</sub>-(R,S)CH(COOH)-(R)Cha-Aze-Pab

49 mg (0.06 mmol) BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Aze-Pab(Z) dissolved in 5 ml 95% ethanol and 1 ml water was hydrogenated in the presence of 5% Pd/C for 4.5 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave a residue which was dissolved in 2 ml water and 0.2 ml 1M hydrochloric acid. Freeze drying gave 32 mg (93%) of the product.

25 The <sup>1</sup>H-NMR spectrum of the title compound in D<sub>2</sub>O exhibits two sets of strongly overlapping signals arising from the two diastereomers. Additionally resolved resonances of a minor rotamer, integrating to approximately 15% also appears in the spectrum.

30 <sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 1.03-2.00 (m, 13H), 2.32-2.53 (m, 1H), 2.72-2.96 (m, 1H), 3.06-3.28 (m, 2H), 4.10-4.55 (m, 4H), 4.62 (bs, 2H), 5.00-5.10 (m, 1H), 7.55-7.68 (m, 2H), 7.80-7.94 (m, 2H)

35 Resolved signals from the minor rotamer appears at δ 0.65 (m), 0.80 (m), 4.00 (m), 5.24 (m), 5.35 (m).

$^{13}\text{C}$ -NMR (75 MHz  $\text{D}_2\text{O}$ ): amidin and carbonyl carbons:  
 $\delta$  167.2, 169.0, 171.0, 172.3 and 174.1.

Example 11

5

**$\text{HOOC}-\text{CH}_2-(\text{RorS})\text{CH}(\text{COOH})-\text{Cha-Aze-Pab/a} \times 2 \text{ HCl}$**

(i)  **$\text{BnOOC}-\text{CH}_2-(\text{RorS})\text{CH}(\text{COOBn})-(\text{R})\text{Cha-Aze-Pab(Z)}/\text{a}$**

10 A mixture of 2.0 g (3.8491 mmol) H-(R)Cha-Aze-Pab(Z) (See Example 8) and 1.37 g dibenzyl maleate in 10 ml 95% ethanol was stirred at ambient temperature for 4 days. After removal of the ethanol in vacuo, the residue was subjected to flash chromatography using ethyl acetate/methanol 98/2 as eluent to give 1.024 g (32%) of  
15  **$\text{BnOOC}-\text{CH}_2-(\text{R,S})\text{CH}(\text{COOBn})-(\text{R})\text{Cha-Aze-Pab(Z)}$** . The two diastereomers were separated by RPLC using ( $\text{CH}_3\text{CN}/0.1 \text{ M NH}_4\text{OAc}$  65/35) as eluent. This diastereomer eluted first from the column. After removal of the acetonitrile in  
20 vacuo the water phase was extracted three times with ethyl acetate. The organic phase was washed once with water dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to yield 0.352 g of the title compound as a pure stereoisomer.

25

(ii)  **$\text{HOOC}-\text{CH}_2-(\text{RorS})\text{CH}(\text{COOH})-(\text{R})\text{Cha-Aze-Pab/a} \times 2 \text{ HCl}$**

350 mg (0.43 mmol)  **$\text{BnOOC}-\text{CH}_2-(\text{RorS})\text{CH}(\text{COOBn})-(\text{R})\text{Cha-Aze-Pab(Z)}/\text{a}$**  (The diastereomer from (i) above) dissolved in  
30 15 ml 95% ethanol and 3 ml water was hydrogenated in the presence of 5% Pd/C for 4.5 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave a residue which was dissolved in 5 ml water and 1.0 ml 1M hydrochloric acid. Freeze drying gave 214 mg (87%) of the  
35 product as a pure stereoisomer.

<sup>1</sup>H-NMR (300 MHz, MeOD, mixture of two rotamers): δ 0.85-1.93 (m, 13H), 2.25-2.38 (m, 1H), 2.60-2.75 (m, 1H), 2.88 (dd, 2H), 3.92 (t, 1H), 4.15-4.25 (m, 2H), 4.30-4.43 (m, 1H), 4.56 (AB-system, 2H), 4.76-4.86 (m, 1H, partially obscured by the solvent signal), 7.59 (d, 2H), 7.78 (d, 2H).

Resolved signals arising from the minor rotamer appears at δ 0.70, 2.95, 3.82, 4.00, 5.08 and 7.83

<sup>13</sup>C-NMR (75 MHz D<sub>2</sub>O): amidine and carbonyl carbons: δ 166.9, 168.8, 171.7, 172.3 and 173.8.

#### Example 12

HOOC-CH<sub>2</sub>-(RorS)CH(COOH)-(R)Cha-Aze-Pab/b x 2 HCl

(i) BnOOC-CH<sub>2</sub>-(RorS)CH(COOBn)-(R)Cha-Aze-Pab(Z)/b

The title compound was obtained by using the same procedure as described in Example 11 above on BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Aze-Pab(Z). This diastereomer came out after the first one from the column. Yield 0.537 g.

(ii) HOOC-CH<sub>2</sub>-(RorS)CH(COOH)-(R)Cha-Aze-Pab/b x 2 HCl

530 mg (0.65 mmol) BnOOC-CH<sub>2</sub>-(RorS)CH(COOBn)-(R)Cha-Aze-Pab(Z)/b dissolved in 15 ml 95% ethanol and 3 ml water was hydrogenated in the presence of 5% Pd/C for 5 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave a residue which was dissolved in 6 ml water and 1.0 ml 1M hydrochloric acid. Freeze drying gave 290 mg (78%) of the product.

<sup>1</sup>H-NMR (300 MHz, MeOD, mixture of two rotamers): δ 0.86-1.90 (m, 13H), 2.30-2.42 (m, 1H), 2.60-2.75 (m, 1H), 2.75-

2.85 (m, 1H), 2.95-3.05 (m, 1H), 3.65-3.71 (m, 1H), 4.00-4.10 (m, 1H), 4.14-4.24 (m, 1H), 4.36-4.62 (m, 3H), 4.78-4.86 (m, 1H partially obscured by the solvent signal), 7.57 (d, 2H), 7.75 (d, 2H).

5

Resolved signals arising from a minor rotamer appears at  $\delta$  0.78, 2.92, 3.82, 5.36 and 7.80

$^{13}\text{C}$ -NMR (75 MHz  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  166.8, 169.0, 172.0, 172.4 and 175.2.

10

### Example 13

$\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pab x 2 HCl}$

15

(i)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pab(Z)}$

20

A mixture of 182 mg (0.35 mmol) H-(R)Cha-Aze-Pab(Z) (See Example 8) and 62.5 mg (0.385 mmol) benzyl acrylate in 1.5 ml 95% ethanol was stirred at room temperature for 4 days. The solvent was removed in vacuo and the residue was subjected to flash chromatography using ethyl acetate/methanol 9:1 as eluent to give 200 mg (84%) of the title compound.

25

(ii)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pab x 2 HCl}$

30

195 mg (0.29 mmol)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pab(Z)}$  dissolved in 10 ml 95% ethanol and 2 ml water was hydrogenated in the presence of 5% Pd/C for 4 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave a residue which was dissolved in 2 ml water and 0.4 ml 1M hydrochloric acid. Freeze drying gave 130 mg (86%) of the product.

35

$^1\text{H}$ -NMR (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  0.98-1.27 (m, 2H), 1.30-1.90 (m, 11H), 2.27-2.35 (m, 1H), 2.65-2.74 (m, 1H), 2.77 (t,

2H), 3.32 (t, 2H), 4.10 (t, 1H), 4.17-4.25 (m, 1H), 4.40-4.49 (m, 1H), 4.55 (AB, 2H), 4.83-4.90 (m, 1H), 7.58 (d, 2H), 7.77 (d, 2H).

- 5  $^{13}\text{C}$ -NMR (125 MHz  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.0, 168.9, 172.4 and 174.6.

Example 14

- 10  $\text{HOOC-CH}_2\text{-NH-CO-CH}_2\text{-(R)Cha-Aze-Pab} \times 2 \text{ HCl}$

(i)  $\text{BnOOC-CH}_2\text{-NH-CO-CH}_2\text{-(R)Cha-Aze-Pab(Z)}$

- 15 A mixture of 0.212 g ( 0.408 mmole ) H-(R)Cha-Aze-Pab(Z) (See Example 8), 0.124 g ( 0.89 mmole )  $\text{K}_2\text{CO}_3$  and 0.128 g ( 0.449 mmole )  $\text{BnOOC-CH}_2\text{-NH-CO-CH}_2\text{-Br}$  (See preparation of starting materials) in 6 ml acetonitrile was stirred at 50°C for two hours. After evaporation of the solvent the residue was dissolved in water and ethyl acetate. The water layer was extracted twice with ethyl acetate and the combined organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. The product was purified by flash chromatography using a stepwise gradient of ethyl acetate/tetrahydrofuran ( 85/15, 4/1, 7/3 ) to yield 25 0.190 g ( 64% ) of the title compound.

- 30  $^1\text{H}$ -NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.75-2.1 (m, 13H), 2.43 (m, 1H), 2.56 (d, 1H), 2.79 (m, 1H), 3.0-3.15 (m, 2H; thereof 3.05 (d, 1H)), 3.89-4.18 (m, 5H), 4.8-4.98 (m, 2H), 5.15 (s, 2H), 5.18 (s, 2H), 7.2-7.47 (m, 12H), 7.72 (t, NH), 7.86 (d, 2H), 8.14 (bs, NH), 8.31 (dd, NH), 9.42 (bs, NH)

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.5, 168.7, 169.22, 169.83, 171.7, 175.5

35

(ii)  $\text{HOOC-CH}_2\text{-NH-CO-CH}_2\text{-(R)Cha-Aze-Pab} \times 2 \text{ HCl}$



0.19 g ( 0.26 mmole ) of  $\text{BnOOC-CH}_2\text{-NH-CO-CH}_2\text{-(R)Cha-Az - Pab(Z)}$  was mixed with 0.075 g 5 % Pd/C, 1.5 ml 1N HCl-solution, 3 ml water and 17 ml ethanol and the mixture was hydrogenated at atmospheric pressure for one hour.  
5 Filtration of the catalyst, evaporation of the solvent followed by freeze drying from water gave 144 mg (97 %) of the title compound.

$^1\text{H-NMR}$  (  $\text{D}_2\text{O}$ , 300 MHz, two rotamers 4:1):  $\delta$  0.88-1.88 (m, 13H), 2.25-2.42 (m, 1H), 2.63-2.89 (m, 1H), 3.94 (s, 2H), 3.99 (apparent doublet, 2H), 4.16 (t, 1H), 4.28 (q, 1H), 4.41 (q, 1H), 4.56 (s, 2H), 4.98 (dd, 1H), 7.53 (d, 2H), 7.77 (d, 2H),  
10

Resolved signals from the minor rotamer appears at  $\delta$  0.50 (bq), 0.77 (bq), 5.21 (dd), 7.56 (d) and 7.81 (d).  
15

$^{13}\text{C-NMR}$  (  $\text{D}_2\text{O}$ , 75 MHz ): The carbonyls and amidinecarbon at  $\delta$  166.8, 166.9, 168.6, 172.3 and 173.4.  
20

Resolved signals from the minor rotamer appears at  $\delta$ : 166.6, 169.6 and 172.0

#### Example 15

25

H-(R)Cha-Pro-Pab x 2 HCl

(i) Boc-(R)Cha-Pro-Pab(Z)

0.135 ml (1.1 mmol) pivaloyl chloride was added at room temperature to a stirred mixture of 0.155 ml (1.1 mmol) triethyl amine and 405 mg (1.1 mmol) Boc-(R)Cha-Pro-OH (See preparation of starting materials) in 5 ml DMF. After 3 h 340 mg (1.1 mmol) H-Pab(Z) (See preparation of  
30 starting materials) in 5 ml DMF was added and stirring was continued over night. The reaction mixture was diluted with water and extracted with ethyl  
35

acetate/toluene 1:1. The organic phase was dried ( $\text{MgSO}_4$ ) and the solvent was removed in vacuo to give a residue which was subjected to flash chromatography using ethyl acetate as eluent. The yield was 309 mg (49%).

5

(ii) H-(R)Cha-Pro-Pab(Z)

Hydrogen chloride was bubbled through a solution, until saturation, of 1.246 g (2 mmol) Boc-(R)Cha-Pro-Pab(Z) in 20 ml ethyl acetate at room temperature. After 30 minutes sodium carbonate solution (10%) was added and the organic phase which separated was dried ( $\text{K}_2\text{CO}_3$ ). The drying agent was washed with methylene chloride and the solvent was evaporated from the combined organic phases to give 1.11 g (100%) of the title compound.

10  
15

(iii) H-(R)Cha-Pro-Pab x 2 HCl

100 mg (0.19 mmol) H-(R)Cha-Pro-Pab(Z) dissolved in 15 ml ethanol was hydrogenated in the presence of 38 mg 10% Pd/C for 1.5 h. Dilution of the reaction mixture with distilled water and removal of the catalyst by filtration followed by removal of the ethanol in vacuo and freeze drying gave the title compound as a colorless powder. The peptide was finally converted to the dihydrochloride by dissolution in hydrochloric acid followed by freeze drying to give 90 mg (100%) of the title compound.

20  
25

$^1\text{H}$ -NMR (300 MHz,  $\text{D}_2\text{O}$ );  $\delta$  1.0-2.0 (m, 13H), 2.0-2.3 (m, 3H), 2.3-2.5 (m, 1H), 3.6-3.7 (m, 1H), 3.8-3.9 (m, 1H), 4.3-4.5 (t, 1H), 4.5-4.6 (m, 3H), 7.4-7.6 (m, 3H), 7.6-7.9 (m, 2H).

30

$^{13}\text{C}$ -NMR (75 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.2, 170.0, 174.9.

35

Example 16

HOOC-CH<sub>2</sub>-(R)Cha-Pro-Pab x 2 HCl

5 (i) BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z)

A mixture of 268 mg (0.5 mmol) H-(R)Cha-Pro-Pab(Z) (See Example 15), 90 µl (0.55 mmol) benzyl bromoacetate and 181 mg (1.3 mmol) K<sub>2</sub>CO<sub>3</sub> in 2 ml acetonitrile was sonicated at 40°C for 2.5 h. The mixture was filtered through hyflo and the solvent was removed in vacuo to give a residue which was subjected to flash chromatography using ethyl acetate as eluent to give 194 mg (57%) of the title compound.

15

HOOC-CH<sub>2</sub>-(R)Cha-Pro-Pab x 2 HCl

194 mg (0.28 mmol) BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z) dissolved in 10 ml ethanol was hydrogenated in the presence of 77 mg 10% Pd on charcoal for 3 h. The reaction mixture was diluted with water and the catalyst was removed by filtration. Evaporation of the ethanol in vacuo followed by freeze drying gave a white residue. Hydrochloric acid was added and the resulting solution was finally freeze dried to give 115 (68%) of the desired product.

<sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O); δ 1.0-1.2 (m, 2H), 1.2-1.5 (m, 3H), 1.5-2.0 (m, 8H), 2.0-2.3 (m, 3H), 2.3-2.5 (m, 1H), 3.6-3.8 (m, 1H), 3.8-4.0 (m, 3H), 4.4-4.7 (m, 4H), 7.5-7.7 (d, 2H), 7.7-7.9 (d, 2H).

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 167.1, 168.2, 169.3, 174.6.

35 Example 17

HOOC-CH<sub>2</sub>-(M)(R)Cha-Pr -Pab

## (i) Boc-(Me) (R)Cha-Pro-Pab(Z)

To a solution of 0.8 g (1.67 mmol) of Boc-(Me) (R)Cha-Pro-OSu (See preparation of starting materials) in 3 ml DMF was added a solution of 0.562 g (1.85 mmol) of H-Pab(Z) (See preparation of starting materials) in 3 ml of DMF, and the pH of the resulting solution was adjusted to 8-9 with N-methylmorpholine, whereafter the solution was stirred at room temperature for 2 days.

The solution was poured onto water, and the resulting mixture was extracted with 3x25 ml of ethyl acetate. The organic solution was washed with 1M KHSO<sub>4</sub> solution, 10% NaHCO<sub>3</sub> solution, water and brine, and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent gave 0.65 g (60%) of the title compound as a yellowish white powder.

## (ii) Me-(R)Cha-Pro-Pab(Z)

A solution of 0.60 g (0.92 mmol) of Boc-(Me) (R)Cha-Pro-Pab(Z) in 50 ml of EtOH was saturated with HCl at 0°C, and the solution was stored in refrigerator overnight. The resulting solution was evaporated to dryness, and the residue was dissolved in a Na<sub>2</sub>CO<sub>3</sub> solution, extracted with 3x25 ml ethyl acetate. The extract was washed with brine and evaporated to give 0.4 g (79%) of the compound as a white fluffy powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 0.8-1.0 (m, 2H), 1.1-1.4 (m, 5H), 1.4-1.55 (m, 1H), 1.6-1.9 (m, 10H), 1.9-2.05 (m, 2H), 2.05-2.2 (m, 2H), 2.19 (s, 3H), 2.4-2.5 (m, 1H), 3.28 (dd, 1H), 3.41 (q, 1H), 3.62 (m, 1H), 4.42 (m, 2H), 4.61 (d, 1H), 5.2 (s, 2H), 7.2-7.45 (m, 7H), 7.72 (t, 1H), 7.79 (d, 2H).

(iii) BnOOC-CH<sub>2</sub>-(Me) (R)Cha-Pro-Pab(Z)

A mixture of 0.40 g (0.73 mmol) of Me-(R)Cha-Pro-Pab(Z),

0.17 g  $\text{BnOOC-CH}_2\text{Br}$  and 0.20 g (2 equiv.) of  $\text{K}_2\text{CO}_3$  (mortared) in 15 ml of  $\text{CH}_3\text{CN}$  was stirred at room temperature overnight. The resulting mixture was evaporated, ethyl acetate was added, and the mixture was washed with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ), and evaporated. The crude product (0.69 g) was subjected to flash chromatography ( $\text{CH}_2\text{Cl}_2/\text{MeOH}$  10/1) yielding 0.39 g (77%) of a light yellow very viscous oil.

10  $\text{HOOC-CH}_2\text{-(Me) (R)Cha-Pro-Pab}$

To a solution of 0.39 g (0.56 mmol) of  $\text{BnOOC-CH}_2\text{-(Me) (R)Cha-Pro-Pab(Z)}$  in 30 ml of EtOH was added 0.1 g of Pd/C (10%), and the substance was hydrogenated at atmospheric pressure. The solution was filtered and evaporated, whereafter the remaining syrupy material was freeze dried to yield 0.25 g (95%) of the compound as a white crystalline powder.

20  $^1\text{H-NMR}$  (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  0.85-1.1 (m, 2H), 1.1-1.4 (m, 6H), 1.5-1.85 (m, 9H), 1.9-2.05 (m, 3H), 2.05-2.15 (m, 1H), 2.15-2.3 (m, 1H), 2.57 (s, 3H), 3.32 (d, 1H), 3.55-3.75 (m, 2H), 3.95-4.1 (m, 2H), 4.35-4.5 (m, 3H), 7.55 (d, 2H), 7.72 (d, 2H).

25  $^{13}\text{C-NMR}$  (75 MHz,  $\text{CD}_3\text{OD}$ ): amidine and carbonyl carbons:  $\delta$  168.4, 171.5, 174.7, 175.1.

Example 18

30  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pro-Pab} \times 2 \text{ HCl}$

(i)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pro-Pab(Z)}$

35 A mixture of 149 mg (0.28 mmol)  $\text{H-(R)Cha-Pro-Pab(Z)}$  (See Example 15) and 66 mg (0.4 mmol) benzyl acrylate in 1.5 ml ethanol was kept at room temperature for 36 h. The

solvent was removed in vacuo and the residue was subjected to flash chromatography using ethyl acetate as eluent to give 124 mg (64%) of the desired product.

5 (ii)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pro-Pab x 2 HCl}$

124 mg (0.18 mmol)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pro-Pab(Z)}$  dissolved in 10 ml ethanol was hydrogenated for 1 h in the presence of 55 mg 10% Pd/C. The catalyst was removed by filtration and the solvent was removed in vacuo. The residue was dissolved in hydrochloric acid and the resulting solution was freeze dried to give 87 mg (79%) of the title compound.

15  $^1\text{H-NMR}$  (300 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  1.0-2.0 (m, 13H), 2.0-2.2 (m, 3H), 2.2-2.4 (m, 1H), 2.7-2.8 (t, 2H), 3.2-3.3 (m, 1H), 3.3-3.4 (m, 1H), 3.5-3.7 (m, 1H), 3.7-3.9 (m, 1H), 4.3-4.6 (m, 4H), 7.4-7.6 (m, 2H), 7.7-7.8 (m, 2H).

20  $^{13}\text{C-NMR}$  (75 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.0, 168.3 and 174.6 (Two carbons are overlapping).

Example 19

25  $\text{HOOC-CH}_2\text{-CH}_2\text{-(Me) (R)Cha-Pro-Pab x 2 HCl}$

(i)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(Me) (R)Cha-Pro-Pab(Z)}$

To a solution of 274 mg (0.5 mmol) of  $\text{Me-(R)Cha-Pro-Pab(Z)}$  (See Example 17) in 5 ml of EtOH (99%) was added 97.3 mg (0.6 mmol) of benzyl acrylate and the reaction was stirred at room temperature. After 72 h an additional amount of 16.2 mg (0.1 mmol) of benzyl acrylate was added and the stirring continued for 24 h. The solvent was evaporated and the residue was subjected to flash chromatography ( $\text{CH}_2\text{Cl}_2/\text{MeOH}(\text{NH}_3\text{-saturated})$ , 95/5) to give 198 mg (56%) of the title compound.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.8-2.0 (several m, 16 H), 2.14 (s, 3H), 2.24-2.33 (m, 2H), 2.38-2.46 (m, 1H), 2.67 (t, 2H), 3.32-3.40 (m, 2H), 3.71 (m, 1H), 4.36-4.44 (m, 2H), 4.58 (m, 1H), 5.03 (apparent s, 2H), 5.20 (s, 2H), 7.25-7.37 (m, 10H), 7.43 (d, 2H), 7.64 (t, 1H (NH)), 7.81 (d, 2H).

<sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 164.7, 167.9, 171.7, 172.3 and 172.6.

(ii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(Me)(R)Cha-Pro-Pab x 2 HCl

To a solution of 198 mg (0.27 mmol) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(Me)(R)Cha-Pro-Pab(Z) in 10 ml EtOH and 1 ml 1M HCl was added 60 mg of 5 % Pd/C (containing 50 % H<sub>2</sub>O by weight) and the mixture was hydrogenated at atmospheric pressure for 4 h. The catalyst was filtered off and the solvent was evaporated. The remaining oil was dissolved in water and freeze dried to give the title compound in a quantitative yield.

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O): δ 1.08-1.2 (m, 2H), 1.2-1.42 (m, 4H), 1.68-1.91 (m, 5H), 1.93-2.08 (m, 2H), 2.09-2.26 (m, 3H), 2.49 (m, 1H), 2.95 (m, 2H), 3.03 (s, 3H), 3.60 (apparent bs, 2H), 3.82 (m, 1H), 3.98 (m, 1H), 4.53 (m, 1H), 4.61 (bs, 2H), 4.64 (m, 1H), 7.63 (d, 2H), 7.97 (d, 2H).

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 167.2, 167.8 and 174.5. Two peaks are probably overlapping.

#### Example 20

HOOC-CH<sub>2</sub>-(RorS)CH(COOH)-(R)Cha-Pro-Pab/a x 2 HCl

(i) BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Pro-Pab(Z)

A mixture of 0.50 g (0.94 mmol) of H-(R)Cha-Pro-Pab(Z) (See Example 15) and 0.28 g (0.94 mmol) of dibenzyl maleate in 20 ml of EtOH was kept at room temperature for 5 days. Evaporation of the solvent followed by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH as eluent gave 0.15 g (19 %) of the diastereomeric mixture.

<sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 0.7-2.1 (m, 17 H), 2.3-2.4 (m, 1 H), 2.5-2.8 (m, 2 H), 3.2-3.7 (m, 4 H), 4.46 (d, 1 H), 4.65 (bd, 1 H), 4.81 (d, 1 H), 4.9-5.1 (m, 3 H), 5.20 (s, 2 H), 7.1- 7.4 (m, 15 H), 7.4-7.5 (m, 2 H), 7.6-7.8 (m, 3 H).

(ii) HOOC-CH<sub>2</sub>-(RorS)CH(COOH)-(R)Cha-Pro-Pab/a x 2 HCl

A mixture of 0.15 g (0.18 mmol) of BnOOC-CH<sub>2</sub>-(R,S)CH(COOBn)-(R)Cha-Pro-Pab(Z) was dissolved in 5 ml of ethanol and was hydrogenated over 5% Pd/C at atmospheric pressure for 1 h. to give HOOC-CH<sub>2</sub>-(R,S)CH(COOH)-(R)Cha-Pro-Pab. The two diastereomers were separated by RPLC using (CH<sub>3</sub>CN/0.1 M NH<sub>4</sub>OAc 15/85) as eluent followed by freeze drying from HCl. This diastereomer eluted first from the column. Yield 19 mg (18 %).

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O, mixture of two rotamers) major rotamer: δ 1.0-2.0 (m, 15H), 2.15 (m, 2H), 2.44 (m, 1H), 3.00 (bd, 1H), 3.05 (bd, 1 H), 3.69 (m, 1H), 3.84 (m, 1H), 3.97 (bs, 1H), 4.5-4.7 (m, 3H), 7.62 (d, 2H), 7.87 (d, 2H).

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 167.2, 168.3, 173.8, 174.6 and 178.2.

#### Example 21

35

HOOC-CH<sub>2</sub>-(R rS)CH(COOH)-(R)Cha-Pr -Pab/b x 2 HCl



The title compound was obtained by using the same procedure as described in Example 20 on  $\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Cha-Pro-Pab}$ . This diastereomer came out after the first one from the column. Yield 19 mg (18 %).

5

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ , mixture of two rotamers) major rotamer:  $\delta$  1.0-2.0 (m, 14H), 2.15-2.25 (m, 3H), 2.44 (m, 1H), 3.11 (bd, 1H), 3.19 (bd, 1H), 3.71 (m, 1H), 3.92 (m, 1H), 4.03 (bs, 1H), 4.5-4.7 (m, 3H), 7.58 (d, 2H), 7.84 (d, 2H).

10

Resolved signals arising from the minor rotamer appears at:  $\delta$  7.66 (d) and 7.91 (d).

15

$^{13}\text{C-NMR}$  (75 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.3, 168.5 and 174.7. Two carbons are probably overlapping.

#### Example 22

20

$\text{HOOC-CH}_2\text{-NH-CO-CH}_2\text{-(R)Cha-Pro-Pab} \times 2 \text{ HCl}$

(i)  $\text{BnOOC-CH}_2\text{-NH-CO-CH}_2\text{-(R)Cha-Pro-Pab(Z)}$

25

0.246 g (0.460 mmole) of  $\text{H-(R)Cha-Pro-Pab(Z)}$  (See Example 15), 0.140 g (1.01 mmole)  $\text{K}_2\text{CO}_3$  and 0.145 g (0.506 mmole)  $\text{BnOOC-CH}_2\text{-NH-CO-CH}_2\text{-Br}$  (See preparation of starting materials) was mixed in 6 ml acetonitrile. The mixture was stirred at  $50^\circ\text{C}$  for 2 h 30 minutes, the solvent was evaporated and the residue was partitioned between water and ethyl acetate. The layers were separated and the water layer was extracted one more time with ethyl acetate. The combined organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to yield 0.350 g of an oil. The crude product was purified by flash chromatography using a stepwise gradient of  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  97/3, 95/5, 92.5/7.5 to yield 0.227 g (67%) of the title compound.

30

35

<sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): δ 25.0, 26.0, 26.2, 26.4, 26.7, 32.4, 34.2, 34.4, 40.8, 40.9, 42.9, 46.7, 50.5, 58.4, 60.2, 67.0, 67.2, 127.5, 127.8, 128.2, 128.3, 128.4, 128.5, 128.6, 128.6, 134.1, 135.2, 137.0, 142.6, 164.7, 168.9, 169.3, 170.4, 172.2, 175.0

(ii) HOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R)Cha-Pro-Pab x 2 HCl

0.089 g (0.12 mmole) BnOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z) was mixed with 30 mg 5 % Pd/C and dissolved in 10 ml acetic acid. The mixture was hydrogenated at atmospheric pressure for one and a half hour. Filtration of the catalyst through hyflo and freeze drying with 1ml 1N hydrochloric acid gave 0.058 g (82%) of the desired product.

<sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 0.9-2.2 (m, 16H), 2.25-2.47 (m, 1H), 3.55-3.7 (m, 1H), 3.7-4.1 (m, 5H), 4.42 (t, 1H), 4.48-4.6 (m, 3H), 7.51 (d, 2H), 7.77 (d, 2H)

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 166.8, 167.1, 168.2, 173.6 and 174.6

### Example 23

EtOOC-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pab x HOAc

(i) EtOOC-CH=CH-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z)

H-(R)Cha-Pro-Pab(Z) (See Example 15) (275 mg, 0.51 mmol) was treated with K<sub>2</sub>CO<sub>3</sub> (141 mg, 1.02 mmol) and BrCH<sub>2</sub>CH=CHCOOEt (108 mg, 0.56 mmol) in CH<sub>3</sub>CN (10 ml) at 20°C for 20 h. The solvent was evaporated and the residue was dissolved in EtOAc (5 ml)/H<sub>2</sub>O (2 ml). The organic layer was separated, dried (Na<sub>2</sub>SO<sub>4</sub>), and

concentrated yielding 397 mg of an oil which was purified by flash chromatography using EtOAc/hexane, 1/4 as eluent to give 252 mg (77%) of the title compound.

5

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 0.8-1.05 (m, 2H), 1.1-1.45 (m, 3H), 1.3 (t, 3H), 1.5-1.9 (m, 8H), 1.95-2.05 (m, 1H), 2.1-2.15 (m, 1H), 2.45-2.55 (m, 1H), 3.0 and 3.15 (two d, 2H), 3.35-3.45 (m, 2H), 3.55-3.65 (m, 1H), 4.15 (q, 2H), 4.3 (d, 1H), 4.6-4.7 (m, 2H), 5.2 (s, 2H), 5.85 (d, 1H), 6.75 (dt, 1H), 5.3-5.4 (m, 4H), 7.45 (d, 2H), 7.85 (d, 2H).

10

<sup>13</sup>C-NMR (75.0 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 165.7, 171.2 and 175.7 (two peaks are probably overlapping).

15

(ii) EtOOC-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pab x HOAc

20

EtOOCCH=CHCH<sub>2</sub>-(R)Cha-Pro-Pab(2) (250 mg, 0.38 mmol) was dissolved in ethanol and hydrogenated in the presence of 5 % Pd/C during approximately 2 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave after purification by RPLC using (CH<sub>3</sub>CN/0.1 M NH<sub>4</sub>OAc) as eluent 70 mg (36%) of the desired product.

25

<sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD): δ 0.9-1.05 (m, 2H), 1.15-1.55 (m, 5H), 1.25 (t, 3H), 1.6-1.85 (m, 7H), 1.95-2.6 (m, 8H), 3.55-3.65 (m, 2H), 3.8 (m, 1H), 4.1 (q, 2H), 4.45 (m and d, 2H), 4.55 (d, 1H), 7.55 and 7.75 (two d, 4H).

30

<sup>13</sup>C-NMR (75.0 MHz, CD<sub>3</sub>OD): amidine and carbonyl carbons: δ 168.3, 173.2, 174.6 and 174.9.

35

Example 24**Ph(4-COOH)-SO<sub>2</sub>-(R)Cha-Pro-Pab x HCl****5 (i) Ph(4-COOH)-SO<sub>2</sub>-(R)Cha-Pro-Pab(Z)**

64 mg (0.32 mmol) 4-chlorosulfonyl-benzoic acid was added at ice bath temperature to a solution of 156 mg (0.29 mmol) H-(R)Cha-Pro-Pab(Z) (See Example 15) and 59 mg  
10 (0.58 mmol) triethyl amine in 4 ml methylene chloride. The mixture was slowly allowed to reach room temperature and after 24 hours it was washed with water and dried (Na<sub>2</sub>SO<sub>4</sub>). Removal of the solvent in vacuo and purification of the residue by flash chromatography using ethyl  
15 acetate/methanol 9:1 followed by methylene chloride/methanol 3:1 as eluents gave 82 mg (39%) of the product.

**20 (ii) Ph(4-COOH)-SO<sub>2</sub>-(R)Cha-Pro-Pab x HCl**

80 mg (0.11 mmol) Ph(4-COOH)-SO<sub>2</sub>-(R)Cha-Pro-Pab(Z) was hydrogenated over 5 % Pd/C in EtOH. The catalyst was filtered off, the solvent evaporated and the crude  
25 product was purified by RPLC using (CH<sub>3</sub>CN/0.1 M NH<sub>4</sub>OAc 1/4) as eluent and finally converted to the hydrochloride salt by freeze drying from HCl which gave 21 mg (29%) of the product.

30 <sup>1</sup>H-NMR (300 MHz, CD<sub>3</sub>OD, mixture of two rotamers): δ 0.45-1.82 (m, 13H), 1.90-2.30 (m, 4H), 2.95-4.16 (several m, total 3H), 4.35-4.68 (m, 3H), 7.54 (d, 2H), 7.74 (d, 1H), 7.80 (d, 1H), 7.90-8.00 (m, 2H), 8.05-8.22 (m, 2H)

35 <sup>13</sup>C-NMR (75 MHz, CD<sub>3</sub>OD): amidine and carbonyl carbons: δ 168.4, 173.4, 173.9 and 174.2

MS m/z 584 ( $M^+ + 1$ )

Example 25

5 H-(R)Cha-Pic-Pab x 2 HCl

(i) Boc-(R)Cha-Pic-Pab(Z)

10 3.57 g (18.6 mmol) EDC was added at  $-15^{\circ}\text{C}$  to a mixture of  
7.11 g (18.6 mmol) Boc-(R)Cha-Pic-OH (See preparation of  
starting materials), 9.07 g (74.2 mmol) DMAP and 5.26 g  
15 (18.6 mmol) H-Pab(Z) (See preparation of starting  
materials) in 200 ml DMF. The temperature was allowed to  
rise to  $20^{\circ}\text{C}$  over night. The solvent was removed in vacuo  
and toluene and water was added. The organic phase was  
washed with water, 1M  $\text{KHSO}_4$ , 10%  $\text{Na}_2\text{CO}_3$  and brine. Drying  
( $\text{MgSO}_4$ ) and evaporation of the solvent in vacuo gave 13.63  
20 g of a residue which was subjected to flash chromatography  
on silica gel using ethyl acetate/toluene 2:1 as  
eluent to give 9.5 g (79%) of the title compound.

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.7-1.0 (m, 2H), 1.0-2.2 (m,  
25H), 2.3-2.5 (m, 1H), 2.9-3.1 (m, 1H), 3.8 (d, 1H), 4.3  
(dd, 1H), 4.4-4.6 (m, 2H), 5.1 (s, 2H), 5.1-5.3 (m, 2H),  
25 7.2-7.3 (m, 5H), 7.35 (d, 2H), 7.4-7.5 (m, 1H), 7.75 (d,  
2H).

$^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$   
156.8, 164.6, 168.2, 170.0 and 173.4.

30

(ii) H-(R)Cha-Pic-Pab(Z)

Hydrogen chloride was bubbled through a solution of 9.5  
g (14.7 mmol) Boc-(R)Cha-Pic-Pab(Z) in 100 ml ethyl  
35 acetate at room temperature until saturation. After 10  
minutes  $\text{Na}_2\text{CO}_3$  solution (10%) was added and the organic  
phase which separated was dried ( $\text{K}_2\text{CO}_3$ ) and the solvent

was removed in vacuo to give the title compound in quantitative yield.

5  $^1\text{H-NMR}$  (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  0.85-1.05 (m, 2H), 1.15-1.90 (m, 16H), 2.25-2.35 (m, 1H), 3.20-3.30 (m, 1H), 3.80-3.90 (d, 1H), 3.90-4.0 (m, 1H), 4.4-4.5 (two d, 2H), 4.7 (br s, 5H) 5.15 (s, 2H), 5.20 (m, 1H), 7.25-7.45 (m, 7H), 7.85 (d, 2H).

10 (iii) H-(R)Cha-Pic-Pab x 2 HCl

55 mg (0.1 mmol) H-(R)Cha-Pic-Pab(Z) dissolved in a mixture of 5 ml ethanol and 0.45 ml 1M hydrochloric acid was hydrogenated in the presence of 33 mg 10% Pd/C for 15 1.5 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave a residue which was subjected to RPLC using 0.1 M  $\text{NH}_4\text{OAc}/\text{CH}_3\text{CN}$  as eluent. The purified peptide was finally converted to the dihydrochloride salt by dissolution in hydrochloric acid 20 followed by freeze drying. The yield was 17 mg (35%) of the title compound

25  $^1\text{H-NMR}$  (300 MHz,  $\text{D}_2\text{O}$ , 2 rotamers, 3:1 mixture):  $\delta$  1.0-2.0 (m, 18H), 2.33 (d, 1H), 3.4-3.5 (m, 1H), 3.8-3.9 (m, 1H), 4.4-4.8 (m, 3H), 5.15-5.25 (m, 1H), 7.5-7.7 (m, 2H), 7.8-8.0 (m, 2H).

Resolved signals from the minor rotamer appears at  $\delta$ : 0.5-0.7 (m) and 3.0-3.1 (m)

30

$^{13}\text{C-NMR}$  (75 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.3, 171.6 and 173.6.

Resolved signals for the minor rotamer appears at  $\delta$  170.6 and 172.4.

35

Example 26

HOOC-CH<sub>2</sub>-(R)Cha-Pic-Pab x 2 HCl

5 (i) BnOOC-CH<sub>2</sub>-(R)Cha-Pic-Pab(Z)

10 A mixture of 742 mg (1.35 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25), 230 ml (1.45 mmol) benzyl bromoacetate and 558 mg (4 mmol) K<sub>2</sub>CO<sub>3</sub> in 4 ml acetonitrile was sonicated at 40°C for 40 minutes. The solvent was removed and the residue was subjected to flash chromatography to give 720 mg (77%) of the desired product.

15 <sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>); δ 0.8-1.0 (m, 2H), 1.1-1.9 (m, 16H), 2.1-2.4 (br s, 1 or 2H), 2.4 (d, 1H), 3.0 (m, 1H), 3.25 (d, 1H), 3.45 (d, 1H), 3.55-3.65 (m, 1H), 3.7 (m, 1H), 4.35 (dd, 1H), 4.55 (dd, 1H), 4.80 (two d, 2H), 5.2 (s, 2H), 5.3 (m, 1H), 7.2-7.4 (m, 12H), 7.8 (d, 2H).

20 <sup>13</sup>C-NMR (125 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 164.5, 167.9, 170.5, 173.4 and 175.5.

(ii) HOOC-CH<sub>2</sub>-(R)Cha-Pic-Pab x 2 HCl

25 509 mg (0.73 mmol) BnOOC-CH<sub>2</sub>-(R)Cha-Pic-Pab(Z) dissolved in 25 ml ethanol was hydrogenated in the presence of 259 mg 10% Pd/C for 4 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo gave a residue which was dissolved in distilled water.  
30 Hydrochloric acid was added and the solution was finally freeze dried to give 281 mg (79%) of the title compound.

35 <sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O, mixture of rotamers 4:1): major rotamer: δ 1.0-2.0 (m, 18H), 2.25-2.40 (m, 1H), 3.4-3.5 (m, 1H), 3.8-3.95 (m, 3H), 4.55-4.65 (two d, 2H), 5.15 (m, 1H), 7.55-7.75 (m, 2H), 7.8-8.0 (m, 2H).

$^{13}\text{C}$ -NMR (125 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.3, 169.9, 170.3 and 173.5.

Resolved signal for the minor rotamer appears at  $\delta$  166.9, 169.2 and 172.0

5

Example 27

$\text{HOOC-CH}_2\text{-(RorS)CH(COOH)-(R)Cha-Pic-Pab/a} \times 2 \text{ HCl}$

10 (i)  $\text{BnOOC-CH}_2\text{-(R,S)CH(COOBn)-(R)Cha-Pic-Pab(Z)}$

A mixture of 592 mg (1.1 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25) and 332 mg (1.1 mmol) dibenzyl maleate in 1 ml ethanol was kept at room temperature for 1 week. The solvent was removed in vacuo and the residue was subjected to flash chromatography using methanol/methylene chloride as eluent to give 275 mg (30%) the diastereomeric mixture.

15

20 (ii)  $\text{HOOC-CH}_2\text{-(RorS)CH(COOH)-(R)Cha-Pic-Pab/a} \times 2 \text{ HCl}$

275 mg  $\text{BnOOC-CH}_2\text{-(R,S)CH(COOBn)-(R)Cha-Pic-Pab(Z)}$  dissolved in 20 ml 95% ethanol was hydrogenated for 18 hours in the presence of 75 mg 10% Pd/C. The mixture was filtered through hyflo and the solvent was removed in vacuo. Addition of water followed by freeze drying gave 166 mg of  $\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Cha-Pic-Pab}$ . The two diastereomers were separated by RPLC using ( $\text{CH}_3\text{CN}/0.1 \text{ M NH}_4\text{OAc } 1/4$ ) as eluent followed by freeze drying from HCl. This diastereomer eluted first from the column. Yield 9 mg.

$^1\text{H}$ -NMR (300 MHz,  $\text{D}_2\text{O}$ , mixture of rotamers):  $\delta$  1.0-2.0 (m, 18H), 2.25-2.4 (m, 1H), 3.0-3.2 (m, 2H), 3.4 (t, 1H), 3.8 (d, 1H), 4.05 (t, 1H), 4.5-4.7 (m, 3H), 5.2 (s, 1H), 7.55 (d, 2H), 7.9 (d, 2H).

35



Resolved signals from the minor rotamer appears at  $\delta$  4.0(t) and 7.7(d).

Example 28

5

$\text{HOOC-CH}_2\text{-(RorS)CH(COOH)-(R)Cha-Pic-Pab/b} \times 2 \text{ HCl}$

The title compound was obtained by using the same procedure as described in Example 27 on  $\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Cha-Pic-Pab}$ . This diastereomer came out after the first one from the column.

10

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ , mixture of rotamers;  $\delta$  1.0-2.0 (m, 18H), 2.25-2.4 (m, 1H), 3.0-3.2 (m, 2H), 3.5 (t, 1H), 3.85 (d, 1H), 4.15 (s, 1H), 4.5-4.7 (m, 3H), 5.15 (s, 1H), 7.55 (d, 2H), 7.8 (d, 2H).

15

Resolved signals from the minor rotamer appear at  $\delta$  4.35(s), 7.65(d) and 7.9(d).

20

Example 29

$\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pic-Pab} \times 2 \text{ HCl}$

(i)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pic-Pab(Z)}$

25

A mixture of 851 mg (1.55 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25) and 269 mg (1.71 mmol) benzyl acrylate in 5 ml ethanol was kept at room temperature for 40 h. The solvent was removed in vacuo and the residue was subjected to flash chromatography using methylene chloride/methanol as eluent to give 812 mg (74%) of the product.

30

$^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.8-1.0 (m, 2H), 1.1-1.9 (m, 16H), 2.3-2.5 (m, 3H), 2.6-2.8 (m, 2H), 3.0 (m, 1H), 3.5 (m, 1H), 3.6-3.7 (m, 1H), 4.3 (dd, 1H), 4.6 (dd, 1H),

35

4.95-5.05 (two d, 2H), 5.2 (s, 2H), 5.3 (m, 1H), 6.5-6.9 (br s, 1H), 7.0-7.1 (m, 1H), 7.2-7.5 (m, 12H), 7.75-7.85 (d, 2H), 9.3-9.7 (br s, 1H).

5 (ii)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pic-Pab} \times 2 \text{ HCl}$  :

780 mg (1.1 mmol)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Pic-Pab(Z)}$  dissolved in 25 ml ethanol was hydrogenated for 4 h in the presence of 306 mg 15% Pd/C. The catalyst was removed  
10 by filtration and the solvent was removed in vacuo. The residue was dissolved in hydrochloric acid and the resulting solution was freeze dried to give 481 mg (78%) of the title compound.

15  $^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  0.95-1.1 (m, 2H), 1.15-1.9 (m, 16H), 2.2-2.3 (m, 1H), 2.7-2.8 (t, 2H), 3.2-3.3 (m, 3H), 3.4-3.5 (m, 1H), 3.75-3.85 (m, 1H), 4.4-4.6 (m, 3H), 5.15 (m, 1H), 7.5-7.6 (m, 2H), 7.8-7.9 (m, 2H), 8.6-8.7 (m, 1H).

20  $^{13}\text{C-NMR}$  (125 MHz,  $\text{CD}_3\text{OD}$ ): amidine and carbonyl carbons:  $\delta$  170.6, 175.9, 179.5 and 183.5.

Example 30

25

$\text{HOOC-CO-(R)Cha-Pic-Pab} \times \text{HOAc}$

(i)  $\text{EtOOC-CO-(R)Cha-Pic-Pab(Z)}$

30 0.12 g ethyloxalyl chloride was added to a mixture of 0.42 g (0.77 mmol)  $\text{H-(R)Cha-Pic-Pab(Z)}$  (See Example 25) and 0.21 g (1.5 mmol)  $\text{K}_2\text{CO}_3$  in 10 ml  $\text{CH}_3\text{CN}$  at room temperature. After 2 hours an additional amount of 0.07 g (0.5 mmol) ethyloxalyl chloride was added. The mixture  
35 was stirred at room temperature over night. The solvent was removed in vacuo. and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$  and washed with water. Evaporation and flash

chromatography (toluene: ethyl acetate 1: 2 followed by  $\text{CH}_2\text{Cl}_2$ : methanol) gave 0.21 g (42%) of the product.

(ii)  $\text{HOOC-CO-(R)Cha-Pic-Pab(Z)}$

5

0.21 g (0.32 mmol)  $\text{EtOOC-CO-(R)Cha-Pic-Pab(Z)}$  was dissolved in 3 ml THF and 0.17 g (4.2 mmol) LiOH dissolved in 3 ml water was added. The mixture was stirred at room temperature over night and then poured onto ethyl acetate/water. The phases were separated and the organic phase was extracted with a  $\text{KHCO}_3$ -solution. The aqueous phase was acidified with 0.5M HCl (pH 1) and extracted with  $\text{CH}_2\text{Cl}_2$ , dried over  $\text{Na}_2\text{SO}_4$  and evaporated to give 80 mg of the product.

15

(iii)  $\text{HOOC-CO-(R)Cha-Pic-Pab} \times \text{HOAc}$

$\text{HOOC-CO-(R)Cha-Pic-Pab(Z)}$  was hydrogenated over 5 % Pd/C in EtOH. The catalyst was filtered off and the solvent evaporated. The residue was subjected to purification by RPLC to give the title compound.

20

$^1\text{H NMR}$  (500 MHz,  $\text{DMSO-d}_6$ );  $\delta$  0.8-1.0 (m, 2H), 1.1-1.75 (m, 15H), 1.86-1.94 (m, 1H), 2.13-2.2 (m, 1H), 3.75-3.81 (m, 1H), 4.32, 4.44 (AB, 2H), 4.71-4.77 (m, 1H), 4.98-5.02 (m, 1H), 7.41 (d, 2H), 7.75 (d, 2H), 8.1-8.15 (m, 1H), 8.22-8.27 (m, 1H), 9.32 (broad s), 9.90 (broad s). The signal of one of the protons (3.25) is partially obscured by the solvent signal.

30

MS  $m/z$  486 ( $\text{M}^+ + 1$ )

### Example 31

35

$\text{HOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab}$

(i)  $\text{MeOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab(Z)}$

0.39 g (0.72 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25) and 0.9 g (0.8 mmol) monomethylmalonate was dissolved in 40 ml  $\text{CH}_2\text{Cl}_2$  and 0.16 g (0.8 mmol) DCC was added. The solution was stirred in room temperature over night. The precipitated DCU was removed by filtration and the filtrate was washed with 0.3M  $\text{KHSO}_4$  and  $\text{KHCO}_3$ -solution and dried ( $\text{NaSO}_4$ ). Evaporation of the solvent followed by flash chromatography using toluen/ethyl acetate (1/3) as eluent gave 0.27 g (58%) of the desired product.

(ii)  $\text{MeOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab}$

90 mg (0.14 mmol)  $\text{MeOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab(Z)}$  was dissolved in 10 ml ethanol and was hydrogenated in presence of 5% Pd/C for 5 hours. Removal of the catalyst by filtration and evaporation of the solvent gave 50 mg (70%) of the title product.

$^1\text{H}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  0.85-1.1 (m, 2H), 1.1-1.9 (m, 16H), 2.35-2.45 (m, 1H), 3.2-3.4 (m, 3H), 3.7 (s, 3H), 3.95-4.05 (m, 1H), 4.4-4.55 (m, 3H), 5.15-5.25 (m, 1H), 7.4-7.55 (m, 2H), 7.7-7.85 (m, 2H).

$^{13}\text{C}$  NMR (75 MHz,  $\text{CD}_3\text{OD}$ ): amidine and carbonyl carbons:  $\delta$  168.2, 168.7, 170.0, 172.4 and 174.6.

MS m/z 514 ( $\text{M}^+ + 1$ )

(iii)  $\text{HOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab}$

To a solution of 0.14 g (0.27 mmol) of  $\text{MeOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab}$  in 5 ml methanol was added 2 ml of 0.5 M NaOH at room temperature. After stirring for 5 hours water was added and the methanol was removed in vacuo. The aqueous phase was freeze dried. The soluble material was extracted out from the insoluble inorganic salts with

absolute ethanol. The remaining solid after evaporation of the ethanol was suspended in water and 70 mg (52%) of the title compound was isolated by filtration.

5  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO-d}_6$ ):  $\delta$  0.8-1.0 (m, 2H), 1.0-1.9 (m, 16H), 2.15-2.30 (m, 1H), 2.58, 2.86 (AB, 2H), 3.8-3.95 (m, 1H), 4.2-4.5 (m, 2H), 4.7-4.85 (m, 1H), 4.95-5.05 (m, 1H), 7.40 (d, 2H), 7.77 (d, 2H), 8.2-8.3 (m, 1H), 9.3-9.4 (m, 1H), 9.90 (broad s, 3H). The signal of one of the  
10 protons (3.21) is partially obscured by the solvent-signal.

$^{13}\text{C}$  NMR (75 MHz,  $\text{DMSO-d}_6$ ): amidine and carbonyl carbons:  
15  $\delta$  165.8, 168.8, 169.9, 172.2 and 172.4.

MS  $m/z$  500 ( $\text{M}^+ + 1$ )

#### Example 32

20  $\text{MeOOC-CH}_2\text{-CO-(R)Cha-Pic-Pab}$

See Example 31 (ii) above.

#### Example 33

25

$\text{H}_2\text{N-CO-CH}_2\text{-(R)Cha-Pic-Pab}$

(i)  $\text{H}_2\text{N-CO-CH}_2\text{-(R)Cha-Pic-Pab(Z)}$

30 Attempted alkylation of 455 mg (0.83 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25) with 80 mg (0.86 mmol) chloroacetamide in 3 ml acetonitrile in the presence of 395 mg (2.86 mmol) potassium carbonate by sonication at 40°C turned out to be an extremely sluggish reaction. Even  
35 the addition of 230 mg (2.6 mmol) lithium bromide did not seem to improve the reaction rate. However, addition of lithium iodide and heating/sonication gave small amounts

of product, according to TLC. Workup by addition of water, extraction with ethyl acetate/toluene, drying of the organic phase ( $\text{MgSO}_4$ ) and removal of the solvent in vacuo gave a residue which was subjected to flash chromatography using  $\text{MeOH}/\text{CH}_2\text{Cl}_2$  as eluent to give 118 mg (24%) of the desired product.

(ii)  $\text{H}_2\text{N}-\text{CO}-\text{CH}_2-(\text{R})\text{Cha-Pic-Pab} \times 2 \text{ HCl}$

118 mg (0.2 mmol)  $\text{H}_2\text{N}-\text{CO}-\text{CH}_2-(\text{R})\text{Cha-Pic-Pab}(\text{Z})$  dissolved in 10 ml 95% ethanol was hydrogenated in the presence of 143 mg 10% Pd/C for 2 h. The mixture was diluted with distilled water and hydrochloric acid and filtered through hyflo. Freeze drying gave 26 mg (24%) of the desired product.

$^1\text{H-NMR}$  (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  0.9-1.1 (m, 2H), 1.1-1.9 (m, 16H), 2.3 (d, 1H), 3.4 (t, 1H), 3.6 (AB-system, 2H), 3.8 (d, 2H), 4.35 (t, 1H), 4.5 (s, 2H), 5.2 (s, 1H), 7.55 (d, 2H), 7.8 (d, 2H).

#### Example 34

##### **Boc-(R)Cha-Pic-Pab**

10 mg (0.015 mmol) Boc-(R)Cha-Pic-Pab(Z) (See Example 25) dissolved in 5 ml ethanol was hydrogenated in the presence of 38 mg 10% Pd/C for 4 h. Removal of the catalyst by filtration and evaporation of the solvent in vacuo followed by dissolution of the residue in water and freeze drying yielded 7.6 mg (95%) of the product.

$^1\text{H-NMR}$  (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  0.9-1.1 (m, 2H), 1.1-1.9 (m, 16H), 2.4 (d, 1H), 3.25 (t, 1H), 4.0 (d, 1H), 4.5 (AB-system, 2H), 4.5-4.6 (m, 1H), 5.25 (s, 1H), 7.45 (d, 2H), 7.75 (d, 2H).

Example 35**Ac-(R)Cha-Pic-Pab x HCl**5      **(i) Ac-(R)Cha-Pic-Pab(Z)**

Acetyl chloride 0.06 g (0.8 mmol) was added to a mixture of 0.37 g (0.68 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25) and 0.19 g (1.35 mmol)  $K_2CO_3$  in 10 ml  $CH_3CN$  at room temperature. After stirring for an additional 30 minutes at room temperature the solvent was removed in vacuo. The residue was dissolved in  $CH_2Cl_2$  and washed with water. Evaporation and flash chromatography using a stepwise gradient of  $CH_2Cl_2$ / MeOH ( 99.9/0.1, 99.8/0.2, 99.6/0.4, 99.2/0.8 and 98.4/1.6) gave 0.24 g (60%) of the product.

**(ii) Ac-(R)Cha-Pic-Pab x HCl**

Ac-(R)Cha-Pic-Pab(Z) was hydrogenated over 5 % Pd/C at atmospheric pressure. After filtration of the catalyst and evaporation of the solvent the crude material was subjected to purification by RPLC using  $CH_3CN$ /0.1 M  $NH_4OAc$  (35/65) as eluent. Removal of the solvent and excess  $NH_4OAc$  followed by freeze drying from 1M HCl gave the title compound.

$^1H$ -NMR (300 MHz,  $CD_3OD$ ):  $\delta$  0.85-1.1 (m, 2H), 1.15-2.0 (m, 19H), 2.35-2.47 (m, 1H), 3.2-3.33 (m, 1H), 3.95-4.05 (m, 1H), 4.46, 4.57 (ABX, 2H), 5.16-5.22 (m, 1H), 7.51 (d, 2H), 7.76 (d, 2H), 8.23 (m, 1H). The signal of one of the protons is totally obscured by the solvent-signal.

$^{13}C$ -NMR (75 MHz,  $CD_3OD$ ): amidine and carbonyl carbons:  $\delta$  168.3, 172.5, 173.8, 175.1

MS m/z 456 ( $M^+ + 1$ )

Example 36**Me-SO<sub>2</sub>-(R)Cha-Pic-Pab x HCl**5      (i) Me-SO<sub>2</sub>-(R)Cha-Pic-Pab(Z)

10      A solution of 48 mg (0.42 mmol) methanesulfonyl chloride in 0.5 ml methylene chloride was added at 0°C to a stirred solution of 209 mg (0.382 mmol) H-(R)Cha-Pic-Pab(Z) (See Example 25) and 0.11 ml (0.763 mmol) triethyl amine in 5 ml of methylene chloride. The reaction mixture was allowed to reach room temperature over night. Washing with water followed by drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation of the solvent in vacuo gave a residue which was subjected to flash chromatography using ethyl acetate/methanol (95/5) as eluent to give 159 mg (67%) of the product.

(ii) Me-SO<sub>2</sub>-(R)Cha-Pic-Pab x HCl

20      150 mg (0.24 mmol) Me-SO<sub>2</sub>-(R)Cha-Pic-Pab(z) dissolved in 5 ml 95% ethanol and 1 ml water was hydrogenated in the presence of 5% Pd/C for 4 h. Removal of the catalyst by filtration, addition of 0.2 ml 1M hydrochloric acid and evaporation of the solvent in vacuo gave a residue which was dissolved in 2 ml water and freeze dried to give 116 mg (86%) of the product.

30      <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD): δ 0.90-1.10 (m, 2H), 1.15-1.85 (m, 15H), 1.90 (bd, 1H), 2.30 (bd, 1H), 2.85 (s, 3H), 3.35 (dt, 1H), 3.90 (bd, 1H), 4.45 (AB-system, 2H) 4.50-4.55 (m, 1H), 5.13 (dd, 1H), 7.50 (d, 2H), 7.75 (d, 2H).

<sup>13</sup>C-NMR (125 MHz D<sub>2</sub>O): amidine and carbonyl carbons: δ 166.8, 173.0 and 174.6.



Example 37

H-(R)Cha-(R,S)betaPic-Pab x 2 HCl

5 (i) Boc-(R)Cha-(R,S)betaPic-Pab(Z)

EDC was added at -18°C to a stirred solution of 1.0 g (2.6 mmol) Boc-(R)Cha-(R,S)betaPic-OH (See preparation of starting materials), 1.28 g (10.5 mmol) DMAP, 0.74 g (2.6 mmol) H-Pab-(Z) (See preparation of starting materials) in 35 ml DMF. The reaction mixture was allowed to reach room temperature over night and the solvent was subsequently removed in vacuo. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and the organic layer was washed succesively with 0.3M KHSO<sub>4</sub>, 15 KHCO<sub>3</sub>-solution and brine. Drying (Na<sub>2</sub>SO<sub>4</sub>) and removal of the solvent gave a residue which was subjected to flash chromatography using heptane:ethyl acetate with 4% methanol as eluent to yield 0.74 g (44%) of the desired product.

20

(ii) H-(R)Cha-(R,S)betaPic-Pab(Z)

0.68 g (1.05 mmol) Boc-(R)Cha-(R,S)betaPic-Pab(Z) was dissolved in ethyl acetate saturated with HCl(g). The solution was stirred for 1 h at room temperature. Water was added and the mixture was made alkaline with K<sub>2</sub>CO<sub>3</sub>. The water phase was extracted with ethyl acetate. The organic phase was then washed with water and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation gave 0.5 g (87%) of the desired product.

30

(iii) H-(R)Cha-(R,S)betaPic-Pab x 2 HCl

65 mg (0.19 mmol) H-(R)Cha-betaPic(R,S)-Pab(Z) was dissolved in 7 ml ethanol and hydr genat d in pr senc of 5% Pd/C f r 4 hours. Rem val f the catalyst by filtrati n, vaporati n of the solv nt and freeze drying

35

from 1M HCl and water gave 41 mg (71%) of the product.

<sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O, 2 diastereomers 4/5, and rotamers);  
δ 0.8-2.16 (m, ), 2.5-2.77 (m, 3H), 3.13-3.43 (m, 3H),  
5 3.68-3.94 (m, 1H), 4.18-4.41 (m, 1H), 4.41-4.52 (m, 3H),  
7.46-7.57 (m, 2H), 7.72-7.83 (m, 2H).

Example 38

10 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)betaPic-Pab x 2 HCl

(i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)betaPic-Pab(Z)

0.21 g (0.38 mmol) H-(R)Cha-(R,S)betaPic-Pab(Z) (See  
15 Example 37) was dissolved in 2 ml ethanol. 0.68 g (0.42  
mmol) benzyl acrylate was added and the solution was  
stirred for 5 days. Evaporation and flash chromatography  
with CH<sub>2</sub>Cl<sub>2</sub> /MeOH (95/5) as eluent gave 0.19 g (70%) of  
the desired product.

20

(ii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)betaPic-Pab x 2 HCl

170 mg (0.24 mmol) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)betaPic-  
Pab(Z) was dissolved in 10 ml ethanol and hydrogenated in  
25 presence of 5% Pd/C for 4 hours. Removal of the catalyst  
by filtration, evaporation of the solvent and freeze  
drying from 1M HCl and water gave 103 mg (77%) of the  
product.

30 <sup>1</sup>H NMR (300 MHz, D<sub>2</sub>O, mixture of 2 diastereomers 4/5 and  
rotamers); δ 0.92-2.03 (m, H), 2.51-2.78 (m, 1H), 3.21-  
3.52 (m, 1H), 3.88-4.01 (m, 1H), 4.07-4.3 (m, 2H), 4.4-  
4.71 (m, 2H), 7.59 (d, 2H), 7.86 (d, 2H)

35 <sup>13</sup>C NMR (300.13 MHz, D<sub>2</sub>O, mixture of 2 diaster omers 4/5  
and rotamers): amidine and carbonyl carbons: δ 167.0,  
168.0, 168.1, 175.9, 176.0, 176.3, 176.4 and 178.2.

Example 39

**HOOC-CH<sub>2</sub>-(R)Cha-Val-Pab x 2 HCl**

5      (i) Boc-(R)Cha-Val-Pab(Z)

10      1.77 g (9.2 mmol) EDC was added at -12°C to a mixture of  
3.41 g (9.2 mmol) Boc-(R)Cha-Val-OH (See preparation of  
starting materials), 2.61 g (9.2 mmol) H-Pab(Z) (See  
preparation of starting materials), and 4.5 g (36.8 mmol)  
DMAP in 50 ml DMF. The reaction mixture was allowed to  
reach room temperature over night and workup by dilution  
with water was followed by extraction with toluene, ether  
and ethyl acetate. Subsequent drying (MgSO<sub>4</sub>) of the  
15      combined organic extracts, removal of the solvent in  
vacuo and flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH as eluent  
gave 2.77 g (47%) of the desired product.

20      (ii) H-(R)Cha-Val-Pab(Z)

Hydrogen chloride was bubbled through a solution of 2.77  
g (4.4 mmol) Boc-(R)Cha-Val-Pab(Z) in 75 ml ethyl  
acetate. After 15 minutes sodium carbonate solution was  
added to pH 10 and the aqueous phase was extracted with  
25      ethyl acetate. Drying (potassium carbonate) and removal  
of the solvent in vacuo gave 1.8 g (77%) of H-(R)Cha-Val-  
Pab(Z).

30      (iii) BnOOC-CH<sub>2</sub>-(R)Cha-Val-Pab(Z)

A mixture of 326 mg (0.61 mmol) H-(R)Cha-Val-Pab(Z), 105  
ml (0.67 mmol) benzyl bromoacetate, and 252 mg (1.83  
mmol) potassium carbonate in 2 ml acetonitrile was  
sonicated for 2.5 h at 40°C. More acetonitrile was added,  
35      in order to dissolve the product, and the mixture was  
filtered and the solvent was removed in vacuo. The  
residue was purified by flash chromatography using

methanol/methylene chloride as eluent. The product was finally crystallised from ethyl acetate to give 124 mg (30%) of colourless crystals.

5 (iv)  $\text{HOOC-CH}_2\text{-(R)Cha-Val-Pab} \times 2 \text{ HCl}$

124 mg (0.18 mmol)  $\text{BnOOC-CH}_2\text{-(R)Cha-Val-Pab(Z)}$  in 20 ml ethanol was hydrogenated for 2 hours in the presence of 25 mg 10% Pd/C. 10 ml of THF was added and the  
10 hydrogenation was continued for another 2 hours at 50°C. The mixture was filtered through hyflo and the filtercake was washed with dilute hydrochloric acid. The organic solvents were removed from the combined filtrates in vacuo. Freeze drying of the remaining solution yielded 55  
15 mg (50%) of the desired compound.

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ );  $\delta$  0.75-1.4 (m, 12H), 1.5-1.9 (m, 7H), 2.0-2.15 (bs, 1H), 3.45 (AB-system, 2H), 4.1 (m, 2H), 4.5 (m, 2H), 7.5 (s, 2H), 7.7 (s, 2H), 8.9 (s, 1H).

20

Example 40

$\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Val-Pab} \times 2 \text{ HCl}$

25 (i)  $\text{H-(R)Cha-(R,S)Val-Pab(Z)}$

The title compound was prepared by coupling Boc-(R)Cha-Val-OH with H-Pab(Z), using the pivaloyl coupling as described for Boc-(R)Cha-Pic-OMe (See preparation of  
30 starting materials). A total epimerization of the valine occurred to give Boc-(R)Cha-(R,S)Val-Pab(Z). The Boc protecting group was removed in the same way as described for Boc-(R)Cha-Val-Pab(Z) (See Example 39) to give the title compound.

35

(ii)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-(R,S)Val-Pab(Z)}$

A solution of 1.007 g (1.9 mmol) H-(R)Cha-(R,S)Val-Pab(Z) and 308 mg (1.9 mmol) benzyl acrylate in 3 ml of ethanol was kept at 40°C over night. The solvent was removed in vacuo and the residue was purified by flash chromatography using methanol/methylene chloride (10/90) as eluent to give 1.086 g (82%) of the title compound.

(iii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Val-Pab x 2 HCl

1.086 g (1.6 mmol) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-(R,S)Val-Pab(Z) was hydrogenated in 25 ml THF and 14 ml 0.5 M hydrochloric acid in the presence of 223 mg 10% Pd/C for 2 hours. Removal of the catalyst by filtration through celite and removal of the THF in vacuo followed by freeze drying of the remaining aqueous solution gave a residue of which approximately 300 mg was subjected to HPLC using 25% acetonitrile in 0.1 M Ammonium acetate buffer as eluent. Two main fractions were isolated, of which the second fraction contained the title compound. 67 mg of title compound, as the dihydrochloride, was isolated.

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O); δ 1.0-1.15 (m, 12H), 1.2-1.4 (m, 7H), 1.65-1.9 (m, 7H), 2.15-2.25 (m, 1H), 2.85 (t, 2H), 3.15-3.2 (m, 1H), 3.3-3.35 (m, 1H), 4.15-4.2 (m, 1H), 4.25 (d, 1H), 4.55-4.65 (AB-system, 2H), 7.65 (d, 2H), 7.85 (d, 2H).

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 167.0, 169.8, 173.96 and 174.04.

#### Example 41

H-(R)Hoc-Aze-Pab x 2 HCl

(i) Boc-(R)Hoc-Aze-Pab(Z)

Prepared in the same way as described for Boc-(R)Cha-Pic-

Pab(Z) (See Example 25) by replacing Boc-(R)Cha-Pic-OH with Boc-(R)Hoc-Aze-OH (See preparation of starting materials). The crude product was subjected to flash chromatography (Toluene/EtOAc 1/6) to give 0.32 g (37%) of the desired product.

(ii) H-(R)Hoc-Aze-Pab(Z)

Boc-(R)Hoc-Aze-Pab(Z) was treated in the same way as described for Boc-(R)Cha-Pic-Pab(Z) in Example 25 to give 0.23 g (88%) of the title compound.

(iii) H-(R)Hoc-Aze-Pab x 2 HCl

20 mg (0.037 mmol) of H-(R)Hoc-Aze-Pab(Z) was dissolved in 3 ml ethanol and hydrogenated in presence of 5% Pd/C for 4 hours at atmospheric pressure. Removal of the catalyst by filtration, evaporation of the solvent and freeze drying from 1M HCl gave 11 mg (63%) of the product.

<sup>1</sup>H NMR (300.13 MHz, D<sub>2</sub>O, mixture of two rotamers 3:1): major rotamer: δ 0.9-2.1 (m, 15H), 2.4-2.6 (m, 1H), 2.7-3.0 (m, 1H), 4.1-4.3 (m, 1H), 4.35-4.56 (m, 1H), 4.65 (s, 2H), 5.0-5.11 (m, 1H), 7.62 (d, 2H), 7.9 (d, 2H). The signal of one of the protons is totally obscured by the H-O-D-signal.

#### Example 42

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Hoc-Aze-Pab x 2 TFA

(i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Hoc-Aze-Pab(Z)

0.067 g (0.41 mmol) benzylacrylate was added to a solution of 0.2 g (0.37 mmol) H-(R)Hoc-Aze-Pab(Z) (See Example 41) in 2 ml ethanol (95%) at room temperature.

The reaction was left at room temperature for 5 days. The solvent was removed in vacuo and the residue was purified with flash chromatography ( $\text{CH}_2\text{Cl}_2$ : MeOH, 96/4) to give 0.16 g (62%) of the desired product.

5

(ii)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Hoc-Aze-Pab} \times 2 \text{ TFA}$

160 mg (0.23 mmol)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Hoc-Aze-Pab(Z)}$  was dissolved in 10 ml ethanol and subjected to hydrogenation at atmospheric pressure in presence of 5% Pd on charcoal for 3 hours. Removal of the catalyst by filtration, evaporation of the solvent and freeze drying from water and TFA gave 120 mg (87%) of the product.

$^1\text{H}$  NMR (300.13 MHz,  $\text{D}_2\text{O}$  2 rotamers 3:1); major rotamer:  $\delta$  0.9-1.9 (m, 13H), 1.94-2.16 (m, 2H), 2.38-2.55 (m, 1H), 2.7-2.97 (m, 3H), 3.2-3.44 (m, 2H), 4.16 (m, 1H), 4.35-4.58 (m, 2H), 4.65 (s, 2H), 5.0-5.12 (m, 1H), 7.63 (d, 2H), 7.87 (d, 2H)

20

$^{13}\text{C}$  NMR (300.13 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  167.3, 168.7, 172.5 and 176.6.

#### Example 43

25

$\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Hoc-Pro-Pab} \times 2 \text{ HCl}$

(i)  $\text{Boc-(R)Hoc-Pro-Pab(Z)}$

Prepared from  $\text{Boc-(R)Hoc-Pro-OH}$  (See preparation of starting materials) in the same way as described for  $\text{Boc-(R)Cha-Pic-Pab(Z)}$  in Example 25. Flash chromatography using ethyl acetate as eluent gave 0.886 g (58 %) of the title compound.

35

$^1\text{H-NMR}$ . (300 MHz,  $\text{CDCl}_3$ );  $\delta$  0.7-0.95 (m, 2H), 0.95-2.1 (m, 27H (there f 1.2 (s, 9H)), 2.1-2.4 (m, 1H), 3.3-3.5 (m,

1H), 3.65-3.95 (m, 1H), 4.0-4.2 (m, 1H), 4.2-4.45 (m, 2H), 4.45-4.6 (d, 1H), 5.15 (apparent bs, 2H), 5.2-5.3 (d, 1H), 7.1-7.4 (m, 7H), 7.65 (m, 1H), 7.7-7.8 (d, 2H), 9.4 (bs, 1H).

5

<sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 156.3, 164.6, 168.1, 171.4 and 172.4.

10 (ii) H-(R)Hoc-Pro-Pab(Z)

40 ml ethyl acetate saturated with hydrogen chloride was added to 0.82 g (1.266 mmol) Boc-(R)Hoc-Pro-Pab(Z) at 0°C. The temperature was allowed to rise to room-temperature. The reaction was not completed after 1.5 h and therefore hydrogen chloride was bubbled through the reaction mixture during 5 minutes. The solvent was evaporated and ethyl acetate and saturated sodium carbonate was added and the phases were separated. The organic phase was washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent evaporated in vacuo to give the title compound in almost quantitative yield.

25 <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>); δ 0.75-0.95 (m, 2H), 0.95-2.4 (m, 17H), 3.3-3.55 (m, 2H), 3.55-3.7 (m, 1H), 4.25-4.45 (m, 2H), 4.5-4.6 (m, 1H), 5.15 (s, 2H), 7.15-7.35 (m, 5H), 7.35-7.45 (m, 2H), 7.6-7.7 (m, 1H), 7.7-7.85 (d, 2H).

30 <sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 164.5, 167.8, 171.4 and 175.3.

(iii) BnOOC-CH<sub>2</sub>-(R,S)CH(COObn)-(R)Hoc-Pro-Pab(Z)

35 To 0.15 g (0.5 mmol) benzyl acrylate in 1.5 ml EtOH (99%) was added 0.273 g (0.498 mmol) H-(R)Hoc-Pro-Pab(Z) and the mixture was stirred at room temperature for 10 days.



The solvent was removed in vacu and the residue was subject d to flash chromatography, using ethyl ac tate as eluent to give 0.103 g (25 %) of  $\text{BnOOC-CH}_2\text{-(R,S)CH(COOBn)-(R)Hoc-Pro-Pab(Z)}$ .

5  
 $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ );  $\delta$  0.75-2.05 (m, 18H), 2.3-2.45 (m, 1H), 2.45-2.8 (m, 3H), 3.15-3.45 (m, 3H), 3.5-3.65 (m, 1H), 4.3-4.5 (m, 2H), 4.55-4.7 (m, 1H), 4.8 (s, 1H), 4.9-5.1 (m, 3H), 5.2 (s, 2H), 7.1-7.2 (m, 1H), 7.2-7.4  
10 (m, 13H), 7.4-7.45 (d, 2H), 7.6-7.8 (m, 3H).

(iv)  $\text{HOOC-CH}_2\text{-(R,S)CH(COOH)-(R)Hoc-Pro-Pab} \times 2 \text{ HCl}$

15 103 mg (0.122 mmol)  $\text{BnOOC-CH}_2\text{-(R,S)CH(COOBn)-(R)Hoc-Pro-Pab(Z)}$  dissolved in 4 ml ethanol (99.5 %) and 0.3 ml chloroform was hydrogenated in the presence of 111 mg 5 % Pd/C for 2 h. Removal of the catalyst by filtration and evaporation of the solvent followed by dissolving in  
20 water and freeze drying showed incomplete hydrogenation. The hydrogenation was continued in the presence of ethanol, 1 N HCl and 5 % Pd/C for 5 hours.

25 Removal of the catalyst by filtration and evaporation of the solvent followed by dissolving in water and freeze drying gave the title compound.

$^1\text{H-NMR}$  (500 MHz,  $\text{CD}_3\text{OD}$ , mixture of two diastereomers);  $\delta$  0.8-1.0 (m, 2H), 1.1-1.4 (m, 6H), 1.6-1.8 (m, 5H), 1.9-  
30 2.15 (m, 5H) 2.25-2.35 (m, 1H), 2.9-3.2 (m, 2H), 3.5-3.65 (m, 1H), 3.7-3.9 (2m, total 1H), 4.15-4.4 (2m, total 1H), 4.4-4.6 (m, 4H), 7.5-7.6 (m, 2H), 7.7-7.85 (m, 2H).

$^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$   
35 167.9, 168.2, 168.3, 172.8, 173.6, 174.3 and 174.4. The signals fr m the tw diastereomers are partly overlapping.

Example 44**HOOC-CH<sub>2</sub>-(R)Hoc-Pic-Pab x 2 HCl****5 (i) Boc-(R)Hoc-Pic-Pab(Z)**

Prepared from Boc-(R)Hoc-Pic-OH (See preparation of starting materials) and H-Pab(Z) (See preparation of starting materials) in the same way as described for Boc-  
10 (R)Cha-Pic-Pab(Z) (See Example 25). Flash chromatography using ethyl acetate as eluent gave 1.3 g (78 %) of the title compound.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 0.75-0.95 (m, 2H), 0.95-2.0 (m,  
15 31H (thereof 1.3 (s, 9H)), 2.4-2.5 (m, 1H), 3.0-3.1 (m, 1H), 3.8 (m, 1H), 4.2-4.45 (m, 2H), 4.45-4.55 (m, 2H), 5.15 (apparent bs, 3H), 5.25-5.3 (m, 1H), 7.0 (bs, 1H), 7.15-7.5 (m, 7H), 7.7-7.85 (d, 2H), 9.45 (bs, 1H).

20 <sup>13</sup>C-NMR (75 MHz, CDCl<sub>3</sub>): amidine and carbonyl carbons: δ 156.6, 164.7, 168.1, 170.0 and 173.0.

**(ii) H-(R)Hoc-Pic-Pab(Z)**

25 100 ml ethyl acetate saturated with hydrogen chloride was added to 1.3 g (1.96 mmol) Boc-(R)Hoc-Pic-Pab(Z) at 0°C. The temperature was allowed to rise to room-temperature. The solvent was evaporated after 40 minutes and ethyl acetate and saturated sodium carbonate was added and the  
30 phases were separated. The organic phase was washed with brine and dried (Na<sub>2</sub>SO<sub>4</sub>) and the solvent evaporated in vacuo to give 0.85 g (77.5 %) of the product.

35 <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>): δ 0.75-0.95 (m, 2H), 1.05-2.3 (m, 25H), 3.0-3.15 (m, 1H), 3.6-3.75 (m, 2H), 4.25-4.4 (m, 2H), 5.15 (apparent bs, 3H), 7.05-7.2 (d, 2H), 7.2-7.35 (m, 4H), 7.35-7.4 (d, 1H), 7.6-7.8 (d, 2H).

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.5, 167.9, 170.8 and 175.7.

5 (iii)  $\text{BnOOC-CH}_2\text{-(R)Hoc-Pic-Pab(Z)}$

0.171 g (0.748 mmol) benzyl bromoacetate was added to a mixture of 0.4 g (0.712 mmol)  $\text{H-(R)Hoc-Pic-Pab(Z)}$  and 0.217 g (1.57 mmol)  $\text{K}_2\text{CO}_3$  in 7 ml acetonitrile. The mixture was heated to  $60^\circ\text{C}$  in oilbath for 1 h. The solvent was removed and ethyl acetate and water was added. The phases were separated and the organic phase was washed with brine and dried ( $\text{Na}_2\text{SO}_4$ ). Evaporation in vacuo gave 0.626 g of a residue which was subjected to flash chromatography using ethyl acetate as eluent, to give 2 products. The first compound eluted from the column was  $(\text{BnOOC-CH}_2)_2(\text{R})\text{Hoc-Pic-Pab(Z)}$  (0.28 g) and the second compound eluted was the title compound (0.27 g).

20  $\text{BnOOC-CH}_2\text{-(R)Hoc-Pic-Pab(Z)}$ :

$^1\text{H}$ -NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.7-0.95 (m, 2H), 1.0-1.75 (m, 18H), 2.3-2.5 (m, 1 or 2H), 2.9-3.05 (m, 1H), 3.2-3.3 (m, 1H), 3.35-3.5 (m, 2H), 3.6-3.7 (m, 1H), 4.35, 4.55 (ABX-system, 2H), 4.75 (s, 2H), 5.15 (apparent s, 3H), 5.25-5.3 (m, 1H), 7.1-7.45 (m, 12H), 7.7-7.8 (d, 2H).

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.6, 167.9, 170.5, 173.4 and 175.0.

30

(iv)  $\text{HOOC-CH}_2\text{-(R)Hoc-Pic-Pab} \times 2 \text{ HCl}$

259 mg (0.365 mmol)  $\text{BnOOC-CH}_2\text{-(R)Hoc-Pic-Pab(Z)}$  dissolved in 7.8 ml ethanol (99.5 %) and 1.2 ml hydrogen chloride (1 N) was hydrogenated in the presence of 280 mg 5 % Pd/C for 4 h. Removal of the catalyst by filtration and evaporation of the solvent followed by dissolving in

water and freeze drying gave 170 mg (83 %) of the title compound.

5  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.4-1.85 (m, 20H), 1.85-2.2 (m, 1H), 2.9-3.2 (m, 1H), 3.4-3.9 (m, 3H), 4.05-4.3 (m, 2H), 4.3-5.05 (m, 2H), 7.1-7.4 (m, 2H), 7.4-7.7 (m, 2H).

10  $^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  167.8, 168.6, 169.6 and 172.3.

Example 45

$(\text{HOOC-CH}_2)_2\text{-(R)Hoc-Pic-Pab} \times 2 \text{ HCl}$

15 (i)  $(\text{BnOOC-CH}_2)_2(\text{R})\text{Hoc-Pic-Pab(Z)}$

The title compound was obtained in the alkylation of H-(R)Hoc-Pic-Pab(Z) as described in Example 44 above.

20  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.7-0.95 (m, 2H), 0.95-1.95 (m, 18H), 2.35-2.5 (m, 1H), 2.9-3.05 (m, 1H), 3.5-3.85 (m, 6H), 4.35-4.55 (m, 2H), 4.9 (2s, 4H), 5.2 (s, 2H), 5.25-5.35 (m, 1H), 7.1-7.45 (m, 16H), 7.5-7.65 (m, 1H), 7.7-7.85 (d, 2H).

25  $^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.7, 167.9, 170.5, 172.0 and 172.4.

(ii)  $(\text{HOOC-CH}_2)_2\text{-(R)Hoc-Pic-Pab} \times 2 \text{ HCl}$

30 153 mg (0.178 mmol)  $(\text{BnOOC-CH}_2)_2\text{-(R)Hoc-Pic-Pab(Z)}$  dissolved in 4.5 ml ethanol (99.5 %) and 0.5 ml hydrogen chloride (1 N) was hydrogenated in the presence of 150 mg 5 % Pd/C for 3.5 h. Removal of the catalyst by filtration and evaporation of the solvent followed by dissolving in water and freeze drying gave 109 mg (99 %) of  $(\text{HOOC-CH}_2)_2\text{-(R)Hoc-Pic-Pab}$  dihydrochloride. This crude material

(80 % purity) was subjected to purification by RPLC using CH<sub>3</sub>CN/0.1 M NH<sub>4</sub>OAc, 1:4 as eluent. Removal of the solvent and excess NH<sub>4</sub>OAc followed by freeze drying from 1 M HCl gave the title compound.

5

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O, mixture of two rotamers): major rotamer: δ 0.95-2.15 (m, 20H), 2.25-2.35 (m, 1H), 3.45-3.55 (m, 1H), 3.95-4.25 (m, 5H), 4.6-4.65 (m, 2H), 4.92-5.01 (m, 1H) 5.15-5.20 (m, 1H), 7.58-7.63 (d, 2H), 7.84-7.89 (d, 2H).

10

Resolved signals arising from the minor rotamer appears at: δ 0.7-0.85 (m), 2.35-3.45 (m), 3.05-3.15 (m), 4.47-4.55 (m), 4.55-4.6 (m), 4.65-4.7 (m), 7.63-7.67 (d), 7.89-7.95 (d).

15

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 168.20, 169.70, 170.20 and 172.71.

#### 20 Example 46

HOOC-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab x 2 HCl

(i) Boc-(R)Pro(3-(S)Ph)-Pro-Pab(Z)

25

To a solution of 570 mg (1.5 mmol) Boc-(R)Pro(3-(S)Ph)-Pro-OH (See preparation of starting materials), 425 mg (1.5 mmol) H-Pab(Z) (See preparation of starting materials) and 733 mg (6 mmol) DMAP in 25 ml CH<sub>3</sub>CN/DMF (1.5/1) was added 310 mg (1.62 mmol) EDC and the mixture was stirred for 23 h at room temperature. Most of the solvent was evaporated and 50 ml water was added to the residue. The water phase was extracted with 1 x 75 and 2 x 50 ml EtOAc. The combined organic phase was washed with 1 x 20 + 1 x 10 ml 1M KHSO<sub>4</sub>, 1 x 15 ml NaHCO<sub>3</sub>(aq), 3 x 15 ml water, 1 x 15 ml brine and dried (MgSO<sub>4</sub>). Filtration and evaporation of the solvent gave 670 mg of an oil

30

35

which was purified by flash chromatography using EtOAc as eluent which gave 529 mg (55 %) of the title compound.

5  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.26 (s, 9H), 1.53-1.88 (m, 3H), 2.1-2.31 (m, 3H), 2.52 (q, 1H), 3.58-3.77 (m, 4H), 4.31 (d, 1H), 4.35 and 4.47 (ABX-system, 2H), 4.65 (dd, 1H), 5.19 (s, 2H), 7.1-7.37 (m, 10H), 7.42 (d, 2H), 7.81 (d, 2H), 8.0 (t, 1H (NH)).

10  $^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  154.6, 164.6, 168.1, 171.1 and 171.3.

(ii) H-(R)Pro(3-(S)Ph)-Pro-Pab(Z)

15 529 mg (0.81 mmol) of Boc-(R)Pro(3-(S)Ph)-Pro-Pab(Z) was dissolved in 15 ml EtOAc/HCl(g,saturated) at room temperature and stirred for 3 h. The solvent was evaporated and the residue was dissolved in 70 ml  $\text{CH}_2\text{Cl}_2$ . the organic phase was washed with 1 x 10 ml 2 M NaOH, 1  
20 x 10 ml water, 1 x 10 ml brine and dried ( $\text{MgSO}_4$ ). Filtration and evaporation of the solvent gave 403 mg (90 %) of the title compound as a white powder.

25  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.44-1.57 (m, 1H), 1.62-1.86 (m, 2H), 1.96-2.35 (m, 3H), 2.45 (q, 1H), 3.05-3.35 (m, 4H), 3.83 (bd, 1H), 4.25-4.45 (m, 2H), 4.53 (m, 1H), 5.19 (s, 2H), 7.16-7.37 (m, 10H), 7.42 (d, 2H), 7.66 (t, 1H, (NH)), 7.77 (d, 2H).

30  $^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.4, 167.9, 171.1 and 173.0.

(iii)  $\text{BnOOC-CH}_2\text{-(R)Pro(3-(S)Ph)-Pro-Pab(Z)}$

35 A mixture of 200 mg (0.36 mmol) H-(R)Pro(3-(S)Ph)-Pro-Pab(Z), 105 mg (0.46 mmol)  $\text{Br-CH}_2\text{-COOBn}$  and 125 mg (0.90 mmol)  $\text{K}_2\text{CO}_3$  in 10 ml  $\text{CH}_3\text{CN}$  was heated to 50°C for 1 h and

30 minutes. The solvent was evaporated and the residu  
was dissolved in 70 ml EtOAc. The organic phase was  
washed with 10 ml water and dried ( $\text{MgSO}_4$ ). Filtration and  
evaporation of the solvent gave 260 mg of an oil. The  
5 crude material was purified by flash chromatography using  
a stepwise gradient of  $\text{CH}_2\text{Cl}_2/\text{MeOH}(\text{NH}_3\text{-saturated})$  (95/5  
followed by 9/1) to give 182 mg (72 %) of the title  
compound as a white solid.

10  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.43-1.82 (m, 3H), 1.96-2.13  
(m, 1H), 2.14-2.22 (m, 1H), 2.26-2.43 (m, 2H), 3.02-3.14  
(m, 2H), 3.24-3.51 (m, 4H), 3.83 (d, 1H), 4.29-4.46 (ABX-  
system centered at 4.37, 2H), 4.58 (dd, 1H), 4.97-5.1  
15 (AB-system centered at 5.03, 2H), 5.19 (s, 2H), 7.16-7.38  
(m, 15H), 7.43 (d, 2H), 7.5-7.8 (m, 3H, one NH).

$^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$   
164.5, 167.9, 171.15, 171.2 and 172.7.

20 (iv)  $\text{HOOC-CH}_2\text{-(R)Pro(3-(S)Ph)-Pro-Pab} \times 2 \text{ HCl}$

0.18 g ( 0.26 mmole ) of  $\text{BnOOC-CH}_2\text{-(R)Pro(3-(S)Ph)-Pro-}$   
 $\text{Pab(Z)}$  was mixed with 0.075 g 5 % Pd/C, 1.0 ml 1N HCl-  
solution, 1 ml water and 10 ml ethanol and the mixture  
25 was hydrogenated at atmospheric pressure for one hour.  
Filtration of the catalyst through hyflo, evaporation of  
the solvent followed by freeze drying twice from water  
gave 129 mg of a crude product . The crude product was  
purified by RPLC using a stepwise gradient of 0.1 M  
30  $\text{NH}_4\text{OAc/CH}_3\text{CN}$  4/1 followed by 3/1. Evaporation followed by  
freeze drying from water and 1N HCl-solution gave 70 mg  
(50%) of the pure product.

35  $^1\text{H-NMR}$  (300 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  1.42-1.60 (m, 1H), 1.65-1.83 (m,  
1H), , 1.83-1.98 (m, 1H), 2.03-2.20 (m, 2H), 2.63 (t, 2H),  
3.28-3.40 (m, 1H), 3.55-3.78 (m, 2H), 3.81-3.96 (AB-  
system central at  $\delta$  3.88, 2 H), 4.06-4.19 (m, 1H), 4.37-

4.61 (AB-system central at  $\delta$  4.49, 2 H), 4.48 (dd, 1H),  
4.70 (d, 1H), 7.35-7.58 (m, 7H), 7.74 (d, 2H)

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$   
5 167.02, 167.2, 169.3 and 174.4.

#### Example 47

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab x 2 HCl

10

(i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab(Z)

To a solution of 190 mg (0.34 mmol) H-(R)Pro(3-(S)Ph)-  
Pro-Pab(Z) (See Example 46) in 7 ml EtOH (99%) was added  
15 114 mg (0.70 mmol) of benzyl acrylate and the reaction  
mixture was stirred at room temperature for 24 h.  
Evaporation of the solvent followed by flash  
chromatography using a stepwise gradient of  
CH<sub>2</sub>Cl<sub>2</sub>/MeOH(NH<sub>3</sub>-saturated) (95/5 followed by 9/1) gave 202  
20 mg (83 %) of the title compound.

$^1\text{H}$ -NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.5-1.71 (m, 2H), 1.74-1.9 (m,  
1H), 1.9-2.05 (m, 1H), 2.2-2.64 (m, 5H), 2.69-2.82 (m,  
2H), 2.84-2.96 (m, 1H), 3.18-3.48 (m, 4H), 4.28-4.44 (m,  
25 2H), 4.61 (m, 1H), 4.48-5.08 (AB-system centered at 5.03,  
2H), 5.19 (s, 2H), 7.15-7.37 (m, 15H), 7.44 (d, 2H),  
7.75-7.85 (m, 3H, one NH).

$^{13}\text{C}$ -NMR (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$   
30 164.6, 168.0, 171.2, 172.5 and 172.9.

(ii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-Pro-Pab x 2 HCl

35 0.20 g ( 0.28 mmole ) of BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Pro(3-(S)Ph)-  
Pr -Pab(Z) was mixed with 0.075 g 5 % Pd/C, 1.0 ml 1N  
HCl-soluti n, 1 ml water and 10 ml ethanol and the



mixture was hydr genat d at atmospheric pr ssure for one hour. Filtration of th catalyst through hyfl , evaporation of the solvent followed by freeze drying twice from water gave 125 mg 79 % of the title compound.

5

$^1\text{H-NMR}$  (300 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  1.44 (m, 1H), 1.65-1.9 (m, 2H), 2.0-2.2 (m, 2H), 2.62 (q, 2H), 2.83 (t, 2H), 3.27-3.4 (m, 1H), 3.4-3.8 (m, 4H), 4.0-4.15 (m, 1H), 4.35-4.6 (m, 3H), 4.68 (d, 1H), 7.35-7.6 (m, 7H), 7.77 (d, 2H)

10

$^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  166.2, 167.1, 174.1 and 174.2.

#### Example 48

15

$\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Tic-Pro-Pab x 2 HCl}$

(i) Boc-(R)Tic-Pro-Pab(Z)

20

Prepared in the same way as described for Boc-(R)Cha-Pic-Pab(Z) (See Example 25) using Boc-(R)Tic-Pro-OH(See preparation of starting materials) instead of Boc-(R)Cha-Pic-OH. Flash chromatography using heptane/EtOAc (4/1) followed by EtOAc as eluents gave 425 mg (37%) of the

25

$^1\text{H NMR}$  (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.35 (s, 9H), 1.95-2.15 (m, 3H), 2.4 (m, 1H), 2.8 (m, 1H), 3.3 (m, 1H), 3.55 (m, 2H), 4.25-4.4 (two m, 2H), 4.55-4.7 (two m, 2H), 7.15-7.5 (m, 10H), 7.85 (d, 2H).

30

$^{13}\text{C-NMR}$  (75.0 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.6, 171.5 and 171.6. (two peaks are probably overlapping)

35

(ii) H-(R)Tic-Pro-Pab(Z)

Boc-(R)Tic-Pro-Pab(Z) (379 mg, 0.59 mmol) was dissolved in EtOAc saturated with HCl(g) and stirred at room temperature. Evaporation of the solvent gave 251 mg (79%) of the title compound as a white powder.

5

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.65-2.15 (two m, 7H), 2.45 (m, 1H), 2.75 (m, 1H), 2.9 (m, 1H), 3.0 (m, 1H), 3.25 (m, 1H), 3.55 (m, 1H), 3.85 (m, 1H), 4.35-4.55 (m, 2H), 4.75 (d, 1H), 4.9 (s, 1H), 5.25 (s, 2H), 6.8-7.45 (several m, 8H), 7.5 and 7.85 (two d, 4H).

10

$^{13}\text{C}$ -NMR (75.0 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  164.5, 171.3 and 172.7 (two peaks are probably overlapping).

15

(iii)  $\text{BnO}_2\text{C}-\text{CH}_2-\text{CH}_2-(\text{R})\text{Tic-Pro-Pab}(\text{Z})$

H-(R)Tic-Pro-Pab(Z) (140 mg, 0.26 mmol) was treated with benzyl acrylat (63 mg, 0.39 mmol) in EtOH (1.3 ml) at 20°C during 48 h. Evaporation of the solvent and flash chromatography using (50% EtOAc/Heptan then 10% MeOH/EtOAc) as eluent afforded 133 mg (73%) of the desired product as a white solid material.

25

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.75-2.0 (two m, 4H), 2.25 (m, 1H), 1.4-1.65 (m, 3H), 2.7-2.95 (two m, 4H), 3.05-3.2 (m, 2H), 3.9 (m, 1H), 4.45 (m, 2H), 4.65 (m, 1H), 5.1 (two d, 2H), 5.25 (s, 2H), 6.85-7.45 (several m, 12H), 7.5 and 7.9 (two d, 4H).

30

$^{13}\text{C}$ -NMR (75.0 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$  171.5, 171.9 and 172.1 (two peaks are probably overlapping).

35

(iv)  $\text{HOOC}-\text{CH}_2-\text{CH}_2-(\text{R})\text{Tic-Pro-Pab} \times 2 \text{ HCl}$

BnO<sub>2</sub>C-CH<sub>2</sub>-CH<sub>2</sub>-(R)Tic-Pr -Pab(Z) (125 mg, 0,17 mmol) was hydrogenated ver 5 % Pd/C using EtOH/HCl as solvent. Filtration of the catlyst and freeze drying gave 73 mg (77%) of the title compound as a white powder.

5

<sup>1</sup>H NMR (500 MHz, D<sub>2</sub>O): δ 2.1-2.35 (two m, 3H), 2.6 (m, 1H), 2.95-3.1 (m, 2H), 3.25-3.5 (two m, 2H), 3.65 (m, 3H), 4.65 (s, 2H), 4.75 (m, 1H), 5.85 (s, 1H), 7.15-7.6 (three m, 4H), 7.6 and 7.85 (two d, 4H).

10

<sup>13</sup>C-NMR (75.0 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 166.9, 167.1 and 174.3 (two peaks are probably overlapping).

15

Example 49

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Aze-Pig x 2 HCl

(i) Boc-(R)Cgl-Aze-Pig(Z)<sub>2</sub>

20

To a mixture of 0.623 g (1.83 mmole) Boc-(R)Cgl-Aze-OH(See preparation of starting materials), 0.816 g (1.92 mmole) H-Pig(Z)<sub>2</sub> (See preparation of starting materials) and 0.89 g (7.3 mmole) DMAP in 10 ml dichloromethane was added 0.368 g (1.92 mmole) of EDC and the mixture was stirred over night. The mixture was diluted and washed with 0.3 M KHSO<sub>4</sub> and once with brine. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to yield 1.4 g of a crude product. Purification by flash chromatography using ethyl acetate as eluent gave 0.3 g (22%) of the pure product.

25

30

(ii) H-(R)Cgl-Aze-Pig(Z)<sub>2</sub>

35

0.3 g (0.4 mmole) Boc-(R)Cgl-Aze-Pig(Z)<sub>2</sub> was mixed with 10 ml dichloromethan and 2.5 ml trifluoroacetic acid. The mixture was stirred f r n and a half h ur. After

evaporation of the solvent the residue was dissolved in dichloromethane and washed twice with 0.2 M NaOH-solution. The combined water layer was extracted one more time with dichloromethane. The combined organic layer was  
5 dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to yield 0.24 g (93%) of the product.

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ , 339K):  $\delta$  0.9-1.9 (m, 15H), 1.94 (bd, 1H), 2.37-2.52 (m, 1H), 2.65-2.8 (m, 1H), 2.9-3.08  
10 (m, 3H), 3.20 (t, 2H), 4.05-4.28 (m, 4H), 4.86 (dd, 1H), 5.16 (s, 4H), 7.2-7.42 (m, 10H), 7.98 (bs, NH).

(iii)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cgl-Aze-Pig(Z)}_2$

0.231 g (0.36 mmole) was dissolved in 2 ml ethanol and 61  
15  $\mu\text{l}$  (0.40 mmole) benzylacrylate was added. The reaction mixture was stirred for five days at room temperature. The mixture was evaporated and the crude product purified by flash chromatography using a stepwise gradient of  
20  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (95/5, 90/10) as eluent to yield 0.218 g (75%) of the pure product.

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ , 335K):  $\delta$  0.93 (bq, 1H), 1.02-1.85 (m, 14H), 1.94 (bd, 1H), 2.33-2.5 (m, 3H), 2.58-2.77 (m,  
25 2H), 2.79-3.02 (m, 4H), 3.17 (t, 2H), 4.0-4.25 (m, 4H), 4.86 (dd, 1H), 5.11 (s, 2H), 5.12 (s, 4H), 7.2-7.4 (m, 15H), 8.03 (bs, NH), 10.35 (bs, NH)

30 (iv)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cgl-Aze-Pig} \times 2 \text{ HCl}$

0.218 g (0.27 mmole) of  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cgl-Aze-Pig(Z)}_2$  was mixed with 0.10 g 5% Pd/C, 1 ml 1M HCl-solution, 1 ml water and 10 ml ethanol and the mixture was  
35 hydrogenated at atmospheric pressure for one hour. Filtration of the catalyst through hyflo, evaporation of the solvent followed by freeze drying twice from water

gav 134 mg (95%) of the title compound.

<sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 1.0-1.4 (m, 7H), 1.55-2.05 (m, 9H), 2.22-2.34 (m, 1H), 2.61-2.76 (m, 1H), 2.88 (t, 2H),  
5 3.08 (bt, 2H), 3.19 (d, 2H), 3.34 (m, 2H), 3.83 (bd, 2H),  
3.95 (d, 1H), 4.29-4.49 (m, 2H), 4.90 (dd, 1H)

<sup>13</sup>C-NMR (75 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ  
10 156.4, 167.6, 172.1 and 174.7

Example 50

HOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pig x 2 HCl

15 (i) Boc-(R)Cgl-Pro-Pig(Z)<sub>2</sub>

0.568 g (2.96 mmol) EDC was added at -15°C to a mixture  
of 1 g (2.82 mmol) Boc-(R)Cgl-Pro-OH (See preparation of  
starting materials), 1.197 g (2.82 mmol) H-Pig(Z)<sub>2</sub> (See  
20 preparation of starting materials) and 1.38 g (11.28  
mmol) DMAP in acetonitrile. The temperature was allowed  
to rise to roomtemperature over night. The solvent was  
evaporated in vacuo and methylenchloride and 1 M KHSO<sub>4</sub>  
was added. The phases were separated and the organic  
25 phase was washed with saturated NaHCO<sub>3</sub>, water and brine,  
drying (Na<sub>2</sub>SO<sub>4</sub>) and evaporation of the solvents gave 2.033  
g of a residue which was subjected to flash chromatography  
using ethylacetate as eluent. This gave two products; 720  
mg (34 %) of the title compound which eluted first from  
30 the column followed by 775 mg (44 %) of Boc-(R)Cgl-Pro-  
Pig(Z) formed by loss of one of the Z-protecting groups.

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>); Some signals, especially in the  
piperidin ring, are selectively broader due to an  
35 intramolecular exchange process. This is especially  
pronounced for the 2- and 6-CH<sub>2</sub> groups of the piperidin  
ring, which exhibit a broad peak ranging from 3.5 to 4.5

ppm.

5  $\delta$  0.85-2.1 (m, 19H), 2.3-2.45 (m, 1H), 2.8-3.2 (m, 4H),  
3.45-3.55 (m, 1H), 3.55-3.65 (m, minor rotamer), 3.8-3.93  
(m, 1H), 3.97-4.1 (m, 1H), 4.52-4.62 (d, 1H), 5.1  
(apparent bs, 5H), 7.12-7.41 (m, 10H).

10  $^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons:  $\delta$   
155.2, 156.3, 171.0 and 172.1.

(ii) H-(R)Cgl-Pro-Pig(Z)<sub>2</sub>

15 720 mg (0.946 mmol) of Boc-(R)Cgl-Pro-Pig(Z)<sub>2</sub> was  
dissolved in 35 ml of TFA/ $\text{CH}_2\text{Cl}_2$ , 1/4 and stirred for 30  
minutes. The solvent was removed in vacuo and  
ethylacetate and 2M NaOH was added. The organic layer was  
washed with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ) and the  
solvent was evaporated in vacuo to give the title  
compound in quantitative yield.

20  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ); Some signals, especially in the  
piperidin ring, are selectively broader due to an  
intramolecular exchange process. This is especially  
pronounced for the 2- and 6- $\text{CH}_2$  groups of the piperidin  
25 ring, which exhibit a broad peak ranging from 3.5 to 4.5  
ppm.

30  $\delta$  0.8-2.15 (m, 19H), 2.22-2.4 (m, 1H), 2.75-2.98 (m, 2H),  
2.98-3.18 (m, 2H), 3.18-3.35 (m, 1H), 3.35-3.5 (qvar.,  
1H), 3.5-3.7 (m, 1H), 4.42-4.58 (d, 1H), 5.1 (s, 4H),  
7.1-7.5 (m, 10H).

$^{13}\text{C-NMR}$  (75 MHz,  $\text{CDCl}_3$ ): amidine and carbonyl carbons;  $\delta$   
154.96, 171.31, 174.82.

35

(iii)  $\text{BnOOC-CH}_2\text{-(R)Cgl-Pr -Pig(Z)}_2$

0.298 g (0.999 mmol)  $\text{BnOOC-CH}_2\text{-OTf}$  (see preparation of starting materials) was added to a mixture of 0.64 g (0.999 mmol)  $\text{H-(R)Cgl-Pro-Pig(Z)}_2$  and 0.531 g (2.996 mmol)  $\text{K}_2\text{CO}_3$  in 6.4 ml acetonitrile and heated to reflux. After 1 h 20 min the mixture was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and the solvent evaporated in vacuo to give 729 mg of a residue which was subjected to flash chromatography using ethylacetate as eluent. This gave two products: 120 mg of  $(\text{BnOOC-CH}_2)_2\text{-(R)Cgl-Pro-Pig(Z)}_2$  which eluted first from the column and 142 mg (18 %) of the title compound.

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ); Some signals, especially in the piperidin ring, are selectively broader due to an intramolecular exchange process. This is especially pronounced for the 2- and 6- $\text{CH}_2$  groups of the piperidin ring, which exhibit a broad peak ranging from 3.5 to 4.6 ppm.

$\delta$  0.94-2.27 (m, 19H), 2.28-2.43 (m, 1H), 2.8-2.98 (m, 2H), 2.98-3.06 (m, 1H), 3.06-3.15 (d, 1H), 3.15-3.25 (m, 1H), 3.3-3.5 (m, 4H), 4.5-4.61 (d, 1H), 5.1 (s, 6H), 7.1-7.6 (m, 15H), 10.52 (bs, 1H).

(iv)  $\text{HOOC-CH}_2\text{-(R)Cgl-Pro-Pig} \times 2 \text{ HCl}$

142 mg (0.176 mmol)  $\text{BnOOC-CH}_2\text{-(R)Cgl-Pro-Pig(Z)}_2$  was hydrogenated in the presence of 0.88 ml 1 M hydrochloric acid, 10 ml ethanol (99.5 %) and 180 mg 5 % Pd/C for 2 h. Removal of the catalyst by filtration on hyflo and millipore filter followed by evaporation of the solvent in vacuo and freeze drying gave 95 mg of  $\text{HOOC-CH}_2\text{-(R)Cgl-Pro-Pig} \times 2 \text{ HCl}$ . This crude material (79 % purity) was purified on RPLC using  $\text{CH}_3\text{CN}/0.1 \text{ M NH}_4\text{OAc}$  15/85 as eluent. Removal of the solvent and excess  $\text{NH}_4\text{OAc}$  by freeze drying, conversion to hydr chloric acid salt by dissolving in 1 M hydr chloric acid followed by freeze drying gave the title compound.

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O); δ 1.1-1.35 (m, 6H), 1.63-2.14 (m, 13H), 2.26-2.36 (m, 1H), 3.01-3.23 (m, 4H), 3.49-3.62 (qvar., 2H), 3.62-3.77 (m, 2H), 3.77-3.88 (apparent d, 2H), 4.18-4.32 (d, 1H), 4.37-4.5 (m, 1H).

5

Example 51**H-(R)Cha-Aze-Pig x 2 HCl**10      **(i) Boc-(R)Cha-Aze-Pig(Z)<sub>2</sub>**

To a well stirred mixture of 86 mg (0.243 mmol) Boc-(R)Cha-Aze-OH (See preparation of starting materials), 100 mg (0.236 mmol) H-Pig(Z)<sub>2</sub> (See preparation of starting materials) and 115 mg (0.944 mmol) DMAP in 5 ml CH<sub>3</sub>CN was added 50 mg (0.260 mmol) EDC and the reaction was stirred for 20 h at room temperature. The solvent was evaporated and the residue was dissolved in 70 ml EtOAc and the organic phase was washed with 3 x 5 ml 1 M KHSO<sub>4</sub>, 1 x 5 ml NaHCO<sub>3</sub>, 3 x 5 ml H<sub>2</sub>O, 1 x 5 ml brine and dried (MgSO<sub>4</sub>). Filtration and evaporation of the solvent gave 141 mg of an oil. The crude product was purified by flash chromatography (36 g SiO<sub>2</sub>) using a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH (97/3 followed by 95/5) to yield 43 mg (24 %) of the title compound.

25

**(ii) H-(R)Cha-Aze-Pig(Z)<sub>2</sub>**

Hydrogen chloride was bubbled through a mixture of 43 mg (0.0565 mmol) Boc-(R)Cha-Aze-Pig(Z)<sub>2</sub> in 10 ml of ethylacetate during 5 minutes. The solvent was evaporated in vacuo and ethyl acetate and 0.1 M NaOH-solution was added. The phases were separated and the organic phase was washed with water and brine and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was evaporated to give 38 mg, which was subjected to flash chromatography using 10 % NH<sub>3</sub>-saturated methanol in ethyl acetate as eluent to give 28 mg of the desired

35



product.

5  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ ); Some signals, especially in the piperidin ring, are selectively broader due to an intramolecular exchange process. This is especially pronounced for the 2- and 6- $\text{CH}_2$  groups of the piperidin ring, which exhibit a broad peak ranging from 3.7 to 4.5 ppm.

10  $\delta$  0.75-1.85 (m, 18H), 2.35-2.53 (m, 1H), 2.62-2.78 (m, 1H), 2.8-3.0 (m, 2H), 3.0-3.28 (m, 2H), 3.28-3.37 (m, 1H), 3.97-4.18 (m, 2H), 4.8-4.9 (m, 1H), 5.1 (s, 4H), 7.2-7.45 (m, 9H), 8.05-8.15 (m, 1H).

15 (iii) H-(R)Cha-Aze-Pig x 2 HCl

28 mg (0.042 mmol) H-(R)Cha-Aze-Pig(Z)<sub>2</sub> dissolved in 2 ml ethanol (99.5 %) and 0.13 ml hydrogen chloride (1 N) was hydrogenated in the presence of 35 mg 5 % Pd/C for 4 h. Removal of the catalyst by filtration and evaporation in vacuo of the solvent followed by dissolving in water and freeze drying gave 12 mg (60 %) of H-(R)Cha-Aze-Pig dihydrochloride.

25  $^1\text{H-NMR}$  (500 MHz, 300 K,  $\text{CD}_3\text{OD}$ ); Some signals, especially in the piperidin ring, are selectively broader due to an intramolecular exchange process. This is especially pronounced for the 2- and 6- $\text{CH}_2$  groups of the piperidin ring, which exhibit a broad peak ranging from 3.7 to 4.5 ppm.

30  $\delta$  0.75-2.1 (m, 18H), 2.2-2.35 (m, 1H), 2.62-2.75 (m, 1H), 3.0-3.12 (t, 2H), 3.12-3.23 (d, 2H), 3.85-3.95 (d, 2H), 3.95-4.0 (dd, 1H), 4.15-4.23 (m, 1H), 4.35-4.42 (m, 1H), 4.72-4.78 (m, 1H).

35

$^{13}\text{C-NMR}$  (75 MHz,  $\text{CD}_3\text{OD}$ ): guanidine:  $\delta$  157.6; carbonyl carbons:  $\delta$  170.0 and 172.6.

Example 52**HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pac x 2 HCl**5      **(i) Boc-(R)Cgl-Aze-Pac(Z)**

To a solution of 0.47 g (1.4 mmol) of Boc-(R)Cgl-Aze-OH  
(See preparation of starting materials), 0.40 g (1.4  
mmol) of H-Pac(Z) (See preparation of starting materials)  
10      and 0.67 g (5.5 mmol) of DMAP in 5 ml of acetonitril was  
added 0.27 g of EDC at 0°C. The mixture was stirred at  
room temperature over night and subsequently diluted  
with ethyl acetate. The solution was washed with KHSO<sub>4</sub>  
(aq) and NaHCO<sub>3</sub> (aq), dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and  
15      evaporated. Flash chromatography using ethyl acetate  
followed by ethyl acetate/methanol 98/2 as eluents gave  
0.25 g (30%) of the title compound as a mixture of 1,4-  
cis- and trans-products with respect to the Pac part of  
the molecule.

20

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>) : δ 0.8-2.0 (m, 29 H; thereof 1.45  
(s, 9 H)), 2.15 and 2.34 (m, 1 H, isomers), 2.45-2.7 (m,  
2 H), 3.0-3.4 (m, 2 H), 3.85 (m, 1 H), 4.14 (m, 1 H), 4.33  
(m, 1 H), 4.85 (m, 1 H), 4.98 (m, 1 H), 5.04 (s, 2 H),  
25      7.25-7.45 (m, 5 H), 7.8-7.9 (m, 1 H), 9.2-9.5 (m, 1 H).

**(ii) H-(R)Cgl-Aze-Pac(Z) x HCl**

Boc-(R)Cgl-Aze-Pac(Z), 0.25 g (0.41 mmol), was dissolved  
30      in 100 ml of ethyl acetate and cooled in an ice bath.  
HCl (g) was bubbled through for 5 min and the solvent  
was evaporated.

<sup>1</sup>H-NMR (300 MHz, MeOD) : δ 0.8-2.0 (m, 22 H), 2.05-2.35  
35      (m, 1 H), 2.4-2.55 (m, 1 H), 2.6-2.75 (m, 1 H), 3.00 (d,  
1 H), 3.05 and 3.37 (multiplets, 0.6 H and 0.4 H  
respectively, isomers), 3.15-3.3 (m, 1 H), 4.05-4.2 (m,

2 H), 4.88 (dd, 1 H), 5.11 (s, 2 H), 7.2-7.45 (m, 5 H), 8.0-8.15 (m, 1 H).

(iii)  $\text{BnO}_2\text{C}-\text{CH}_2-(\text{R})\text{Cgl-Aze-Pac}(\text{Z})$

5

A mixture of 0.17 g (0.33 mmol) of  $\text{H}-(\text{R})\text{Cgl-Aze-Pac}(\text{Z}) \times \text{HCl}$ , 0.11 g (0.37 mmol) of benzyl triflyloxyacetate and 0.14 g (1.0 mmol) of  $\text{K}_2\text{CO}_3$  in 5 ml of acetonitrile was stirred at room temperature for 3 days. The crude material was flash chromatographed with  $\text{EtOAc}/\text{CH}_2\text{Cl}_2/\text{MeOH}$  95/20/5. Yield: 70 mg (32 %).

$^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ ) :  $\delta$  0.85-2.3 (m, 20 H), 2.48 (m, 1 H), 2.63 (m, 1 H), 2.87 (m, 1 H), 3.05-3.25 (m, 1 H), 3.25-3.35 (m, 2 H), 3.38 (dd, 1 H), 3.95 (m, 1 H), 4.08 (m, 1 H), 4.88 (m, 1 H), 5.1-5.2 (m, 4 H), 5.9-6.3 (m, 1 H), 7.25-7.5 (m, 10 H), 8.00 and 8.08 (broad triplets, 1 H, isomers).

20 (iv)  $\text{HO}_2\text{C}-\text{CH}_2-(\text{R})\text{Cgl-Aze-Pac} \times 2 \text{ HCl}$

$\text{BnO}_2\text{C}-\text{CH}_2-(\text{R})\text{Cgl-Aze-Pac}(\text{Z})$ , 70 mg (0.11 mmol), was dissolved in 5 ml of ethanol, and 5% Pd/C and 0.1 ml of conc. HCl were added. The mixture was hydrogenated at atmospheric pressure for 1 h. After filtration and evaporation the product was purified through preparative RPLC using 0.1 M  $\text{NH}_4\text{OAc}/\text{CH}_3\text{CN}$  4/1 as eluent. After change of salt to the hydrochloride and freeze drying the title compound was obtained as a 45/55 mixture of 1,4-cis- and trans-isomers with respect to the Pac part of the molecule. Yield: 40 mg (74%).

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ )  $\delta$  1.1-2.1 (m, 20 H), 2.32 (m, 1 H), 2.52 (m, 1 H), 2.63 (m, 1 H), 2.72 (m, 1 H), 3.1-3.3 (m, 1 H), 3.40 (m, 1 H), 3.8-3.95 (m, 2 H), 4.04 (d, 1 H), 4.39 (m, 1 H), 4.93 (m, 1 H).

$^{13}\text{C}$ -NMR (125 MHz,  $\text{D}_2\text{O}$ ) amidine and carbonyl carbons:  $\delta$  167.7, 172.0, 174.9 and 175.2.

Example 53

5

**H-(R)Cha-Pro-Pac x 2 HCl**

(i) Boc-(R)Cha-Pro-Pac(Z)

10 211 mg (1.1 mmol) EDC was added at 0°C to a stirred solution of 0.4 g (1.1 mmol) H-Pac(Z) x 2 HCl (See preparation of starting materials), 0.4 g (1.1 mmol) Boc-(R)Cha-Pro-OH(See preparation of starting materials), and 0.55 g DMAP in 7 ml acetonitrile. The  
15 reaction mixture was stirred at 0°C for 1 h and at room temperature for 2 h. The solvent was removed in vacuo and the residue was diluted with ethyl acetate and water. The organic phase was washed with acetic acid, water, and sodium hydrogen carbonate solution and dried  
20 ( $\text{MgSO}_4$ ). Removal of the solvent in vacuo gave a residue which was purified by flash chromatography using ethyl acetate as eluent to give 196 mg (27%) of the title compound.

25 (ii) H-(R)Cha-Pro-Pac(Z)

Hydrogen chloride was bubbled through a solution of 196 mg Boc-(R)Cha-Pro-Pac(Z) in 25 ml ethyl acetate. After  
10 minutes the reaction mixture was diluted with methylene chloride and sodium hydroxide solution was  
30 added. The aqueous phase was extracted several times with methylene chloride and the combined organic phases were dried ( $\text{K}_2\text{CO}_3$ ) and the solvent was removed in vacuo to give 86 mg (52%) of the title compound.

35

(iii) H-(R)Cha-Pro-Pac x 2 HCl

The title compound was prepared by hydrogenation of H-(R)Cha-Pro-Pac(Z) in ethanol in the presence of 10% Pd/C.

- 5 H-NMR (300 MHz, D<sub>2</sub>O; A ca: 1:1 mixture of 1,4-cis- and 1,4-trans isomers in the Pac part of the molecule);  $\delta$  1.15-1.3 (q), 1.6-1.85 (m), 1.9-2.0 (m), 2.0-2.1 (d), 2.1-2.15 (m), 2.15-2.2 (m), 2.65-2.7 (m), 2.7-2.8 (m), 2.95-3.0 (d), 3.15-3.2 (d), 5.4 (s), 7.45-7.55 (m).

10

Example 54

H-(R)Cgl-Ile-Pab x 2 HCl

- 15 (i) Boc-(R)Cgl-Ile-Pab(Z)

To a stirred mixture of 1.33 g (3.6 mmol) Boc-(R)Cgl-Ile-OH (See Preparation of starting materials), 1.12 g (3.9 mmol) H-Pab(Z) (See Preparation of starting materials) and 1.76 g (14.4 mmol) DMAP in 50 ml CH<sub>3</sub>CN/DMF (1/1) was added 0.75 g (3.9 mmol) EDC at + 5°C. The reaction mixture was allowed to reach room temperature and left for 60 h. The CH<sub>3</sub>CN was removed by evaporation and the residue was poured out in 100 ml water (a yellow precipitate was formed). The mixture was extracted with 2 x 50 ml EtOAc and the combined organic phase was washed with 2 x 30 ml NaHCO<sub>3</sub>(saturated), 2 x 50 ml 0.2 M HCl, 1 x 50 ml Brine and dried (MgSO<sub>4</sub>). Evaporation followed by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/THF (85/15) as eluent gave 510 mg (24 %) of the title compound.

20

25

30

(ii) H-(R)Cgl-Ile-Pab(Z)

- 35 530 mg Boc-(R)Cgl-Ile-Pab(Z) was dissolved in 14 ml CH<sub>2</sub>Cl<sub>2</sub>/TFA (2.5/1) and stirred for 2 h at room temperature. Evaporation of the solvent followed by

flash chromatography using  $\text{CH}_2\text{Cl}_2/\text{MeOH}(\text{NH}_3\text{-saturated})$  (95/5) as eluent gav the title compound.

(iii) H-(R)Cgl-Ile-Pab x 2 HCl

5

75 mg (0.14 mmol) H-(R)Cgl-Ile-Pab(Z) was hydrogenated over 10 % Pd/C in 5 ml EtOH, which contained an excess HCl(g) to give the dihydrochloride, at atmospheric pressure for 6 h. Addition of 2 g activated charcoal and 20 ml EtOH followed by filtration through celite, evaporation of the solvent and freeze drying from water gave 50 mg (89%) of the title compound as a white powder.

15  $^1\text{H-NMR}$ (500 MHz, MeOD):  $\delta$  0.90 (t, 3H), 0.94 (d, 3H), 1.1-2.0 (m, 14H), 3.83 (bs, 1H), 4.26 (d, 1H), 4.50 (m, 2H), 7.57 (bd, 2H), 7.78 (bd, 2H).

Example 55

20

H-(R)Cgl-Aze-Pab

Hydrogenation of 257 mg (5.08 mmol) H-(R)Cgl-Aze-Pab(Z) (See Example 1 (ii)) over 5 % Pd/C in 6 ml EtOH/ $\text{H}_2\text{O}$  at atmospheric pressure for 6 h followed by filtration of the catalyst, evaporation of the solvent and freeze drying from water gave 200 mg (89 %) of the title compound.

30  $^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  1.0-2.0 (m, 11H), 2.25 (m, 1H), 2.70 (m, 1H), 3.30 (m, 1H), 3.75 (m, 1H), 4.30 (m, 1H), 4.45 (m, 1H), 4.55 (m, 2H), 7.60 (m, 2H), 7.77 (m, 2H).

MS m/z 372 ( $\text{M}^+ + 1$ ).

35

Example 56

HOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab x HOAc

5 (i) BnOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab(Z)

0.250 g (0.47 mmol) H-(R)Cha-Pro-Pab(Z) (See Example 15) dissolved in 5 ml CH<sub>2</sub>Cl<sub>2</sub> was cooled to -10°C and 150 mg (0.48 mmol) of TfOCH<sub>2</sub>COOBn (See Preparation of Starting materials) dissolved in 3 ml CH<sub>2</sub>Cl<sub>2</sub> was added slowly. 200 mg (1.45 mmol) of potassium carbonate was added and the mixture was stirred at room temperature for 20 h. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub>, extracted with water and dried (MgSO<sub>4</sub>). Evaporation of the solvent followed by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH 9/1 as eluent gave 150 mg (46%) of the title compound.

(ii) HOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab x HOAc

20 150 mg (0.2 mmol) BnOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab(Z) was hydrogenated over 50 mg 5 % Pd/C in 20 ml EtOH at atmospheric pressure for 4 h. Filtration of the catalyst, evaporation of the solvent followed by purification by RPLC, using CH<sub>3</sub>CN/0.1 M NH<sub>4</sub>OAc 1/4 as eluent, gave 35 mg (37%) of the title compound.

<sup>1</sup>H-NMR (500 MHz, MeOD): δ 1.00 (m, 1H), 1.20-1.45 (m, 5H), 1.5 (m, 1H), 1.6-1.8 (m, 6H), 1.9-2.1 (m, 6H), 2.25 (m, 1H), 3.25 (m, 1H), 3.5 (m, 1H), 3.85 (m, 1H), 4.15 (m, 1H), 4.35-4.6 (m, 3H), 4.9 (m, partially hidden by the HOD line, 6H), 7.55 (d, 2H), 7.75 (d, 2H).

Example 57

35 MeOOC-CH<sub>2</sub>-(R)Cgl-Az -Pab x 2 HCl

(i) MeOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)

0.186 g (0.841 mmol) TfO-CH<sub>2</sub>-COOMe (See preparation of starting materials) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and slowly added to a mixture of 0.425g (0.841 mmol) H-(R)Cgl-Aze-Pab(Z) (See Example 1), 0.894 g (5.04 mmol) K<sub>2</sub>CO<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> (totally 4.3 ml) at roomtemperatur, and stirred over night. More CH<sub>2</sub>Cl<sub>2</sub> was added and the mixture was washed with water and brine, dried, filtered and the solvent was evaporated in vacuo to give 0.51 g of a residue that was three times subjected to flash chromatography on silica gel using, first CH<sub>2</sub>Cl<sub>2</sub>/THF/MeOH (16/4/1), then CH<sub>2</sub>Cl<sub>2</sub>/THF(2%NH<sub>3</sub>) (8/2) and the last time diethylether/MeOH(NH<sub>3</sub>-saturated) (95/5) as eluent. This gave 0.324 g (67 %) of the title compound.

15

(ii) MeOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab x 2 HCl

220 mg (0.38 mmol) MeOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z) was hydrogenated in the presence of 1.14 ml 1 N HCl, 6.5 ml MeOH and 300 mg Pd/C for 2 h. Removal of the catalyst by filtration on cellite and millipore filter followed by evaporation of the solvent in vacuo and freeze drying twice gave 178 mg (91 %) of the title compound.

<sup>1</sup>H-NMR (500 MHz, D<sub>2</sub>O); δ 1.12-1.4 (m, 5H), 1.68-1.81 (m, 2H), 1.81-1.9 (m, 3H), 1.97-2.1 (m, 1H), 2.29-2.4 (m, 1H), 2.68-2.8 (m, 1H), 3.86 (s, 3H), 4.1 (s, 2H), 4.1-4.5 (d, 1H), 4.36-4.42 (t, 2H), 4.59 (s, 2H), 4.99-5.04 (m, 1H), 7.65-7.7 (d, 2H), 7.8-7.85 (d, 2H).

30

<sup>13</sup>C-NMR (75 MHz, MeOD): amidine and carbonyl carbons; δ 146.78, 167.68, 168.15, 172.29.

#### Example 58

35

EtOOC-CH<sub>2</sub>-(R)Cgl-Az -Pab x 2 HCl



(i)  $\text{EtOOC-CH}_2\text{-(R)Cgl-Az -Pab(Z)}$ 

0.208 g (0.876 mmol)  $\text{TfO-CH}_2\text{-COOEt}$  (See preparation of starting materials) was dissolved in  $\text{CH}_2\text{Cl}_2$  and slowly added to a mixture of 0.443 g (0.876 mmol)  $\text{H-(R)Cgl-Aze-Pab(Z)}$  (See Example 1) and 0.931 g (5.26 mmol)  $\text{K}_2\text{CO}_3$  in  $\text{CH}_2\text{Cl}_2$  (totally 4 ml) cooled on an ice-bath. After 2 h the ice-bath was removed and stirring was continued at roomtemperature for 2 hours. More  $\text{CH}_2\text{Cl}_2$  was added and the mixture was washed with water and brine, dried, filtered and the solvent was evaporated in vacuo to give 0.51 g of a residue that was subjected to flash chromatography using diethylether/ $\text{MeOH}(\text{NH}_3\text{-saturated})$  (95/5) as eluent. This gave 0.387 g (75 %) of the title compound.

(ii)  $\text{EtOOC-CH}_2\text{-(R)Cgl-Aze-Pab x 2 HCl}$ 

395 mg (0.668 mmol)  $\text{EtOOC-CH}_2\text{-(R)Cgl-Aze-Pab(Z)}$  was hydrogenated in the presence of 12 ml  $\text{EtOH}$  (99.5 %) and 390 mg  $\text{Pd/C}$  for 5 h. Removal of the catalyst by filtration on cellite and millipore filter, followed by evaporation of the solvent in vacuo and freeze drying twice, gave 281 mg (88 %) of  $\text{EtOOC-CH}_2\text{-(R)Cgl-Aze-Pab}$ . 2 equivalents of 1 N  $\text{HCl}$  was added, and freeze drying three times gave 288 mg (81 %) of the title compound.

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ );  $\delta$  1.05-1.48 (m, 8H), 1.6-2.05 (m, 6H), 2.15-2.33 (m, 1H), 2.58-2.79 (m, 1H), 3.89-4.0 (m, 3H), 4.2-4.33 (m, 3H), 4.33-4.44 (m, 1H), 4.44-4.66 (m, 2H), 4.91 (m, 1H (partially hidden by the H-O-D signal)), 7.54-7.63 (d, 2H), 7.72-7.84 (d, 2H).

Example 59

$^n\text{BuOOC-CH}_2\text{-(R)Cgl-Aze-Pab x HOAc}$

(i) <sup>n</sup>BuOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)

Prepared in the same way as described for <sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z) (See Example 60 (i)) using TfO-CH<sub>2</sub>-COO<sup>n</sup>Bu as alkylating agent. The crude product was purified by flash chromatography twice, first using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95/1) as eluent and then CH<sub>2</sub>Cl<sub>2</sub>/i-propylalcohol (90/7) to give 324 mg (47 %) of the title compound.

(ii) <sup>n</sup>BuOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab x HOAc

The deprotection was done according to the procedure described in Example 57 (ii). The crude material was purified on RPLC using CH<sub>3</sub>CN (30 %) in 0.05 M NH<sub>4</sub>OAc and 0.05 M HOAc as eluent to give 100 mg (53 %) of the title compound.

<sup>1</sup>H-NMR (300 MHz, MeOD); δ 0.85-2.1 (m, 18H), 2.15-2.37 (m, 1H), 2.58-2.8 (m, 1H), 3.7-5.0 (m, 10H), 4.88-5.0 (partially hidden by the H-O-D signal), 7.46-7.65 (d, 2H), 7.71-7.88 (d, 2H).

<sup>13</sup>C-NMR (75 MHz, MeOD): amidine and carbonyl carbons; δ 146.8, 168.12, 168.2, 172.2.

Example 6030 <sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab x 2 HCl(i) <sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)

0.402 g (1.375 mmol) TfO-CH<sub>2</sub>-COO<sup>n</sup>Hex (See Preparation of starting materials) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and slowly added to a mixture of 0.695 g (1.375 mmol) H-(R)Cgl-Aze-Pab(Z) (See Example 1), 1.463 g (8.25 mmol) K<sub>2</sub>CO<sub>3</sub> in

CH<sub>2</sub>Cl<sub>2</sub> (totally 4 ml) at <-10°C. After 1 h the CO<sub>2</sub>-ice-bath was removed and stirring was continued at room temperature for 45 minutes. More CH<sub>2</sub>Cl<sub>2</sub> was added and the mixture was washed with water and brine, dried, filtered and the solvent was evaporated in vacuo to give 0.828 g of a residue, which was twice subjected to flash chromatography, first using diethylether/MeOH(NH<sub>3</sub>-saturated) (95/5), and then CH<sub>2</sub>Cl<sub>2</sub>/MeOH(NH<sub>3</sub>-saturated) (95/5) as eluent. This gave 0.42 g (47 %) of the title compound.

(ii) <sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab x 2 HCl

Hydrogenation of 400 mg (0.617 mmol) <sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(2) in the presence of 12 ml THF and 400 mg Pd/C for 1.5 h did not give complete de-protection. The hydrogenation was completed in 4 h in the presence of 1.7 ml 1 N HCl, 12 ml MeOH and 340 mg Pd/C. Removal of the catalyst by filtration on cellite and millipore filter, followed by evaporation of the solvent in vacuo and freeze drying twice, gave 287 mg (79 %) of the title compound.

<sup>1</sup>H-NMR (300 MHz, MeOD); δ 0.8-2.13 (m, 22H), 2.13-2.31 (m, 1H), 2.61-2.81 (m, 1H), 3.93-4.15 (m, 3H), 4.15-4.37 (m, 3H), 4.37-4.7 (m, 3H), 4.88-5.0 (m, 1H (partially hidden by the H-O-D signal)), 7.52-7.69 (d, 2H), 7.75-7.9 (d, 2H).

<sup>13</sup>C-NMR (75 MHz, MeOD): amidine and carbonyl carbons; δ 146.84, 167.67, 167.84, 172.17.

#### Example 61

H-(R)Cgl-Pro-Pac x 2 HCl

## (i) Boc-(R)Cgl-Pro-Pac(Z)

377 mg (1.97 mmol) EDC was added at 0°C to a stirred solution of 708 mg (1.95 mmol) of Boc-(R)Cgl-Pro-OH (See Preparation of starting material), 714 mg (2.0 mmol) H-Pac(Z) x 2 HCl (See Preparation of starting material) and 1.078 g (8.8 mmol) DMAP in 12.5 ml acetonitrile. The reaction mixture was allowed to reach room temperature over night. The solvent was removed in vacuo and the residue was first purified by flash chromatography, using 10 % methanol in methylene chloride as eluent, and subsequently by RPLC. Two fractions (51 mg and 150 mg) giving MS m/z = 626 (M + 1) were isolated.

## (ii) H-(R)Cgl-Pro-Pac(Z)

Hydrogen chloride was bubbled into a solution of 141 mg (0.22 mmol) Boc-(R)Cgl-Pro-Pac(Z) in 50 ml ethyl acetate. After 15 minutes 10 % sodium carbonate solution was added and the organic phase was separated and dried (K<sub>2</sub>CO<sub>3</sub>). Evaporation of the solvent gave 71 mg (61 %) of the product.

## (iii) H-(R)Cgl-Pro-Pac x 2 HCl

A mixture of 71 mg (0.14 mmol) H-(R)Cgl-Pro-Pac(Z) and a small spatula of 10 % Pd/C in 10 ml of ethanol was hydrogenated at room temperature and atmospheric pressure for 2 h. The catalyst was removed by filtration and the solvent was removed in vacuo. The residue was dissolved in 50 ml water and 0.6 g 1M hydrochloric acid. Freeze drying yielded 38 mg (58 %) of the title compound

MS m/z 392 (M + 1)

Example 62

**HOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac x HOAc**

5      (i) BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac(Z)

A mixture of 84 mg (0.15 mmol) H-(R)Cha-Pro-Pac(Z) (See Example 53 (ii)), one spatula of potassium carbonate, and 47 mg of TfOCH<sub>2</sub>-COOBn (See Preparation of starting materials) in 3 ml of methylene chloride was stirred at room temperature over night. The reaction mixture was filtered and the solvent was removed in vacuo to give a residue which was subjected to flash chromatography using ethyl acetate/methylene chloride/methanol 95:20:5 as eluent. 29 mg of the desired product was isolated.

(ii) HOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac x HOAc

A mixture of 29 mg BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac(Z) and 37 mg of 10 % Pd-C in 5 ml ethanol was stirred for 4 h at room temperature and atmospheric pressure. Filtration of the catalyst followed by removal of the solvent and purification by RPLC gave the desired compound.

25      MS m/z = 464 (M + 1).

Example 63

**HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pac**

30

(i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pac(Z)

A solution of 0.35 g (0.64 mmol) H-(R)Cgl-Pro-Pac(Z) (See Example 61 (ii) ), 124 mg (0.76 mmol) benzyl acrylate, and 280 µl (2 mmol) triethyl amine in 1 ml ethanol was kept at room temperature for 3 days. Removal of the solvent followed by purification by HPLC gave 18

mg (4 %) of the title compound.

(ii)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cgl-Pro-Pac}$

- 5 A mixture of 18 mg  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cgl-Pro-Pac(Z)}$  and a small spatula of 10 % Pd/C was hydrogenated for 2 h at room temperature and atmospheric pressure in EtOH. Filtration followed by removal of the solvent in vacuo and dissolution in water and freeze drying gave 7 mg (78 %)  
10 of the title compound. MS  $m/z = 464 (M + 1)$ .

Example 64

$\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pac}$

15

(i)  $\text{Boc-(R)Cha-Aze-Pac(Z)}$

- A solution of 0.4 g (1.38 mmol) H-Pac(Z) (See Preparation of starting material of H-Pac(Z) x 2 HCl),  
20 0.5 g (1.41 mmol) Boc-(R)Cha-Aze-OH (See Preparation of starting material), and 0.67 g (5.5 mmol) DMAP in 20 ml acetonitrile was mixed at 0°C with a solution of 0.26 g (1.4 mmol) EDC in 15 ml acetonitrile. The reaction mixture was kept at room temperature over night and the  
25 solvent was subsequently removed in vacuo. The residue was partitioned between ethyl acetate and water. The aqueous phase was extracted once more with ethyl acetate and the combined organic phases were washed with sodium hydrogen sulphate solution, sodium carbonate solution,  
30 and brine and then dried (sodium sulphate). Evaporation of the solvent gave 0.54 g (63 %) of the title compound.

(ii)  $\text{H-(R)Cha-Aze-Pac(Z)}$

- 35 Hydrogen chloride was bubbled into a solution of 0.54 g (0.9 mmol) Boc-(R)Cha-Aze-Pac(Z) in ethyl acetate. The solution was kept in the refrigerator over night and

the solvent was then removed in vacuo and the residue was dissolved in ethyl acetate. The solution was washed with aqueous sodium hydrogen carbonate solution, water and brine and dried (sodium sulphate). Removal of the solvent gave 0.35 g (77 %) of the product.

(iii)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pac(Z)}$

A solution of 180 mg (0.33 mmol) H-(R)Cha-Aze-Pac(Z) and 53 mg (0.33 mmol) benzyl acrylate in ethanol was kept at room temperature for 60 h. The solvent was removed in vacuo and the residue was dissolved in ethyl acetate. The solution was washed with potassium hydrogen sulphate solution and sodium hydrogen carbonate solution and brine. Drying (sodium sulphate) and removal of the solvent in vacuo gave a residue which was purified by flash chromatography, using 10 % methanol in methylene chloride as eluent to yield 150 mg (66 %) of the title compound

(iv)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pac} \times 2 \text{ HCl}$

A mixture of 115 mg  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Pac(Z)}$  and 67 mg of 10 % Pd-C in 10 ml ethanol was hydrogenated for 1.5 h at room temperature and atmospheric pressure. Filtration followed by removal of the solvent in vacuo and dissolution of the residue in water and 1.5 ml of 1M hydrochloric acid gave a solution which was freeze dried to give 30 mg (33 %) of the title compound.

MS m/z 464 (M + 1).

Example 65

$\text{HOOC-CH}_2\text{-(R)Cha-Aze-Pig} \times 2 \text{ HCl}$

(i)  $\text{Boc-(R)Cha-Aze-Pig(Z)}$

0.249 g (1.298 mmol) EDC was added at  $<-15^{\circ}\text{C}$  to a mixture of 0.473 g (1.236 mmol) Boc-(R)Cha-Az -OH (See Preparation of starting materials), 0.404 g (1.236 mmol) H-Pig(Z) x HCl (See Preparation of starting materials) and 0.604 g (4.94 mmol) DMAP in 13.5 ml DMF. The temperature was allowed to rise to roomtemperature over night. The solvent was evaporated in vacuo and EtOAc and 2 M  $\text{KHSO}_4$  was added. The phases were separated and the organic phase was washed with saturated  $\text{Na}_2\text{CO}_3$  and brine. Repeting the extractive procedure, drying ( $\text{Na}_2\text{SO}_4$ ), filtration and evaporation of the solvents gave 0.612 g of a residue which was subjected to flash chromatography using EtOAc/MeOH 9/1 as eluent. This gave 407 mg (53 %) of the title compound.

(ii) H-(R)Cha-Aze-Pig(Z)

0.4 g (0.638 mmol) of Boc-(R)Cha-Aze-Pig(Z) was dissolved in 24.4 ml of TFA/ $\text{CH}_2\text{Cl}_2$  1/4, stirred for 30 minutes on an ice-bath, and for 30 minutes at roomtemperature. The solvent was removed in vacuo and EtOAc and saturated  $\text{Na}_2\text{CO}_3$  was added. The phases were seperated and the organic layer was washed with water and brine, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and the solvent was evaporated in vacuo to give 336 mg (100 %) of the title compound.

(iii)  $\text{BnOOC-CH}_2\text{-(R)Cha-Aze-Pig(Z)}$

89 ml (0.562 mmol)  $\text{BnOOC-CH}_2\text{-Br}$  was slowly added to a mixture of 0.296 g (0.562 mmol) H-(R)Cha-Aze-Pig(Z) and 0.171 g (1.236 mmol)  $\text{K}_2\text{CO}_3$  in 6 ml  $\text{CH}_3\text{CN}$  heated to  $60^{\circ}\text{C}$  on an oilbath. After 1 h 45 minutes the solvent was evaporated, EtOAc was added, and th mixtur was washed with water, dri d ( $\text{Na}_2\text{SO}_4$ ), filtered and th solvent was evap rat d in vacu t give 346 mg of a



residue which was subjected to flash chromatography using  $\text{CH}_2\text{Cl}_2/\text{THF}/\text{MeOH}$  (8/2/1) as eluent. This gave 297 mg (78 %) of the title compound.

5 (iv)  $\text{HOOC}-\text{CH}_2-(\text{R})\text{Cha-Aze-Pig} \times 2 \text{ HCl}$

243 mg (0.36 mmol)  $\text{BnOOC}-\text{CH}_2-(\text{R})\text{Cha-Aze-Pig}(\text{Z})$  was hydrogenated in the presence of 1.7 ml 1 N HCl, 10 ml EtOH (99.5 %) and 300 mg Pd/C for 2 h. Removal of the catalyst by filtration on cellite and millipore filter followed by evaporation of the solvent in vacuo and freeze drying twice gave 166 mg (88 %) of the title compound

15  $^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ );  $\delta$  0.6-1.9 (m, 18H), 2.1-2.27 (m, 1H), 2.52-2.76 (m, 1H), 2.82-3.2 (m, 4H), 3.46-3.61 (m, 1H), 3.61-3.81 (m, 2H), 3.81-4.0 (m, 2H), 4.0-4.24 (m, 2H), 4.24-4.4 (m, 1H).

20 Example 66

$\text{HOOC}-\text{CH}_2-(\text{R})\text{Cha-Pro-Pig} \times 2 \text{ HCl}$

(i)  $\text{Boc}-(\text{R})\text{Cha-Pro-Pig}(\text{Z})$

25

To a mixture of 0.3495 g (0.95 mmole)  $\text{Boc}-(\text{R})\text{Cha-Pro-OH}$  (See Preparation of starting materials), 0.464 g (3.8 mmole) DMAP, 0.310 g (0.95 mmole)  $\text{H-Pig}(\text{Z}) \times \text{HCl}$  (See Preparation of starting materials) in 5 ml  $\text{CH}_2\text{Cl}_2$  was added 0.192 g (1 mmole) of EDC and the mixture was stirred over night at room temperature. The mixture was evaporated and the residue was dissolved in ethyl acetate. The organic phase was washed twice with 0.3 M  $\text{KHSO}_4$  and once with brine. The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. The crude product was purified by flash chromatography using a stepwise gradient of  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (100/0, 97/3,

95/5, 90/10) as eluent to yield 307 mg of the title compound.

(ii) H-(R)Cha-Pro-Pig(Z)

5  
0.306 g (0.48 mmole) Boc-(R)Cha-Pro-Pig(Z) was dissolved in 30 ml HCl saturated ethyl acetate. The mixture was allowed to stand for half an hour. The solvent was evaporated and the residue was dissolved  
10 in CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed twice with 0.2 M NaOH. The combined water layer was extracted once with CH<sub>2</sub>Cl<sub>2</sub> and the combined organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to yield 257 mg (99%) of the title compound.

15  
(iii) BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z)

A mixture of 0.256 g (0.473 mmole) H-(R)Cha-Pro-Pig(Z), 0.144 g (1.04 mmole) K<sub>2</sub>CO<sub>3</sub> and 82 µl (0.521  
20 mmole) of benzylbromoacetate in 6 ml acetonitrile was heated to 60°C for two hours under stirring. The solvent was evaporated and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, washed once with water and once with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and the solvent evaporated.  
25 The crude product was purified by flash chromatography using a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH ( 97/3, 95/5, 90/10 ) as eluent to yield 0.2 g product (90% pure according to RPLC). The final purification was made on a chromatotron (Harrison  
30 research, model 7924T ) on a 2mm silica plate in CH<sub>2</sub>Cl<sub>2</sub>/MeOH 95/5 yielding 0.158 g (48%) of the pure product.

35  
(iv) HOOC-CH<sub>2</sub>-(R)Cha-Pro-Pig x 2 HCl

0.158 g (0.227 mmole) of BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z) was mixed with 0.075 g Pd/C (5%), 1.0 ml 1N HCl-

solution and 10 ml ethanol. The mixture was hydrogenated at atmospheric pressure for one hour. Filtration through cellite and evaporation of the solvent followed by freeze drying twice from water gave 119 mg (97%) of the product

<sup>1</sup>H-NMR (D<sub>2</sub>O, 300 MHz): δ 0.95-1.44 (m, 7H), 1.52 (m, 1H), 1.60-2.20 (m, 13H), 2.39 (m, 1H), 3.07-3.32 (m, 4H), 3.68 (m, 1H), 3.77-4.02 (m, 5H; thereof 3.98 (s, 2H), 4.44-4.58 (m, 2H)

<sup>13</sup>C-NMR (D<sub>2</sub>O, 75 MHz): carbonyl- and guanidine carbons: δ 156.5, 168.3, 169.6, 174.5

15 Example 67

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pig x 2 HCl

(i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z)

20 0.297 g (0.55 mmole) H-(R)Cha-Pro-Pig(Z) (See Example 66 (ii)) was dissolved in 2 ml ethanol and 90 μl (0.59 mmole) benzylacrylate was added. The reaction mixture was stirred for four days at room temperature. The solvent was evaporated and the crude product chromatographed on a chromatotron (Harrison research, model 7924T) using a 2mm silica plate with a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95/5, 90/10) as eluent to yield 0.338 g (87%) of the title compound.

30 (ii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pig x 2 HCl

0.238 g (0.227 mmole) of BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z) was mixed with 0.120 g Pd/C (5%), 1.2 ml 1N HCl-solution and 15 ml ethanol. The mixture was hydrogenated at atmospheric pressure for one hour. Filtration of the catalyst through cellite,

evaporation of the solvent followed by freeze drying twice from water gave 178 mg (95 %) of the title compound.

5  $^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ , 300 MHz):  $\delta$  0.82-1.45 (m, 8H), 1.45-2.15 (m, 13H), 2.29 (m, 1H), 2.83 (t, 2H), 2.9-3.4 (m, 6H), 3.57 (bq, 1H), 3.67-3.87 (m, 3H), 4.25-4.43 (m, 2H)

10  $^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ , 75 MHz): carbonyl- and guanidine carbons:  $\delta$  156.3, 168.2, 174.3, 174.6

MS  $m/z$  479 ( $\text{M}^+ + 1$ )

15 Example 68

$(\text{HOOC-CH}_2)_2\text{-(R)Cgl-Pro-Pig} \times 2 \text{ HCl}$

120 mg (0.126) mmol ( $\text{BnOOC-CH}_2$ )<sub>2</sub>-(R)Cgl-Pro-Pig(Z)<sub>2</sub>  
20 (See Example 50 (iii) ) was hydrogenated in the presence of 0.75 ml 1 N HCl, 7 ml EtOH (99.5 %) and 150 mg Pd/C for 4 h. Removal of the catalyst by filtration on cellite and millipore filter and evaporation of the solvent in vacuo followed by  
25 freeze drying gave 66 mg (90 %) of the title compound

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ );  $\delta$  1.05-1.38 (m, 7H), 1.53-1.64 (d, 1H), 1.64-2.14 (m, 11H), 2.27-2.39 (m, 1H), 3.03-3.28 (m, 4H), 3.58-3.70 (m, 1H), 3.7-3.8 (m, 1H),  
30 3.8-3.9 (d, 2H), 4.07-4.22 (m, 2H), 4.22-4.35 (m, 1H), 4.38-4.5 (m, 1H).

$^{13}\text{C-NMR}$  (75 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons;  
 $\delta$  156.28, 166.73, 170.14, 174.01.

35

Example 69

**HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(HOOC-CH<sub>2</sub>)-(R)Cha-Pr -Pig x 2 HCl**

**(i) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(BnOOC-CH<sub>2</sub>)-(R)Cha-Pro-Pig(Z)**

5 To a cold (ice-bath temperature) mixture of 100 mg  
(0.14 mmol) BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z) (See  
Example 67 (i)) and 80 mg (0.57 mmol) potassium  
carbonate in 4 ml of CH<sub>2</sub>Cl<sub>2</sub> was carefully added a  
10 solution of 64 mg (0.21 mmol) TfO-CH<sub>2</sub>-COOBn dissolved  
in 1 ml CH<sub>2</sub>Cl<sub>2</sub>. The reaction mixture was left at 0°C  
for 30 minutes and then allowed to reach room  
temperature for 2 h after which it was heated to  
reflux for 30 minutes and finally left over night at  
room temperature. Evaporation of the solvent followed  
15 by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (97/3) as  
eluent afforded 65 mg (54 %) of the title compound.

**(ii) HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(HOOC-CH<sub>2</sub>)-(R)Cha-Pro-Pig x 2 HCl**

20 65 mg (0.08 mmol) of BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(BnOOC-CH<sub>2</sub>)-(R)Cha-  
Pro-Pig(Z) was dissolved in 10 ml of EtOH/1M HCl  
(9/1) and hydrogenated over 10 % Pd/C for 3 h at  
atmospheric pressure. Filtration of the catalyst  
evaporation of the solvent followed by freeze drying  
25 from water gave 40 mg (97 %) of the title compound as  
a white powder.

<sup>13</sup>C-NMR (125 MHz, MeOD): amidine and carbonyl carbons:  
δ 157.5, 167.2, 169.1, 173.7 and 174.1.

30

**Example 70**

**HOOC-CH<sub>2</sub>-(R)Cgl-Aze-(R,S)Itp x 2 HCl**

35 **(i) Boc-(R)Cgl-Az -(R,S)Itp(Ts)**

**B c-(R)Cgl-Aze-OH (See Pr parati n f starting**

materials) (400 mg, 1.17 mmol), H-(R,S)Itp(Ts) (See Preparation of starting materials) (366 mg, 1.23 mmol) and DMAP (286 mg, 2.34 mmol) was dissolved in CH<sub>3</sub>CN (6 ml) and cooled to 5°C. EDC (236 mg, 1.23 mmol) was added and the resulting mixture was stirred at room temperature over night. The CH<sub>3</sub>CN was removed and the residue was dissolved in MeOH/EtOAc/H<sub>2</sub>O. The separated organic layer was washed with K<sub>2</sub>CO<sub>3</sub>(sat), 2 M KHSO<sub>4</sub>, brine and dried(Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent resulted in a white solid, 688 mg (85%).

MS m/z 620 (M<sup>+</sup> + 1)

(ii) H-(R)Cgl-Aze-(R,S)Itp(Ts)

Boc-(R)Cgl-Aze-(R,S)Itp(Ts) (500 mg, 0.8 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 ml) and HCl(g) was bubbled through the solution for ca 4 min. After 45 min the solvent was removed by evaporation and the resulting product was dissolved in EtOAc/MeOH/H<sub>2</sub>O and the acidic solution was treated with 2 M NaOH(aq) to pH=8-9. The organic layer was separated and dried(Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent afforded 425 mg (100%) of the title compound as a white solid

MS m/z 520 (M<sup>+</sup> + 1)

(iii) BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-(R,S)Itp(Ts)

H-(R)Cgl-Aze-(R,S)Itp(Ts) (400 mg, 0.77 mmol), Benzyl-2-(para-nitrobenzenesulfonyloxy)acetate (See Preparation of starting materials) (325 mg, 0.92 mmol) and K<sub>2</sub>CO<sub>3</sub> (235 mg, 1.7 mmol) was stirred in CH<sub>3</sub>CN (5 ml) at 45°C. After a few hours the conversion was only 25% and therefore the temperature was increased to 60°C and an additional amount of Benzyl-2-(para-nitrobenzenesulfonyloxy)acetate was added.

The reaction was stirred for 48 h,  
(startingm.:product/25:63), and then worked up. The  
solvent was evaporated and EtOAc/H<sub>2</sub>O was added to the  
residue. The phases were separated and the water-  
5 phase was washed twice with EtOAc and then the  
combined organic phase was washed with K<sub>2</sub>CO<sub>3</sub>(sat), 2 M  
KHSO<sub>4</sub>, H<sub>2</sub>O and dried Na<sub>2</sub>SO<sub>4</sub>). This afforded, after  
back-extraction of the acidic KHSO<sub>4</sub>, some 340 mg which  
was purified by RPLC. This gave 34 mg (7%) of the  
10 title compound

MS m/z 668 (M<sup>+</sup> + 1).

(iv) HOOC-CH<sub>2</sub>-(R)Cgl-Aze-(R,S)Itp x 2 HCl  
15

BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-(R,S)Itp(Ts) (34 mg, 0.05 mmol)  
was dissolved in THF (5 ml) and NH<sub>3</sub>(g) was distilled  
(40 ml) into the reaction flask with a dry-ice  
cooler. Na(s) was added and a deep blue color  
20 appeared. The reaction was stirred for 5 min before  
it was quenched with HOAc (50 µl). The dry-ice cooler  
was removed and the NH<sub>3</sub>(l) was allowed to evaporate.  
To the residue H<sub>2</sub>O and HOAc was added to pH=7. Freeze-  
drying and preparative RPLC gave several fractions  
25 which were analyzed with FAB-MS. Two fractions  
contained the desired compound, 3 mg (10%) after  
freeze-drying with 2.2 eq of 1 M HCl:

MS m/z 424 (M<sup>+</sup> +1).

30

Example 71

HOOC-CH<sub>2</sub>-(R)Cha-Aze-(R,S)Itp

35 (i) Boc-(R)Cha-Aze-(R,S)Itp(Ts)

B c-(R)Cha-Az -OH (See Preparation starting

materials) (169 mg, 0.5 mmol), H-(R,S)Itp(Ts) (See Preparation of starting materials) (155 mg, 0.52 mmol), DMAP (122 mg, 1 mmol) was dissolved in CH<sub>3</sub>CN (2.5 ml) and cooled to 5°C. EDC x HCl (115 mg, 0.6 mmol) was added and the resulting mixture was stirred at room temperature over night. Extra (0.5 eq) H-(R,S)Itp(Ts) and EDC was added after stirring over night. The reaction mixture was stirred an additional night and worked up as described in the Boc-(R)Cgl-Aze-(R,S)Itp(Ts) (See Example 70) case above. This gave 260 mg of crude product. Purification by RPLC gave 180 mg (57%) of the title compound

MS m/z 634 (M<sup>+</sup> +1).

(ii) H-(R)Cha-Aze-(R,S)Itp(Ts)

Boc-(R)Cha-Aze-(R,S)Itp(Ts) (180 mg, 0.28 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 ml) and HCl(g) was bubbled through the solution for ca 4 min. After 45 min the solvent was removed by evaporation and the resulting product was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with 2 M NaOH to pH=8-9. The phases were separated and the organic phase was dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated to yield 163 mg (ca 100%):

MS m/z 534 (M<sup>+</sup> +1).

(iii) BnOOC-CH<sub>2</sub>-(R)Cha-Aze-(R,S)Itp(Ts)

H-(R)Cha-Aze-(R,S)Itp(Ts) (80 mg, 0.15 mmol), K<sub>2</sub>CO<sub>3</sub> (45 mg, 0.33 mmol) and Br-CH<sub>2</sub>COOBn (39 mg, 0.17 mmol) was stirred in CH<sub>3</sub>CN (1.5 ml) at 60°C for 2.5 h. The solvent was evaporated and the residue was dissolved in EtOAc/H<sub>2</sub>O. The phases were separated and organic phase was washed with 10% citric acid and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent gave a 171 mg of



crude product, which was purified by RPLC yielding 53 mg (52%) of the title compound.

MS m/z 681 ( $M^+ + 1$ ).

5

(iv)  $\text{HOOC-CH}_2\text{-(R)Cha-Aze-(R,S)Itp}$

BnOOC- $\text{CH}_2\text{-(R)Cha-Aze-(R,S)Itp(Ts)}$  (50 mg, 0.07 mmol) was treated as described for BnOOC- $\text{CH}_2\text{-(R)Cgl-Aze-(R,S)Itp(Ts)}$  (See Example 70 (iv)). This gave a product mixture which was purified on a RPLC yielding 12 mg of a 1:1 mixture of the title compound together with a reduced form which appear at mass 439 (m/z).

15 MS m/z 438 ( $M^+ + 1$ )

#### Example 72

H-(R)Cha-Pic-(R,S)Itp x 2 HCl

20

(i) Boc-(R)Cha-Pic-(R,S)Itp(Ts)

At roomtemperature 2.1 g (5.5 mmol) Boc-(R)Cha-Pic-OH (See Preparation of starting materials), 1.0 g (8.2 mmol) DMAP and 1.7 g (5.8 mmol) H-(R,S)Itp(Ts) (See Preparation of starting materials) was dissolved in 40 mL acetonitrile. After a few minutes of stirring 1.1 g (5.8 mmol) EDC was added and the stirring was continued for 60 hours. The solvent was removed in vacuo and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$ , washed with water, 0.3M  $\text{KHSO}_4$  and  $\text{KHCO}_3$  (aq) and dried( $\text{Na}_2\text{SO}_4$ ). Evaporation of the solvent and filtration through Silica gel gave 2.43 g (67%) of the product.

35

MS m/z 661 ( $M^+ + 1$ )

## (ii) Boc-(R)Cha-Pic-(R,S)Itp

2.4 g (3.6 mmol) Boc-(R)Cha-Pic-(R,S)Itp(Ts) was dissolved in 15 mL THF and  $\text{NH}_3$  (g) was condensed into the flask followed by addition of Na. The reaction was quenched after 5 min with acetic acid and the  $\text{NH}_3$  and the THF was evaporated. The residue was freeze-dried from water and purified by RPLC ( $\text{CH}_3\text{CN}/0.1\text{M NH}_4\text{OAc}$ , 6/4) to give 0.93 g (51%) of the desired product.

MS m/z 507 ( $\text{M}^+ + 1$ )

## (iii) H-(R)Cha-Pic-(R,S)Itp x 2 HCl

At roomtemperature 50 mg (0.099 mmol) Boc-(R)Cha-Pic-(R,S)Itp was dissolved in ethylacetate saturated with HCl (g). After stirring 2 h the solvent was removed in vacuo. The residue was freeze-dried from water three times to give 35 mg (74%) of the desired product.

MS m/z 407 ( $\text{M}^+ + 1$ )

25 Example 73

$\text{HOOC-CH}_2\text{-(R)Cha-Pic-(R,S)Itp x 2 HCl}$

## (i) Boc-(R)Cha-Pic-(R,S)Itp(Z)

At roomtemperature 0.84 g (1.66 mmol) Boc-(R)Cha-Pic-(R,S)Itp (See Example 72) was dissolved in 10 mL  $\text{CH}_2\text{Cl}_2$  and 10 mL 0.5M NaOH. 0.29 mL (1.82 mmol) Z-Cl was added dropwise. After stirring for 3 h the phases was separated and the organic phase was washed with water and dried over  $\text{Na}_2\text{SO}_4$ . Evaporation and flash chromatography (ethylacetate/heptane 9/1) gave 0.5 g

(47%) of the desired product.

MS m/z 641 ( $M^+ + 1$ )

5 (ii) H-(R)Cha-Pic-(R,S)Itp(Z)

At roomtemperature 0.5 g (0.78 mmol) Boc-(R)Cha-Pic-(R,S)Itp(Z) was dissolved in ethylacetate saturated with HCl. Water was added and the mixture was made  
10 basic with  $K_2CO_3$ . The phases were separated. The waterphase was extracted with  $CH_2Cl_2$  and the organic phase was washed with water. The combined organic phase was then dried ( $Na_2SO_4$ ). Evaporation of the solvent gave 0.3 g (71%) of the desired product.

15

MS m/z 541 ( $M^+ + 1$ )

(iii)  $BnOOC-CH_2-(R)Cha-Pic-(R,S)Itp(Z)$

20 0.29 g (0.5 mmol) H-(R)Cha-Pic-(R,S)Itp(Z), 0.15 g (1 mmol)  $K_2CO_3$  was taken up in 25 mL acetonitrile. 154 mg (0.6 mmol) benzylbromoacetate was added and the mixture was stirred at 50°C for 4 h. evaporation and purification by RPLC  
25 (acetonitrile:0.1M  $NH_4OAc$  70:30) gave about 200 mg of the desired product.

(iv)  $HOOC-CH_2-(R)Cha-Pic-(R,S)Itp \times 2 HCl$

30 200 mg  $BnOOC-CH_2-(R)Cha-Pic-(R,S)Itp(Z)$  was dissolved in ethanol. A small spoon of 10% Pd on charcoal was added and the mixture was hydrogenated for 4 h.  
Filtration through hyflo, evaporation of the solvent followed by freeze-drying from water gave 53 mg of the  
35 desired product.

$^1H$  NMR (300.13 MHz,  $D_2O$ );  $\delta$  1.0-2.35 (overlapping m,

22H), 3.28-3.51 (m, 5H), 3.51-3.64 (m, 1H), 3.75-4.03 (m, 3H), 5.03-5.14 (s broad, 1H). The signal of one of the protons is partially obscured by the H-O-D-signal.

5

MS m/z 465 ( $M^+ + 1$ )

Example 74

10

H-(R)Cgl-Pro-(R,S)Hig x 2 HCl

(i) Boc-(R)Cgl-Pro-(R,S)Hig(Z)

15

To a mixture of 1.0 g (2.95 mmole) Boc-(R)Cgl-Pro-OH (See Preparation of starting materials), 1.44 g (11.8 mmole) DMAP, 1.12 g (3.25 mmole) H-(R,S)Hig(Z) (See Preparation of starting materials) in 15 ml  $CH_2Cl_2$  was added 0.62 g (3.2 mmole) of EDC and the mixture was stirred at room temperature over night. The solvent was evaporated and the residue was dissolved in ethyl acetate. When the organic layer was washed twice with a 0.3 M  $KHSO_4$ -solution an oil separated from the organic layer. The ethyl acetate layer was dried ( $Na_2SO_4$ ) and filtered. The oil and the water layer was then extracted with  $CH_2Cl_2$ . The organic layer was dried ( $Na_2SO_4$ ), filtered and combined with the EtOAc phase from above. Evaporation and purification of the crude product on a chromatotron (Harrison research, model 7924T) using a 2mm silica plate with a stepwise gradient of  $CH_2Cl_2$ /MeOH (97/3, 95/5, 90/10) as eluent to yielded 1.1 g (59%) of the title compound.

20

25

30

35

(ii) H-(R)Cgl-Pro-(R,S)Hig x 2 HCl

81 mg (0.13 mmole) of Boc-(R)Cgl-Pro-(R,S)Hig(Z) was dissolved in 50 ml ethyl acetate saturated with HCl.

The mixture was allowed to stand for one hour, evaporated and the residue was dissolved in 10 ml ethanol. 40 mg Pd/C (5%), 1 ml water and 0.5 ml 1 M HCl-solution was added and the mixture was hydrogenated at atmospheric pressure over night. Filtration of the catalyst through cellite and evaporation of the solvent followed by freeze drying 3 times from water gave the title compound in 75% yield.

$^1\text{H-NMR}$  ( $\text{D}_2\text{O}$ , 300 MHz):  $\delta$  0.95-1.35 (m, 5H), 1.50-2.45 (m, 15H), 3.02 (bt, 1H), 3.1-3.8 (m, 7H), 4.13 (d, 1H), 4.38 (bd, 1H)

$^{13}\text{C-NMR}$  ( $\text{D}_2\text{O}$ , 75 MHz): carbonyl and guanidinecarbons:  $\delta$  154.8, 168.9, 174.4

MS  $m/z$  393 ( $\text{M}^++1$ )

#### Example 75

$\text{HOOC-CH}_2\text{-(R)Cgl-Pro-(R,S)Hig} \times 2 \text{ HCl}$

(i)  $\text{H-(R)Cgl-Pro-(R,S)Hig(Z)}$

1 g (1.6 mmole) Boc-(R)Cgl-Pro-(R,S)Hig(Z) (See Example 74 (i)) was dissolved in 100 ml ethyl acetate saturated with HCl, and the mixture was allowed to stand for one hour. The mixture was evaporated and the residue was dissolved in  $\text{CH}_2\text{Cl}_2$ . The organic layer was washed twice with 0.2 M NaOH-solution, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated to yield 0.825 g (98%) of title compound.

(ii)  $\text{BnOOC-CH}_2\text{-(R)Cgl-Pro-(R,S)Hig(Z)}$

0.442 g (0.839 mmole)  $\text{H-(R)Cgl-Pro-(R,S)Hig(Z)}$ , 0.256 g (1.85 mmol)  $\text{K}_2\text{CO}_3$  and 145  $\mu\text{l}$  (0.521 mmole) of

bensylbromoacetate was mixed in 12 ml THF. The mixture was stirred at 40°C for one hour and at room temperature over night. After evaporation of the solvent the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed once with water and once with brine. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated and the crude product was purified on a chromatotron (Harrison research, model 7924T) using a 2mm silica plate with a stepwise gradient of CH<sub>2</sub>Cl<sub>2</sub>/MeOH (97/3, 95/5, 90/10) as eluent to yield 0.165 g (29%) of the title compound.

(iii) HOOC-CH<sub>2</sub>-(R)Cgl-Pro-(R,S)Hig x 2 HCl

0.165 g (0.25 mmole) of BnOOC-CH<sub>2</sub>-(R)Cgl-Pro-(R,S)Hig(Z) was mixed with 0.050 g Pd/C (5%), 0.7 ml 1 M HCl-solution and 10 ml ethanol. The mixture was hydrogenated at atmospheric pressure for four hours. Filtration of the catalyst through cellite and evaporation of the solvent followed by freeze drying twice from water gave 0.1 g (75%) of the product.

<sup>1</sup>H-NMR (D<sub>2</sub>O, 300 MHz): δ 1.05-1.45 (m, 5H), 1.55-2.5 (m, 15H), 3.08 (bt, 1H), 3.2-4.05 (m, 9H), 4.30 (d, 1H), 4.44 (m, 1H)

<sup>13</sup>C-NMR (D<sub>2</sub>O, 75 MHz): carbonyl and guanidinecarbons: δ 154.9, 167.2, 169.4, 174.1

### Example 76

H-(R)Cha-Pro-(R,S)Hig x 2 HCl

(i) Boc-(R)Cha-Pro-(R,S)Hig(Z)

0.72 g (1.95 mmole) Boc-(R)Cha-Pro-OH (See Preparation of starting materials), 0.95 g (7.8

mmole) DMAP, 0.74 g (2.14 mmole) 82% pure H-(R,S)Hig(Z) (See Preparation of starting materials) in 10 ml CH<sub>2</sub>Cl<sub>2</sub> was added 0.486 g (2.54 mmole) of EDC and the mixture was stirred at room temperature for 3 days. The mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with water, twice with 0.3M KHSO<sub>4</sub>-solution and once with brine. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated and the crude product was purified by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH 95/5 as eluent to yield 0.450 g (33%) of the product.

(ii) H-(R)Cha-Pro-(R,S)Hig x 2 HCl

50 mg (0.078 mmole) of Boc-(R)Cha-Pro-(R,S)Hig(Z) was dissolved in 20 ml ethyl acetate saturated with HCl. The mixture was allowed to stand for one hour, evaporated and the residue was dissolved in 10 ml ethanol. 20 mg Pd/C (5%) and 0.3 ml 1 M HCl-solution was added and the mixture was hydrogenated at atmospheric pressure for two hours. Filtration of the catalyst through celite and evaporation of the solvent followed by freeze drying twice from water gave 28 mg (76%) of the title compound.

<sup>1</sup>H-NMR (D<sub>2</sub>O, 300 MHz): δ 0.9-1.6 (m, 6H), 1.6-2.5 (m, 16H), 3.09 (t, 1H), 3.31 (t, 1H), 3.37-3.74 (m, 4H), 3.81 (m, 1H), 4.35-4.47 (m, 2H)

<sup>13</sup>C-NMR (D<sub>2</sub>O, 75 MHz): carbonyl and guanidinecarbons: δ 154.9, 169.8, 174.5

Example 77

H-(R)Cgl-Aze-Rig x 2 HCl

(i) Boc-(R)Cgl-Aze-Rig(Z)

T a solution of 0.50 g (1.6 mmol) of H-Rig(Z) (See Preparation of starting materials) , 0.59 g (1.6 mmol) of Boc-(R)Cha-Aze-OH( See preparation of starting materials), 0.84 g (6.9 mmol) of  
5 dimethylaminopyridine in 30 ml of acetonitrile and 5 ml of dimethylformamide was added 0.33 g (1.7 mmol) of N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride. The reaction was allowed to stir for 3 days then evaporated and partitioned between aqueous  
10 potassium hydrogen sulfate and methylene chloride. The methylene chloride layer was washed with aqueous sodium bicarbonate and water, dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated. The crude material was suction filtered through a pad of silica gel with methylene  
15 chloride/methanol 9/1 to give 0.78 g (76%) of the desired compound after evaporation.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 0.8-1.9 (m, 27 H), 2.4-2.6 (m, 2 H), 2.78 (bt, 2 H), 3.15-3.4 (m, 2 H), 3.80  
20 (bt, 1 H), 4.0-4.4 (m, 4 H), 4.75 (bt, 1 H), 4.97 (bd, 1 H), 5.08 (s, 2 H), 7.1-7.4 (m, 7 H), 7.74 (b, 1 H).

(ii) H-(R)Cgl-Aze-Rig(Z) x 2 HCl  
25

A flask containing Boc-(R)Cgl-Aze-Rig(Z), 0.76 g (1.2 mmol), in 50 ml of ethyl acetate was cooled in an ice bath. Dry HCl was bubbled through for 5 min and the solution was evaporated to give 0.74 g (100%) of the  
30 dihydrochloride as a white powder.

<sup>1</sup>H-NMR (300 MHz, MeOD): δ 1.1-2.0 (m, 18 H), 2.23 (m, 1 H), 2.68 (m, 1 H), 3.15-3.45 (m, 4 H), 3.72 (bd, 1 H), 3.9-4.0 (bd, 2 H), 4.27 (m, 1 H), 4.39 (m, 1 H),  
35 4.78 (m, 1 H), 5.30 (s, 2 H), 7.3-7.5 (m, 5 H).

(iii) H-(R)Cgl-Az -Rig x 2 HCl



A flask containing a solution of 20 mg of H-(R)Cgl-Aze-Rig(Z) and a small amount of 5% Pd/C was hydrogenated at atmospheric pressure for 1 h. The mixture was filtered through celite and evaporated.  
5 The residue was lyophilized with a few drops of conc. HCl added to give the product. Yield: 8 mg (52%).

<sup>1</sup>H-NMR (300 MHz, D<sub>2</sub>O): δ 1.1-2.0 (m, 18 H), 2.37 (m, 1 H), 2.75 (m, 1 H), 3.08 (bt, 2 H), 3.39 (bt, 2 H),  
10 3.8-4.0 (m, 3 H), 4.35-4.5 (m, 2 H), 4.90 (m, 1 H).

<sup>13</sup>C-NMR (75.5 MHz, D<sub>2</sub>O): guanidine and carbonyl carbons: δ 172.2, 169.4, 156.4.

15 Example 78

HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Rig x 2 HCl

20 (i) BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Rig(Z)

A mixture of 0.20 g (0.33 mmol) of H-(R)Cgl-Aze-Rig(Z) (See Example 77), 0.13 g of potassium carbonate, 80 mg of sodium iodide, 10 ml of tetrahydro-furane and 10 ml of acetonitrile was  
25 heated at 60°C for 10 h. The solvents were evaporated and the crude material was flash chromatographed on silica gel using methylene chloride/methanol 92/8 as eluent. Yield: 0.13 g (58%).

30 <sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ 0.9-2.1 (m, 18 H), 2.45 (m, 1 H), 2.61 (m, 1 H), 2.81 (m, 2 H), 2.88 (d, 1 H), 3.2-3.5 (m, 4 H), 3.94 (m, 1 H), 4.0-4.25 (m, 3 H), 4.85 (m, 1 H), 5.12 (s, 2 H), 5.14 (s, 2 H), 6.9-7.2 (b. 2 H), 7.2-7.5 (m, 10 H), 7.95 (m, 1 H).

35

(ii) HOOC-CH<sub>2</sub>-(R)Cgl-Aze-Rig x 2 HCl

A mixture of 0.12 g (0.18 mmol) of  $\text{BnOOC-CH}_2\text{-(R)Cgl-Az-Rig(Z)}$ , 5 ml of ethan 1, 3 drops of conc. HCl and a small amount of 5% Pd/C was hydrogenated at atmospheric pressure for 1 h. The mixture was  
5 filtered through celite and evaporated. The residue was lyophilized in water to give 91 mg (98%) of the product.

$^1\text{H-NMR}$  (500 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  1.1-1.9 (m, 17 H), 2.00 (m, 1 H), 2.29 (m, 1 H), 2.70 (m, 1 H), 3.10 (m, 2 H), 3.34 (t, 2 H), 3.83 (bd, 2 H), 3.89 (dd, 2 H), 4.00 (d, 1 H), 4.35 (m, 2 H), 4.87 (m, 1 H).  
10

$^{13}\text{C NMR}$  (125.8 MHz,  $\text{D}_2\text{O}$ ): guanidine and carbonyl carbons:  $\delta$  171.8, 169.6, 167.7, 156.3.  
15

#### Example 79

$\text{HOOC-CH}_2\text{-(R)Cha-Pro-Rig} \times 2 \text{ HCl}$   
20

(i)  $\text{Boc-(R)Cha-Pro-Rig(Z)}$

To a solution of 0.25 g (0.82 mmol) of 4-aminoethyl-1-benzyloxycarbonylamidino piperidine( $\text{H-Rig(Z)}$ ), (See  
25 preparation of starting materials), 0.32 g (0.82 mmol) of  $\text{Boc-(R)Cha-Pro-OH}$  (See Preparation of starting materials), 0.40 g (3.3 mmol) of dimethylaminopyridine in 10 ml of acetonitrile and 2 ml of dimethylformamide was added 0.165 g (0.86 mmol)  
30 of  $\text{N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide}$  hydrochloride. The reaction was allowed to stir for 3 days then evaporated and partitioned between aqueous potassium hydrogen sulfate and methylene chloride. The methylene chloride layer was washed with aqueous  
35 sodium bicarbonate and water, dried ( $\text{Na}_2\text{SO}_4$ ) and evap rat d. The NMR spectrum f the crude product was satisfactory and th product which contained som

dimethylformamide was used in the next step without further purification.

5  $^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.8-2.2 (m, 32 H; thereof 1.41 (s, 9 H)), 2.34 (m, 1 H), 2.77 (bt, 2 H), 3.10 (m, 1 H), 3.29 (m, 1 H), 3.40 (m, 1 H), 3.83 (m, 1 H), 4.17 (m, 2 H), 4.30 (m, 1 H), 4.54 (m, 1 H), 5.07 (m, 1 H), 5.08 (s, 2 H), 7.03 (m, 1 H), 7.05-7.4 (m, 7 H).

10

(ii) H-(R)Cha-Pro-Rig(Z)

15 A flask containing the crude product of Boc-(R)Cha-Pro-Rig(Z) in 100 ml of ethyl acetate was cooled in an ice bath. Dry HCl was bubbled through for 5 min and the solution was evaporated to get rid of the excess of HCl. The product was dissolved in water and the extracted twice with ethyl acetate to remove the dimethylformamide from the previous step. The  
20 aqueous phase was made alkaline with  $\text{NaHCO}_3$  (aq) and extracted twice with methylene chloride. The combined organic phase was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. Yield: 0.37 g (81%) over two steps.

25  $^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ )  $\delta$  0.8-2.4 (m, 24 H), 2.82 (bt, 2 H), 3.26 (m, 2 H), 3.42 (bq, 1 H), 3.70 (m, 2 H), 4.19 (m, 2 H), 4.49 (bd, 1 H), 5.11 (s, 2 H), 6.9-7.5 (m, 8 H).

30 (iii)  $\text{BnOOC-CH}_2\text{-(R)Cha-Pro-Rig(Z)}$

A mixture of 0.18 g (0.32 mmol) of H-(R)Cha-Pro-Rig(Z), an excess of potassium carbonate and 10 ml of acetonitrile was heated at  $60^\circ\text{C}$  for 2 h. The solvents  
35 wer evaporated and the crude material was flash chromatographed n silica gel using methylene chlorid /methanol 95/5 as eluent. Yield: 0.20 g

(88%).

<sup>1</sup>H-NMR (300 MHz, CDCl<sub>3</sub>) δ 0.8-2.1 (m, 23 H), 2.37 (m, 1 H), 3.1-3.5 (m, 7 H), 4.0-4.2 (m, 2 H), 4.54 (m, 1 H), 5.1 (m, 4 H), 6.9-7.5 (m, 13 H).

(iv) HOOC-CH<sub>2</sub>-(R)Cha-Pro-Rig x 2 HCl

A mixture of 0.15 g (0.21 mmol) of BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Rig(Z), 10 ml of ethanol, 4 drops of conc. HCl and a small amount of 5% Pd/C was hydrogenated at atmospheric pressure for 1 h. The mixture was filtered through celite and evaporated. The residue was lyophilized in water to give 95 mg (64%) of the product.

<sup>1</sup>H-NMR (500 MHz, MeOD) δ 0.85- 2.1 (m, 23 H), 2.30 (m, 1 H), 3.10 (m, 2 H), 3.25 (m, 1 H), 3.35 (m, 1 H), 3.54 (m, 1 H), 3.85-4.0 (m, 3 H), 4.03 (d, 1 H), 4.41 (m, 1 H), 4.50 (m, 1 H).

<sup>13</sup>C-NMR (125.8 MHz, D<sub>2</sub>O): guanidine and carbonyl carbons: δ 174.0, 168.9, 168.1, 157.5.

#### Example 80

HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Aze-Rig x 2 HCl

(i) Boc-(R)Cha-Aze-Rig(Z)

To a solution of 0.25 g (0.82 mmol) of 4-aminoethyl-1-benzyloxy-carbonylamidino piperidine (H-Rig(Z)), (See preparation of starting materials), 0.31 g (0.86 mmol) of Boc-(R)Cha-Aze-OH (See preparation of starting materials), 0.40 g (3.3 mmol) of dimethylaminopyridine in 10 ml of acetonitrile and 2 ml of dimethylformamide was added 0.17 g (0.86 mmol)

of N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride. The reaction was allowed to stir for 3 days then evaporated and partitioned between aqueous potassium hydrogen sulfate and methylene chloride. The methylene chloride layer was washed with aqueous sodium bicarbonate and water, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. The crude product which contained some dimethylformamide was used in the next step without further purification.

10

$^1\text{H-NMR}$  (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.85 (m, 1 H), 0.97 (m, 1 H), 1.1-1.75 (m, 26 H; thereof 1.41 (s, 9 H)), 1.82 (bd, 1 H), 2.53 (m, 2 H), 2.77 (bt, 2 H), 3.25 (m, 2 H), 4.03 (q, 1 H), 4.08 (m, 1 H), 4.18 (m, 2 H), 4.29 (m, 1 H), 4.78 (m, 1 H), 4.97 (m, 1 H), 5.09 (s, 2 H), 7.1-7.4 (m, 7 H), 7.65 (m, 1 H).

15

(ii) H-(R)Cha-Aze-Rig(Z)

A flask containing the crude product of Boc-(R)Cha-Aze-Rig(Z) in 100 ml of ethyl acetate was cooled in an ice bath. Dry HCl was bubbled through for 5 min and the solution was evaporated to get rid of the excess of HCl. The product was dissolved in water and the extracted twice with ethyl acetate to remove the dimethylformamide from the previous step. The aqueous phase was made alkaline with  $\text{NaHCO}_3$  (aq) and extracted twice with methylene chloride. The combined organic phase was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and evaporated. Yield: 0.31 g (70%) over two steps.

25

30

$^1\text{H-NMR}$  (300 MHz,  $\text{CDCl}_3$ )  $\delta$  0.8-1.9 (m, 20 H), 2.48 (m, 1 H), 2.73 (m, 1 H), 2.85 (bt, 2 H), 3.25 (m, 1 H), 3.35 (m, 2 H), 4.05 (q, 1 H), 4.1-4.25 (m, 3 H), 4.86 (m, 1 H), 5.12 (s, 2 H), 6.9-7.2 (m, 2 H), 7.2-7.45 (m, 5 H), 7.93 (m, 1 H).

35

(iii)  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Rig(Z)}$ 

A solution of 0.31 g (0.57 mmol) of H-(R)Cha-Aze-Rig(Z) and 93 mg (0.57 mmol) of benzyl acrylate in 5 ml of ethanol was allowed to stand at room temperature for one week. It was evaporated and flash chromatographed on silica gel using methylene chloride/methanol 94/6 as eluent. Yield: 0.20 g (49%).

10

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  0.8-1.0 (m, 2 H), 1.1-1.8 (m, 18 H), 2.48 (m, 1 H), 2.54 (bt, 2 H), 2.68 (m, 2 H), 2.81 (bt, 2 H), 2.87 (m, 1 H), 3.20 (m, 1 H), 3.25 (m, 1 H), 3.31 (m, 1 H), 4.04 (q, 1 H), 4.1-4.2 (m, 3 H), 4.84 (dd, 1 H), 5.05-5.15 (m, 4 H), 7.0-7.5 (m, 12 H), 8.03 (m, 1 H).

15

(iv)  $\text{HOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Rig} \times 2 \text{ HCl}$ 

The title compound was made and purified in the same way as described in Example 80 from 0.20 g (0.28 mmol) of  $\text{BnOOC-CH}_2\text{-CH}_2\text{-(R)Cha-Aze-Rig(Z)}$ . Yield: 30 mg (19%) of the dihydrochloride salt.

20

$^1\text{H}$ -NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  1.0-1.9 (m, 20 H), 2.33 (m, 1 H), 2.70 (m, 1 H), 2.83 (m, 2 H), 3.10 (m, 2 H), 3.3-3.4 (m, 4 H), 3.85 (bd, 2 H), 3.92 (m, rotamer), 4.14 (t, 1 H), 4.17 (m, rotamer), 4.31 (m, 1 H), 4.46 (m, 1 H), 4.89 (m, 1 H), 5.18 (m, rotamer).

25

30

$^{13}\text{C}$  NMR (125.8 MHz,  $\text{D}_2\text{O}$ ) guanidine and carbonyl carbons:  $\delta$  175.4, 171.8, 168.8, 156.3.

Example 81

35

$\text{HOOC-CH}_2\text{-(R)Cha-Pro-(S)Itp} \times 2 \text{ HCl}$

## (i) Boc-(R)Cha-Pr -(S)Itp(Ts)

At roomtemperature 0.87 g (2.36 mmol) Boc-(R)Cha-Pro-OH (See preparation of startingmaterials), 0.78 g (4.72 mmol) DMAP and 0.70 g (2.36 mmol) H-(S)Itp(Ts) (See preparation of starting materials) was dissolved in 12 mL acetonitrile. After stirring 20 minutes 0.59 g (3.07 mmol) EDC was added. After 18 hours the solvent was removed in vacuo and the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, washed with water, citric acid (10%), KHCO<sub>3</sub> (aq), water and dried with Na<sub>2</sub>SO<sub>4</sub>. Evaporation gave 1.74 g (> 100% yiled (purity of about 60%)) of the desired product. Which was used in the next step without further purification.

FAB-MS: m/z = 647 (M<sup>+</sup> + 1)

## (ii) H-(R)Cha-Pro-(S)Itp(Ts)

The Boc-protecting group was removed in the same way as described for Boc-(R)Cha-Pic-(R,S)Itp(Z) (See Example 72 (ii)) to give 0.75 g (81%) of the title compound.

FAB-MS: m/z = 547 (M<sup>+</sup> + 1)

(iii) BnOOC-CH<sub>2</sub>-(R)Cha-Pro-(S)Itp(Ts)

0.75 g (1.37 mmol) H-(R)Cha-Pro-(S)Itp(Ts), 0.38 g (2.74 mmol) K<sub>2</sub>CO<sub>3</sub> was taken up in 15 mL acetonitrile. 0.39 g (1.65 mmol) benzylbromoacetate was added and the mixture was stirred at 50°C for 2 h. Evaporation of the solvent followed by flash chromatography using ethylacetate/methanol 95/5 as eluent gave about 530 mg of the desired product.

FAB-MS: m/z = 695 (M<sup>+</sup> + 1)

(iv)  $\text{HOOC-CH}_2\text{-(R)Cha-Pro-(S)Itp} \times 2 \text{ HCl}$

0.53 g (0.76 mmol)  $\text{BnOOC-CH}_2\text{-(R)Cha-Pro-(S)Itp(Ts)}$  was dissolved in 15 mL THF.  $\text{NH}_3$  (g) was condensed into the flask and Na (m) was added. The reaction was quenched after 30 min with acetic acid and the  $\text{NH}_3$  and the THF was evaporated. The residue was freeze dried from water and the crude product was purified by RPLC (acetonitrile/0.1M HOAc 15/85) gave 0.25 g (61%) of the desired product after freeze-drying from aqueous HCl.

$^1\text{H-NMR}$  (500.13 MHz,  $\text{D}_2\text{O}$ );  $\delta$  0.9-2.09 (overlapping m, 20H), 2.22-2.35 (m, 1H), 3.2-3.36 (m, 4H), 3.44-3.62 (overlapping m, 2H), 3.7-3.8 (m, 1H), 3.87-3.99 (m, 2H), 4.33-4.48 (overlapping m, 2H).

$^{13}\text{C-NMR}$  (500.13 MHz,  $\text{D}_2\text{O}$ ); carbonyl- and guanidinecarbons:  $\delta$  154.3, 168.1, 169.0 and 174.2

#### Example 82

$\text{H-(R)Cha-Pro-(R,S)Nig} \times 2 \text{ HCl}$

(i)  $\text{Boc-(R)Cha-Pro-(R,S)Nig(Z)}$

174 mg (0.471 mmole)  $\text{Boc-(R)Cha-Pro-OH}$  (See preparation of starting materials), 229 mg (1.87 mmole) DMAP, 130 mg (0.471 mmole)  $\text{H-(R,S)Nig(Z)}$  (See Preparation of starting materials) was mixed in 2 ml  $\text{CH}_2\text{Cl}_2$  and 117 mg (0.61 mmole) of EDC was added and the mixture was stirred for four days. The mixture was diluted with  $\text{CH}_2\text{Cl}_2$  and washed with water, twice with 0.3 M  $\text{KHSO}_4$ -solution and once with brine. The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and vaporated and the crude product was purified twice by flash chromatography using  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  95/5 as



eluent the first time and  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  97/3 as eluent the second time to yield 0.104 g (35%) of the title compound.

5 MS m/z 627 ( $\text{M}^+ + 1$ )

(ii) H-(R)Cha-Pro-(R,S)Nig x 2 HCl

10 10 mg (0.016 mmole) of Boc-(R)Cha-Pro-(R,S)Nig(Z) was dissolved in 15 ml ethyl acetate saturated with HCl. The mixture was allowed to stand for half an hour. The mixture was evaporated and the residue was dissolved in 6 ml ethanol and 8 mg 5% Pd/C (5%) and 0.1 ml 1 M HCl-solution was added and the mixture was  
15 hydrogenated at atmospheric pressure for one and a half hour. After filtration through hyflo and evaporation of the solvent gave 4 mg of the title compound the product

20  $^1\text{H}$ -NMR (300 MHz,  $\text{D}_2\text{O}$ ):  $\delta$  0.9-1.58 (m, 6H), 1.58-2.45 (m, 13H), 2.65 (m, 1H), 3.19 (m, 1H), 3.34 (d, 2H), 3.4-3.73 (m, 4H), 3.82 (m, 1H), 4.34-4.49 (m, 2H).

$^{13}\text{C}$ -NMR (75 MHz,  $\text{D}_2\text{O}$ ): carbonyl and guanidinecarbons:  
25  $\delta$  155.1, 169.9 and 174.8.

### Example 83

H-(R)Pro-Phe-Pab x 2 HCl

30

(i) Boc-(R)Pro-Phe-Pab(Z)

To a mixture of 1.2 g (3.31 mmol) Boc-(R)Pro-Phe-OH (See preparation of starting materials) and 1.70 g (13.91 mmol) DMAP in 40 ml  $\text{CH}_3\text{CN}$  at room temperature  
35 was added 0.98 g (3.35 mmol) H-Pab(Z) (See preparation of starting materials) dissolved in 1 ml

DMF. After stirring for 2 h the reaction mixture was cooled to - 18°C and 0.66 g (3.48 mmol) EDC was added portion wise and the reaction was left at room temperature over night. The solvent was evaporated and the residue was dissolved in 100 ml EtOAc and washed with 1 x 30 ml water, 3 x 30 ml 0.3 M KHSO<sub>4</sub>, 1 x 30 ml Na<sub>2</sub>CO<sub>3</sub>, 1 x 30 ml water and dried. Evaporation of the solvent followed by flash chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (95/5) as eluent gave 0.691 g (38%) of the title compound.

(ii) H-(R)Pro-Phe-Pab(Z)

0.673 g Boc-(R)Pro-Phe-Pab(Z) was dissolved in 30 ml EtOAc and the solution was saturated with HCl(g) for a few minutes (a white solid precipitated out from the solution). The solvent and excess HCl was evaporated and 60 ml EtOAc was added to the residue and the organic phase was washed with 2 x 20 ml 2 M NaOH. The washing water was extracted with 1 x 25 ml EtOAc which was combined with the other EtOAc-phase and the combined organic phase was washed with water, dried and evaporated to give 560 mg (98%) of the desired product.

<sup>1</sup>H-NMR (500 MHz, CDCl<sub>3</sub>): δ 1.5-1.74 (m, 3H), 1.98-2.05 (m, 1H), 2.78-2.85 (m, 1H), 2.90-2.96 (m, 1H), 3.0-3.2 (ABX-system centered at 3.1, 2H), 3.62 (dd, 1H), 4.3-4.45 (ABX-system centered at 4.37, 2H), 4.58 (q, 1H), 5.22 (s, 2H), 6.96 (bt, 1H), 7.1-7.4 (m, 10H), 7.46 (d, 2H), 7.76 (d, 2H), 8.12 (d, 1H).

(iii) H-(R)Pro-Phe-Pab x 2 HCl

200 mg H-(R)Pro-Phe-Pab(Z) was dissolved in 10 ml 95 % EtOH and 2 ml of water and the mixture was hydrogenated over 5 % Pd/C at atmospheric pressure

f r 5 h. Filtration of the catalyst and addition of 1 ml 1 M HCl followed by evaporation and freeze drying from water gave the title compound in 88 % yield.

5 <sup>1</sup>H-NMR (500 MHz, CD<sub>3</sub>OD): δ 1.51-1.59 (m, 1H), 1.69-1.80 (m, 1H), 1.87-1.97 (m, 1H), 2.19-2.29 (m, 1H), 2.90 (dd, 1H), 3.20-3.33 (m, 3H, partially hidden by the solvent peak), 4.27 (m, 1H), 4.43-4.54 (AB-system centered at 4.48, 2H), 4.75-4.81 (m, 1H), 4.87 (s, 10 2H), 7.2-7.3 (m, 5H), 7.45 (d, 2H), 7.75 (d, 2H).

<sup>13</sup>C-NMR (125 MHz, D<sub>2</sub>O): amidine and carbonyl carbons: δ 166.7, 170.1 and 173.4.

15 Example 84

HOOC-CH<sub>2</sub>-(R)Pro-Phe-Pab x 2 HCl

(i) BnOOC-CH<sub>2</sub>-(R)Pro-Phe-Pab(Z)

20

To a slurry of 244 mg (0.463 mmol) H-(R)Pro-Phe-Pab(Z) (See Example 83) and 159.9 mg (1.157 mmol) K<sub>2</sub>CO<sub>3</sub> in 8 ml DMF/CH<sub>3</sub>CN (5/3) was added 127.2 mg (0.555 mmol) benzylbromo acetate dissolved in 2 ml 25 DMF and the mixture was stirred at 60°C for 1.5 h and at room temperature over night. The solvent was evaporated and the residue was dissolved in 50 ml EtOAc, washed with 2 x 20 ml water and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent followed by flash 30 chromatography using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (9/1) as eluent gave 176 mg (56 %) of the title compound as a white solid.

<sup>1</sup>H-NMR (300 MHz), CDCl<sub>3</sub>): δ 1.45-1.80 (m, 3H), 2.06 (m, 1H), 2.54 (m, 1H), 2.92-3.28 (m, 6H), 4.3-4.5 (ABX-system centered at δ = 4.4, 2H), 4.60 (dd, 1H), 5.10 (apparent s, 2H), 5.2 (apparent s, 2H), 7.1-7.4 (m, 15H), 7.43 (d, 2H), 7.75 (d, 2H), 7.932 (d, 1H). 35

(ii)  $\text{HOOC-CH}_2\text{-(R)Pro-Phe-Pab} \times 2\text{HCl}$

170 mg (0.252 mmol) of  $\text{BnOOC-CH}_2\text{-(R)Pro-Phe-Pab(Z)}$  was dissolved in 12 ml EtOH/water (5/1) and hydrogenerated over 5 % Pd/C at atmospheric for 4.5 h. The catalyst was filtered off, the solvent evaporated and the residue freeze dried from  $\text{HCl(aq)}$  to give the title compound.

$^1\text{H-NMR}$  (500 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  1.62 (m, 1H), 1.82 (m, 1H), 2.08 (m, 1H), 2.38 (m, 1H), 2.90 (dd, 1H), 3.25-3.35 (m, 2H), partially hidden by the solvent peak), 3.80 (m, 1H), 4.08-4.19 (AB-system centered at  $\delta=4.19$ , 2H), 4.39 (m, 1H), 4.45-4.58 (AB-system centered at  $\delta=4.50$ , 2H), 4.80 (m, 1H), 7.20-7.35 (m, 5H), 7.45 (d, 2H), 7.75 (D, 2H).

$^{13}\text{C-NMR}$  (125 MHz,  $\text{D}_2\text{O}$ ): amidine and carbonyl carbons:  $\delta$  166.8, 169.1, 169.5 and 173.2.

#### Example 85

#### **H-(R)Phe-Phe-Pab**

(i)  $\text{Boc-(R)Phe-Phe-Pab(Z)}$

$\text{Boc-(R)Phe-Phe-OH}$  (16.4 mmol) (see preparation of starting materials),  $\text{Pab(Z)-HCl}$  (18.0 mmol) and 4-dimethylaminopyridine (24.6 mmol) were dissolved in 50 mL of acetonitrile. The solution was cooled to ice-water temperature and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (21.3 mmol) was added. The cooling bath was removed and the reaction mixture was stirred over night. The solvent was then evaporated under reduced pressure, the residue dissolved in 50 mL of ethyl acetate and the resulting solution extracted with 50 mL of water.  $\text{Boc-(R)Phe-}$

Phe-Pab(Z) precipitating from the two-phase mixture was filtered and washed with water yielding 8.7 g (78%) after drying under vacuum at 45°C for 24 h. <sup>1</sup>H NMR (200 MHz, d-CHCl<sub>3</sub> and d<sub>4</sub>-CH<sub>3</sub>OH); δ 8.35-7.00 (m, 19H), 4.63 (t, 1H), 4.3-4.1 (m, 1H), 3.40-2.70 (m, 6H), 1.30 (s, 9H).

(ii) H-(R)Phe-Phe-Pab(Z)

10 Boc-(R)Phe-Phe-Pab(Z) (10.3 mmol) was slurried in 70 mL of ethylacetate and 31 mL of 3.3 M ethylacetate/HCl was added. The slurry was stirred for 4 h after which the hydrochloride salt of H-(R)Phe-Phe-Pab(Z) was filtered off and washed with  
15 several portions of ethylacetate. The salt was dissolved in a mixture of 50 mL of methylenechloride, 50 mL of 1 M potassiumcarbonate and ca 5 mL of ethanol. The organic layer was collected and the solvent was removed under reduced pressure yielding  
20 5.0 g of H-(R)Phe-Phe-Pab(Z) (84%). <sup>1</sup>H NMR (200 MHz, d<sub>6</sub>-DMSO); δ 9.1 (s, 2H), 8.59 (m, 1H), 8.1 (m, 1H), 7.90 (d, 2H), 7.4-7.0 (m, 17H), 5.09 (s, 2H), 4.58 (m, 1H), 4.31 (m, 2H), 3.1-2.7 (m, 4H).

25 (iii) H-(R)Phe-Phe-Pab(Z) (0.42 mmol) was dissolved in 10 mL of tetrahydrofuran and 1 mL of water. Palladium on charcoal (42 mg) was charged to the solution and the mixture was hydrogenated at 45 psi hydrogen pressure in a Parr shaking apparatus for 2  
30 days. After complete hydrogenolysis the mixture was diluted with methanol and the catalyst was filtered off. Evaporation of the solvents gave crude H-(R)Phe-Phe-Pab which was purified by chromatography on neutral alumina (70-230 Mesh) eluting with  
35 methylenechloride-methanol--ammoniumhydroxide (80:20:2). Yield 76 mg of the title compound (41%). <sup>1</sup>H NMR (200 MHz, d<sub>6</sub>-DMSO); δ 7.61 (d, 2H), 7.4-7.0 (m,

12H), 4.64 (m, 1H), 4.44 (m, 2H), 4.13 (t, 1H), 3.1-2.8 (m, 4H).

### Example 86

5

**HOOC-CO-(R)Phe-Phe-Pab**

(i) MeOOC-CO-(R)Phe-Phe-Pab(Z)

10 H-(R)Phe-Phe-Pab(Z) (0.87 mmol) (see Example 85 (ii))  
was dissolved in 10 mL of tetrahydrofuran. The  
solution was cooled on an icewater bath and  
triethylamine (1.73 mmol) followed by  
methyloxalylchloride (0.95 mmol) were added. The  
15 cooling bath was removed and the reaction mixture  
stirred for 18 h at ambient temperature. The reaction  
mixture was diluted with ethylacetate and extracted  
with water. The organic phase was collected and the  
solvent was removed under reduced pressure yielding  
20 0.45 g of MeOOC-CO-(R)Phe-Phe-Pab(Z) (78%) which was  
used in the next step without further purification.  
TSP-MS found m/z 664 (calculated for  $MH^+$  ( $C_{37}H_{38}N_5O_7$ )  
664).

25 (ii) HOOC-CO-(R)Phe-Phe-Pab(Z)

MeOOC-CO-(R)Phe-Phe-Pab(Z) (0.68 mmol) was dissolved  
in 4 mL of tetrahydrofuran and 2 mL of water.  
Lithiumhydroxide (2.6 mmol) was added and the  
30 reaction mixture was stirred at room temperature for  
1.5 h. After complete hydrolysis the reaction mixture  
was diluted with 25 mL of water and acidified by  
addition of 0.5 mL of acetic acid. The precipitate  
was filtered and washed with several portions of  
35 water yielding 0.40 g of crude HOOC-CO-(R)Ph -Phe-  
Pab(Z) after drying under vacuum at 45°C for 24 h.  
The crude product was slurried in 10 mL of ethanol

and 1 mL of water. The solution was brought to reflux and the insoluble title compound was filtered off, yielding 0.23 g of HOOC-CO-(R)Phe-Phe-Pab(Z) (41% over two steps). <sup>1</sup>H NMR (200 MHz, d<sub>6</sub>-DMSO); δ 8.62 (m, 2H), 8.41 (d, 1H), 7.89 (d, 2H), 7.4-6.9 (m, 17H), 5.10 (s, 2H), 4.54 (m, 2H), 4.34 (m, 2H), 3.2-2.6 (m, 4H).

(iii) HOOC-CO-(R)Phe-Phe-Pab

10

HOOC-CO-(R)Phe-Phe-Pab(Z) (0.20 mmol) was slurried in 20 mL of tetrahydrofuran and 5 mL of water. Palladium on charcoal (52 mg) was charged to the solution and the mixture was hydrogenated at 45 psi hydrogen pressure in a Parr shaking apparatus for 2 days. After complete hydrogenolysis the mixture was diluted with 40 mL of methanol and the catalyst was filtered off. Evaporation of the solvents yielded 50 mg of the title compound (49%). <sup>1</sup>H NMR (200 MHz, d<sub>6</sub>-DMSO); δ 9.2(s), 8.78(d), 8.60(m), 7.91(m), 7.79(d, 2H), 7.35-6.8(m, 12H), 4.6-4.0(m, 4H), 3.0-2.6(m, 4H).

20

Example 87

25 HOOC-CH<sub>2</sub>-(R)Phe-Phe-Pab

(i) BnOOC-CH<sub>2</sub>-(R)Phe-Phe-Pab(Z)

H-(R)Phe-Phe-Pab(Z) (0.87 mmol) (see Example 85 (ii)) and potassium carbonate (2.6 mmol) were slurried in 10 mL of acetonitrile. Iodobenzylacetate (0.95 mmol) was added to the mixture and the solution was heated to 30°C and stirred at that temperature for 2 days. After complete alkylation the solvent was removed and the residue dissolved in 10 mL of ethylacetate. The solution was rapidly extracted with 10 mL of water and from the collected organic phase the title

30

35

compound precipitates.  $\text{BnOOC-CH}_2\text{-(R)Phe-Phe-Pab(Z)}$  was filtered off and dried under vacuum at  $45^\circ\text{C}$  for 24 h yielding 177 mg  $\text{BnOOC-CH}_2\text{-(R)Phe-Phe-Pab(Z)}$  (28%).  $^1\text{H}$  NMR (200 MHz,  $\text{CDCl}_3$ );  $\delta$  7.79(d, 2H), 7.5-7.1(m, 22H), 6.55(t, 1H), 5.21(s, 2H), 5.03(s, 2H), 4.64(m, 1H), 4.41(m 2H), 3.3-2.6(m, 7H).

(ii)  $\text{BnOOC-CH}_2\text{-(R)Phe-Phe-Pab(Z)}$

10  $\text{BnOOC-CH}_2\text{-(R)Phe-Phe-Pab(Z)}$  (0.32 mmol) was slurried in 30 mL of tetrahydrofuran and 3 mL of water. Palladium on charcoal (41 mg) was charged to the solution and the mixture was hydrogenated at 45 psi hydrogen pressure in a Parr shaking apparatus for 2  
15 days. After complete hydrogenolysis the mixture was diluted with 40 mL of water and the catalyst was filtered off. Evaporation of the solvents yielded 95 mg of the title compound (59%). TSP-MS found  $m/z$  502 (calculated for  $\text{MH}^+$  ( $\text{C}_{28}\text{H}_{32}\text{N}_5\text{O}_4$ ) 502).

20 Example 88

**H-(R)Cha-Pro-Mig**

25 (i)  $\text{Boc-(R)Cha-Pro-Mig(Z)}$

To a stirred mixture of 0.344 g (0.93 mmol)  $\text{Boc-(R)Cha-Pro-OH}$  (see preparation of starting materials), 0.245 g (0.93 mmol) of  $\text{H-Mig(Z)}$  (see  
30 preparation of starting materials) and 0.227 g (1.86 mmol) of DMAP in 10 mL  $\text{CH}_3\text{CN}$  was added 0.232 g (1.21 mmol) of EDC at  $-10^\circ\text{C}$ . The reaction mixture was allowed to reach roomtemperature and left for 5 days. The  $\text{CH}_3\text{CN}$  was evaporated and the residue was dissolved  
35 in EtOAc and washed with  $\text{H}_2\text{O}$ ,  $\text{NaHCO}_3$  (aq) and brine. The organic layer was dried with  $\text{Na}_2\text{SO}_4$  and evaporated. The crude product was purified by flash



chr matography using a  
gradient of EtOAc/MeOH, 95/5 to 90/10, as eluent to  
yield 0.340 g (60 %) of the title compound.

5 (ii) H-(R)Cha-Pro-Mig(Z)

0.34 g (0.55 mmol) Boc-(R)Cha-Pro-Mig(Z) was  
dissolved in 8 mL of EtOAc saturated with HCl(g) and  
stirred for 10 min. at roomtemperature. 10 mL of a  
10 saturated solution of KOH(aq) was added dropwise. The  
layers were separated and the aqueous phase was  
extracted with 3x8 mL EtOAc. The organic layers were  
combined, washed with brine, dried with Na<sub>2</sub>SO<sub>4</sub> and  
evaporated to yield 0.286 g (100 %) of the title  
15 compound.

(iii) H-(R)Cha-Pro-Mig

0.050 g (0.132 mmol) of H-(R)Cha-Pro-Mig(Z) was  
20 dissolved in 3 mL MeOH and hydrogenated over 10 %  
Pd/C at atmospheric pressure over night. The solution  
was filtered through celite and the solvent  
evaporated to yield 0.040 g (80 %) of the title  
compound.

25 <sup>1</sup>H-NMR (500 MHz, MeOD): δ 0.92-1.02 (m, 2H), 1.18-1.47  
(m, 6H), 1.66-1.73 (m, 4H), 1.85-2.04 (m, 4H), 2.17-  
2.22 (m, 1H), 2.95-2.98 (m, 1H), 3.12-3.16 (m,  
1H), 3.47-3.55 (m, 2H), 3.62-3.66 (m, 1H), 3.75-3.78  
30 (m, 1H), 3.85-3.89 (m, 1H), 4.05-4.12 (m, 3H), 4.34-  
4.37 (m, 1H).

Signals from a minor rotamer appear at: δ 3.4, 3.7,  
4.13-4.16, 4.3.

35

MS m/z 379 (M<sup>+</sup> + 1)

Example 89**H-(R)Cha-Pro-Dig**5     **(i) Boc-(R)Cha-Pro-Dig(Z)**

To a stirred mixture of 0.280 g (0.76 mmol) Boc-(R)Cha-Pro-OH (see preparation of starting materials), 0.210 g (0.76 mmol) of H-Dig(Z) (see  
10 preparation of starting materials) and 0.186 g (1.52 mmol) of DMAP in 8 mL CH<sub>3</sub>CN was added 0.189 g (0.99 mmol) of EDC at -10°C. The reaction mixture was allowed to reach roomtemperature and left for 4 days. The CH<sub>3</sub>CN was evaporated and the residue was dissolved  
15 in EtOAc and washed with H<sub>2</sub>O, NaHCO<sub>3</sub> (aq) and brine. The organic layer was dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated. The crude product was purified by flash chromatography using a gradient of EtOAc/MeOH, 95/5 to 90/10, as eluent to  
20 yield 0.210 g (44 %) of the title compound.

**(ii) H-(R)Cha-Pro-Dig(Z)**

0.210 g (0.33 mmol) Boc-(R)Cha-Pro-Dig(Z) was  
25 dissolved in 8 mL of EtOAc saturated with HCl(g) and stirred for 10 min. at roomtemperature. 8 mL of a saturated solution of KOH(aq) was added dropwise. The layers were separated and the aqueous phase was extracted with 3x8 mL EtOAc. The organic layers were  
30 combined, washed with brine, dried with Na<sub>2</sub>SO<sub>4</sub> and evaporated to yield 0.146 g (83 %) of the title compound.

**(iii) H-(R)Cha-Pro-Dig**

35     0.046 g (0.087 mmol) of H-(R)Cha-Pro-Dig(Z) was dissolved in 3 mL MeOH and hydrogenated over 10 %

Pd/C at atmospheric pressure over night. The solution was filtered through celite and the solvent evaporated to yield 0.040 g (100 %) of the title compound.

5

$^1\text{H-NMR}$  (500 MHz, MeOD):  $\delta$  0.90-1.04 (m, 2H), 1.10-1.47 (m, 6H), 1.66-1.74 (m, 4H), 1.78-2.05 (m, 4H), 2.13-2.21 (m, 1H), 2.74-2.83 (m, 1H), 2.94-2.99 (m, 1H), 3.15-3.29 (m, 1H), 3.44-3.57 (m, 2H), 3.65-3.87 (m, 3H), 4.07-4.25 (m, 3H), 4.35-4.39 (m, 2H).

10

Signals from a minor rotamer appear at:  $\delta$  4.29-4.32.

MS  $m/z$  393 ( $M^+ + 1$ )

15

#### Example 90

#### **H-(R)Cha-Aze-Dig**

20 (i) Boc-(R)Cha-Aze-Dig(Z)

The title compound was prepared from Boc-(R)Cha-Aze-OH and H-Dig(Z) (see preparation of starting material) according to the procedure for Boc-(R)Cha-Pro-Dig(Z) in a yield of 0.253 g (54 %).

25

(ii) H-(R)Cha-Aze-Dig(Z)

The title compound was prepared from Boc-(R)Cha-Aze-Dig(Z) according to the procedure for Boc-(R)Cha-Pro-Dig(Z) in a yield of 0.210 g (100 %).

30

(iii) H-(R)Cha-Aze-Dig

35 0.060 g (0.117 mmol) of H-(R)Cha-Aze-Dig(Z) was dissolved in 3 mL MeOH and hydrogenated over 10 % Pd/C at atmospheric pressure over night. The solution

was filtered through celite and the solvent evaporated to yield 0.042 g (95 %) of the title compound.

5 <sup>1</sup>H-NMR (500 MHz, MeOD):  $\delta$  0.91-1.02 (m, 2H), 1.18-1.48  
(m, 6H), 1.66-1.90 (m, 8H), 2.15-2.17 (m, 1H), 2.66-  
2.68 (m, 1H), 2.80-2.83 (m, 1H), 3.14-3.29 (m, 1H),  
3.39-3.44 (m, 1H), 3.72-3.80 (m, 2H), 4.01-4.04 (m,  
10 1H), 4.14-4.23 (m, 2H), 4.48-4.49 (m, 1H), 4.60-4.64  
(m, 1H).

Signals from a minor rotamer appear at:  $\delta$  2.25, 2.6,  
4.3, 4.67.

15 MS m/z 379 ( $M^+ + 1$ ).

#### Examples of pharmaceutical preparations

The compound according to the invention can be  
20 formulated in solid dosage forms for oral  
administration such as plain tablets, coated tablets  
or modified release tablets. Liquid or solid-  
semisolid dosage forms for rectal administration.  
Lyophilized substance or liquids as emulsion or  
25 suspension for parenteral use. Liquid solid or  
semisolid dosage forms for topical administration.

In pressurized aerosols or in dry powder inhalers for  
oral or nasal inhalation.

30

#### Example P1

##### Tablets for oral administration

35

1000 tablets are prepared from the following  
ingredients:

	Active compound	100 g
	Lact se	200 g
	Polyvinyl pyrrolidone	30 g
	Microcrystalline cellulose	30 g
5	Magnesium stearate	6 g

The active constituent and lactose are mixed with an aqueous solution of polyvinyl pyrrolidone. The mixture is dried and milled to form granules. The microcrystalline cellulose and then the magnesium stearate are then admixed. The mixture is then compressed in a tablet machine giving 1000 tablets, each containing 100 mg of active constituent.

15

Example P2**Solution for parenteral administration**

20 A solution is prepared from the following ingredients:

	Active compound	5 g
	Sodium chloride for injection	6 g

25

Sodium hydroxide for pH adjustment ad pH 5-7  
Water for inj. up to 1000 ml

30 The active constituent and the sodium chloride are dissolved in the water. The pH is adjusted with 2 M NaOH to pH 3-9 and the solution is filled into sterile ampoules.

Example P3

35

Tablets for oral administration

- |   |                               |       |
|---|-------------------------------|-------|
|   | 1. Active compound            | 150 g |
|   | 2. Sodium aluminium silicate  | 20 g  |
|   | 3. Paraffin                   | 120 g |
|   | 4. Microcrystalline cellulose | 20 g  |
| 5 | 5. Hydroxy propyl cellulose   | 5 g   |
|   | 6. Sodium stearyl fumarate    | 3 g   |

1-4 are mixed and an aqueous solution of 5 is added. The mixture is dried and milled and 6 is admixed. The mix is then compressed in a tablet machine.

#### Example B6

##### **Inhaler powder**

15

The active compound is micronized in a jet mill to a particle size suitable for inhalation (mass diameter < 4 $\mu$ m).

20

100 mg of the micronized powder is filled into a powder multidose inhaler (Turbohaler®). The inhaler is equipped with a dosing unit which delivers a dose of 1 mg.

25

##### Biology

##### Determination of Thrombin clotting Time (TT):

30

Human thrombin (T 6769, Sigma Chem Co) in buffer solution, pH 7.4, 100  $\mu$ l, and inhibitor solution, 100  $\mu$ l, were incubated for one min. Pooled normal citrated human plasma, 100  $\mu$ l, was then added and the clotting time measured in an automatic device (KC 10, Amelung).

35

The clotting time in seconds was plotted against the inhibitor concentration, and the IC<sub>50</sub>TT was determined

by interpolation.

IC<sub>50</sub>TT is the concentration of inhibitor that doubles the thrombin clotting time for human plasma.

5  
Determination of Activated Partial Thromboplastin Time (APTT)

10 APTT was determined in pooled normal human citrated plasma with the reagent PTT Automated 5 manufactured by Stago. The inhibitors were added to the plasma (10 µl inhibitor solution to 90 µl plasma) and APTT was determined in the mixture by use of the coagulation analyser KC10 (Amelung) according to the instructions  
15 of the reagent producer. The clotting time in seconds was plotted against the inhibitor concentration in plasma and the IC<sub>50</sub>APTT was determined by interpolation.

20 IC<sub>50</sub>APTT is defined as the concentration of inhibitor in plasma that doubled the Activated Partial Thromboplastin Time.

Determination of thrombin time ex vivo

25 The inhibition of thrombin after oral administration of the compounds were examined in conscious rats that two days prior to the experiment were equipped with a catheter for blood sampling from the carotid artery.  
30 On the experimental day blood samples were withdrawn at fixed times after the administration of the compound into plastic tubes containing 1 part sodium citrate solution (0.13 mol per L.) and 9 parts of blood. The tubes were centrifuged to obtain platelet  
35 poor plasma. The plasma was used for determination of thrombin time as described below.

The citrated rat plasma, 100  $\mu$ l, was diluted with a saline solution, 0.9%, 100  $\mu$ l, and plasma coagulation was started by the addition of human thrombin (T 6769, Sigma Chem Co, USA) in a buffer solution, pH 7.4, 100  $\mu$ l. The clotting time was measured in an automatic device (KC 10, Amelung, Germany).

Determination of the inhibition constant  $K_i$  for plasma kallikrein

$K_i$  determinations were made with a chromogenic substrate method, and performed on a Cobas Bio centrifugal analyzer manufactured by Roche (Basel, Switzerland). Residual enzyme activity after incubation of human plasma kallikrein with various concentrations of test compound was determined at three different substrate concentrations, and measured as change in optical absorbance at 405 nm and 37°C.

Human plasma kallikrein (E.C.3.4.21.34, Chromogenix AB, Mölndal, Sweden), 250  $\mu$ l of 0.4 nkat/ml in buffer (0.05 mol/l Tris-HCl, pH 7.4, 10.15 adjusted with NaCl) with bovine albumin 5 g/l (cat no 810033, ICI Biochemicals Ltd, High Wycombe, Bucks, GB), was incubated for 300 s with 80  $\mu$ l of test compound solution in 0.15 mol/l NaCl containing albumin 10 g/l. An additional 10  $\mu$ l of water was supplied in this step. Then 40  $\mu$ l of kallikrein substrate (S-2302, Chromogenix AB, 1.25, 2.0 or 4.0 mmol/l in water) was added together with another 20  $\mu$ l of water, and the absorbance change monitored.

$K_i$  was evaluated from Dixon plots, i.e. diagrams of inhibit r concentration versus  $1/(\Delta A/\text{min})$ , where the data for the different substrate concentrations form straight lines which intercept at  $x = -K_i$ .



## ABBREVIATIONS


	Ac =	acetyl
	aq =	aqueous
5	Aze =	Azetidine-2-carboxylic acid
	betaPic =	Piperidine-3-carboxylic acid
	Boc =	tert-butyloxycarbonyl
	Boc-Dig(Z) =	3-(N-tert-butyloxycarbonyl- aminoethyl)-1-(N-benzyloxy- carbonylamidino) azetidine
10	Boc-Mig(Z) =	3-(N-ter-butyloxycarbonyl- aminomethyl)-1-(N-benzyloxy- carbonylamidino) azetidine
	Boc-Pig(Z) =	4-(N-tert-butyloxycarbonyl- aminomethyl)-1-(N-benzyloxy- carbonylamidino) piperidine
15	Boc-Pig(Z) <sub>2</sub> =	4-(N-tert-butyloxycarbonyl- aminomethyl)-1-(N,N'-dibenzyloxy- carbonylamidino) piperidine
20	Brine =	saturated water/NaCl solution
	Bn =	benzyl
	Bu =	butyl
	Cgl =	Cyclohexyl glycine
	Cha =	$\beta$ -cyclohexyl alanine
25	CME-CDI =	1-Cyclohexyl-3-(2-morpholinoethyl) carbodiimide metho-p-toluenesulfonate
	DBU =	1,8-diazabicyclo[5.4.0]undec-7-ene
	DCC =	dicyclohexyl carbodiimide
	DCU =	dicyclohexyl urea
30	DMAP=	N,N-dimethyl amino pyridine
	DMF =	dimethyl formamide
	DMSO =	dimethyl sulphoxide
	EDC =	1-(3-Dimethylaminopropyl)-3- ethylcarbodiimide hydrochloride
35	Et =	ethyl
	EtOAc =	thyl ac tat
	EtOH =	ethan l

	Gly =	glycine
	h =	hours
	HCl =	hydrochloric acid
	Hex =	hexyl
5	HOAc =	acetic acid
	HOBT =	N-hydroxy benzotriazole
	Hoc =	Homocyclohexyl alanine
	Hop =	Homophenyl alanine
	HOSu =	N-hydroxysuccinimide
10	H-Dig(Z) =	3-aminoethyl-1-(N-benzyloxycarbonyl-amidino) azetidine
	H-Dig =	3-aminoethyl-1-amidino azetidine
	H-(R,S)Hig(Z)=	(3RS)-1-(N-benzyloxycarbonylamidino)-3-aminoethyl pyrrolidine
15	H-(R,S)Hig =	(3RS)-1-amidino-3-aminoethyl pyrrolidine
	H-Hig =	1-amidino-3-aminoethyl pyrrolidine
	H-(R,S)Itp(Ts)=	(4RS)-1,3-diaza-2-tosylimino-4-aminoethylcyclohexane
20	H-(R,S)Itp =	(4RS)-1,3-diaza-2-imino-4-aminoethylcyclohexane
	H-(S)Itp(Ts)=	(4S)-1,3-diaza-2-tosylimino-4-aminoethylcyclohexane
	H-(S)Itp =	(4S)-1,3-diaza-2-imino-4-aminoethylcyclohexane
25	H-Itp =	1,3-diaza-2-imino-4-aminoethyl cyclohexane
	H-Mig(Z) =	3-aminomethyl-1-(N-benzyloxycarbonylamidino) azetidine
30	H-Mig =	3-aminomethyl-1-amidino azetidine
	H-(R,S)Nig(Z)=	(3RS)-1-(N-benzyloxycarbonylamidino)-3-aminomethyl pyrrolidine
	H-(R,S)Nig =	(3RS)-1-amidino-3-aminomethyl pyrrolidine
35	H-Nig =	1-amidino-3-aminomethyl pyrrolidine
	H-Pab =	1-amidin -4-aminomethyl benzene



	H-Pab(Z) =	4-aminomethyl-1-(N-benzyloxy carbonylamidino) benzene
	H-Pac =	1-amidino-4-aminomethyl cyclohexane
5	H-Pac(Z) =	4-aminomethyl-1-(N-benzyloxy carbonylamidino) cyclohexane
	H-Pig =	4-aminomethyl-1-amidino piperidine
	H-Pig(Z) =	4-aminomethyl-1-(N-benzyloxycarbonyl- amidino)-piperidine
10	H-Pig(Z) <sub>2</sub> =	4-aminomethyl-1-(N,N'-dibenzyloxy carbonylamidino) piperidine
	H-Rig(Z) =	4-aminoethyl-1-(N-benzyloxy- carbonylamidino)piperidine
	H-Rig =	4-aminoethyl-1-N-amidino piperidine
	Me =	methyl
15	MeOH =	methanol
	Mpa =	mega pascal
	Ms =	mesyl
	NMM =	N-methyl morpholine
	Pd/C =	palladium on charcoal
20	Pgl =	phenyl glycine
	Phe =	phenyl alanine
	Pic =	pipecolinic acid
	Pro =	proline
25	RPLC =	Reverse phase high performace liquid chromathography
	Tf =	trifluoromethylsulfonyl
	TFA =	trifluoroacetic acid
	THF =	tetrahydrofuran
30	Tic =	1-carboxy-1,2,3,4-tetrahydro- isoquinoline
	Ts =	tosyl
	Val =	valine
	Z =	benzyloxy carbonyl

35 Prefixes n, s, i and t have their usual meanings: normal, is, sec and tertiary. The stereochemistry for the amino acids is by default (S) if not otherwise stated.


ABBREVIATIONS (continued, the wavy lines on the nitrogen atoms in the structural formulas below signify the bond position of the fragment.)





Pab (n=1)

Pig (n=1)  
Rig (n=2)

Pac (n=1)




Itp (n=2)

Nig (n=1)  
Hig (n=2)Mig (n=1)  
Dig (n=2)

## CLAIMS






1. A compound of the general formula



Formula I

wherein:

A<sup>1</sup> represents a structural fragment of Formula  
IIa, IIb, IIc, IId or IIe;



wherein:

k is an integer 0, 1, 2, 3 or 4;

m is an integer 1, 2, 3 or 4;

5 q is an integer 0, 1, 2, or 3;

R<sup>1</sup> represents H, an alkyl group having 1 to 4 carbon atoms, or R<sup>11</sup>OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group R<sup>17</sup>-(CH<sub>2</sub>)<sub>p</sub>-, wherein p is 0, 1 or 2 and R<sup>17</sup> is methyl, phenyl, OH, COOR<sup>12</sup>, CONHR<sup>12</sup>, where R<sup>12</sup> is H or an alkyl group having 1 to 4 carbon atoms, and R<sup>11</sup> is H or an alkyl group having 1 to 6 carbon atoms, or

R<sup>1</sup> represents Ph(4-COOR<sup>12</sup>)-CH<sub>2</sub>-, where R<sup>12</sup> is as defined above, or

20 R<sup>1</sup> represents R<sup>13</sup>-NH-CO-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where R<sup>13</sup> is H or an alkyl group having 1 to 4 carbon atoms or -CH<sub>2</sub>COOR<sup>12</sup>, where R<sup>12</sup> is as defined above, or

30 R<sup>1</sup> represents R<sup>12</sup>OOC-CH<sub>2</sub>-OOC-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where R<sup>12</sup> is as defined above, or

35 R<sup>1</sup> represents R<sup>14</sup>SO<sub>2</sub>-, Ph(4-COOR<sup>12</sup>)-SO<sub>2</sub>-, Ph(3-COOR<sup>12</sup>)-SO<sub>2</sub>-, Ph(2-COOR<sup>12</sup>)-SO<sub>2</sub>- where R<sup>12</sup> is as defined above and R<sup>14</sup> is an alkyl group having 1-4 carbon atoms, or

R<sup>1</sup> represents -CO-R<sup>15</sup>, where R<sup>15</sup> is an alkyl group

having 1-4 carbon atoms, r

$R^1$  represents  $-\text{CO}-\text{OR}^{15}$ , where  $R^{15}$  is as defined above,  
or

5

$R^1$  represent  $-\text{CO}-(\text{CH}_2)_p-\text{COOR}^{12}$ , where  $R^{12}$  is as defined  
above and p is an interger 0, 1 or 2, or

10

$R^1$  represents  $-\text{CH}_2\text{PO}(\text{OR}^{16})_2$ ,  $-\text{CH}_2\text{SO}_3\text{H}$  or  
 $-\text{CH}_2-(5-(1\text{H})-\text{tetrazolyl})$  where  $R^{16}$  is, individually at  
each occurrence, H, methyl or ethyl;

15

$R^2$  represents H or an alkyl group having 1 to 4 carbon  
atoms or  $R^{21}\text{OOC-alkyl-}$ , where the alkyl group has 1 to 4  
carbon atoms and where  $R^{21}$  is H or an alkyl group  
having 1 to 4 carbon atoms;

20

$R^3$  represents an alkyl group having 1-4 carbon atoms,  
and the alkyl group may or may not carry one or more  
fluorine atoms, or

25

$R^3$  represents a cyclopentyl, cyclohexyl- or a phenyl  
group which may or may not be substituted with an alkyl  
group having 1 to 4 carbon atoms, or

30

$R^3$  represents a phenyl group substituted with a  $\text{OR}^{31}$   
group, where  $R^{31}$  is H or an alkyl group having 1 to 4  
carbon atoms and k is 0, 1, or

$R^3$  represents a 1-naphthyl or 2-naphthyl group and k is  
0, 1, or

35

$R^3$  represent a cis- or trans-decalin group and k is 0,  
1, or


$R^3$  represents 4-pyridyl, 3-pyrrolidyl or 3-indolyl  
which may r may not b substitut d with a  $\text{OR}^{31}$  gr up,

where  $R^{31}$  is as defined above and  $k$  is 0, 1, or

$R^3$  represents  $\text{Si}(\text{Me})_3$  or  $\text{CH}(\text{R}^{32})_2$ , wherein  $\text{R}^{32}$  is a cyclohexyl- or a phenyl group;

$R^4$  represents H, an alkyl group having 1 to 4 carbon atoms, a cyclohexyl or a phenyl group;

$A^2$  represents a structural fragment of Formula IIIa, IIIb or IIIc



wherein:

$p$  is an integer 0, 1 or 2;

$m$  is an integer 1, 2, 3 or 4;

$Y$  represents a methylene group, or

$Y$  represents an ethylene group and the resulting 5-membered ring may or may not carry one or two fluorine atoms, a hydroxy group or an oxo group in position 4, or may or may not be unsaturated, or

$Y$  represents  $-\text{CH}_2-\text{O}-$ ,  $-\text{CH}_2-\text{S}-$ ,  $-\text{CH}_2-\text{SO}-$ , with the heteroatom functionality in position 4, or



Y represents a n-propylene group and the resulting 6-membered ring may or may not carry in position 5 one fluorine atom, a hydroxy group or an oxo group, carry two fluorine atoms in one of positions 4 or 5 or be unsaturated in position 4 and 5, or carry in position 4 an alkyl group with 1 to 4 carbon atoms, or

Y represents  $-\text{CH}_2-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{S}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{SO}-\text{CH}_2-$ , or

Y represent  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ;

$\text{R}^3$  is as defined above;

$\text{R}^5$  represents H or an alkyl group having 1 to 4 carbon atoms, or


$\text{R}^5$  represents  $-(\text{CH}_2)_p-\text{COOR}^{51}$ , where p is 0, 1 or 2 and  $\text{R}^{51}$  is H or an alkyl group having 1 to 4 carbon atoms;

n is an integer 0, 1, 2, 3 or 4;


B represents a structural fragment of Formula IVa, IVb, IVc or IVd

25


30




IVa



IVb



IVc



IVd

35

wh rein:

r is an interger 0 or 1;

5  $x^1$  represent  $\text{CH}_2$  or NH or is absent;

$x^2$  represents  $\text{CH}_2$ , NH or  $\text{C}=\text{NH}$ ;

10  $x^3$  represents NH,  $\text{C}=\text{NH}$ ,  $\text{N}-\text{C}(\text{NH})-\text{NH}_2$ ,  $\text{CH}-\text{C}(\text{NH})-\text{NH}_2$ ,  $\text{CH}-\text{NH}-\text{C}(\text{NH})-\text{NH}_2$  or  $\text{CH}-\text{CH}_2-\text{C}(\text{NH})-\text{NH}_2$ ;

$x^4$  represents  $\text{CH}_2$  or NH;

15  $x^5$  represents  $\text{C}(\text{NH})-\text{NH}_2$  or  $\text{NH}-\text{C}(\text{NH})-\text{NH}_2$ ;


$x^6$  represents CH or N;

$\text{R}^6$  is H or an alkyl group having 1-4 carbon atoms;

20 either the compound as such or stereoisomers thereof or in the form of a physiologically acceptable salt.

2. A compound of the general formula

25




Formula V

30

wherein:

35  $\text{A}^1$  represents a structural fragment of Formula IIa, IIb, IIc, II d r IIe;



wherein:

20

k is an integer 0, 1, 2, 3 or 4;

m is an integer 1, 2, 3 or 4;

25

q is an integer 0, 1, 2, or 3;

30

$R^1$  represents  $R^{11}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted in the position which is alpha to the carbonyl group, and the alpha substituent is a group  $R^{17}-(CH_2)_p$ -, wherein p is 0, 1 or 2 and  $R^{17}$  is,  $COOR^{12}$ ,  $CONHR^{12}$ , where  $R^{12}$  is H, an alkyl group having 1 to 4 carbon atoms or a benzyl group, and  $R^{11}$  is H or an alkyl group having 1 to 6 carbon atoms, or a benzyl group, or

35

$R^1$  represents  $Ph(4-COOR^{12})-CH_2$ -, where  $R^{12}$  is as defined above, or

5  $R^1$  represents  $R^{13}$ -NH-CO-alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{13}$  is H or an alkyl group having 1 to 4 carbon atoms or  $-CH_2COOR^{12}$ , where  $R^{12}$  is as defined above, or

10  $R^1$  represents  $R^{12}OOC-CH_2-OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and is possibly substituted alpha to the carbonyl with an alkyl group having 1 to 4 carbon atoms and where  $R^{12}$  is as defined above, or

15  $R^1$  represents  $R^{14}SO_2$ -,  $Ph(4-COOR^{12})-SO_2$ -,  $Ph(3-COOR^{12})-SO_2$ -,  $Ph(2-COOR^{12})-SO_2$ - where  $R^{12}$  is as defined above and  $R^{14}$  is an alkyl group having 1-4 carbon atoms, or

20  $R^1$  represents  $-CO-R^{15}$ , wherein  $R^{15}$  is an alkyl group having 1-4 carbon atoms, or

$R^1$  represents  $-CO-OR^{15}$ , where  $R^{15}$  is as defined above, or

25  $R^1$  represent  $-CO-(CH_2)_p-COOR^{12}$ , where  $R^{12}$  is as defined above and p is an interger 0, 1 or 2, or

30  $R^2$  represents H or an alkyl group having 1 to 4 carbon atoms or  $R^{21}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and where  $R^{21}$  is H or an alkyl group having 1 to 4 carbon atoms or a benzyl group;

$R^3$  represents an alkyl group having 1-4 carbon atoms, and the alkyl group may or may not carry one or more fluorine atoms, or

35

$R^3$  represents a cyclopentyl, cyclohexyl- or a phenyl group which may or may not be substituted with an alkyl

group having 1 to 4 carbon atoms,  $r$

$R^3$  represents a phenyl group substituted with a  $OR^{31}$  group, where  $R^{31}$  is H or an alkyl group having 1 to 4 carbon atoms and  $k$  is 0, 1, or

$R^3$  represents a 1-naphthyl or 2-naphthyl group and  $k$  is 0, 1, or


$R^3$  represent a cis- or trans-decalin group and  $k$  is 0, 1, or

$R^3$  represents 4-pyridyl, 3-pyrrolidyl or 3-indolyl which may or may not be substituted with a  $OR^{31}$  group, where  $R^{31}$  is as defined above and  $k$  is 0, 1, or


$R^3$  represents  $Si(Me)_3$  or  $CH(R^{32})_2$ , wherein  $R^{32}$  is a cyclohexyl- or a phenyl group;

$R^4$  represents H, an alkyl group having 1 to 4 carbon atoms, a cyclohexyl or a phenyl group;


$A^2$  represents a structural fragment of Formula IIIa, IIIb or IIIc



IIIa



IIIb



IIIc

wher in:

p is an interger 0, 1 or 2;

5 m is an integer 1, 2, 3 or 4;

Y represents a methylene group, or

10 Y represents an ethylene group and the resulting 5-membered ring may or may not carry one or two fluorine atoms, a hydroxy group or an oxo group in position 4, or may or may not be unsaturated, or

15 Y represents  $-\text{CH}_2-\text{O}-$ ,  $-\text{CH}_2-\text{S}-$ ,  $-\text{CH}_2-\text{SO}-$ , with the heteroatom functionality in position 4, or

20 Y represents a n-propylene group and the resulting 6-membered ring may or may not carry in position 5 one fluorine atom, a hydroxy group or an oxo group, carry two fluorine atoms in one of positions 4 or 5 or be unsaturated in position 4 and 5, or carry in position 4 an alkyl group with 1 to 4 carbon atoms, or

25 Y represents  $-\text{CH}_2-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{S}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{SO}-\text{CH}_2-$ , or

Y represent  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ;

$\text{R}^3$  is as defined above;

30  $\text{R}^5$  represents H or an alkyl group having 1 to 4 carbon atoms, or


$\text{R}^5$  represents  $-(\text{CH}_2)_p-\text{COOR}^{51}$ , where p is 0, 1 or 2 and  $\text{R}^{51}$  is H or an alkyl group having 1 to 4 carbon atoms;

35 n is an int ger 0, 1, 2, 3 r 4;


B r presents a structural fragment f Formula IVa, IVb, IVc or IVd

5


10




IVa



IVb



IVc



IVd

15

wherein:

r is an interger 0 or 1;

X<sup>1</sup> represent CH<sub>2</sub> or NH or is absent;

20

X<sup>2</sup> represents CH<sub>2</sub>, NH or C=NH;

X<sup>3</sup> represents NH, C=NH, N-C(NH)-NH<sub>2</sub>, CH-C(NH)-NH<sub>2</sub>, CH-NH-C(NH)-NH<sub>2</sub> or CH-CH<sub>2</sub>-C(NH)-NH<sub>2</sub>;

25

X<sup>4</sup> represents CH<sub>2</sub> or NH;

X<sup>5</sup> represents C(NH)-NH<sub>2</sub> or NH-C(NH)-NH<sub>2</sub>;

30

X<sup>6</sup> represents CH or N;

R<sup>6</sup> is H or an alkyl group having 1-4 carbon atoms;

D is Z or (Z)<sub>2</sub>;

35

Z is a benzyloxy carbonyl gr up;

either the compound as such or stereoisomers thereof or in the form of a physiologically acceptable salt.

5 3. A compound according to claims 1 or 2 wherein  $A^1$  is a structural fragment of formula IIa or IIb.

10 4. A compound according to one or more of the precedings claims 1-3 wherein  $R^1$  represents  $R^{11}OOC$ -alkyl-, where the alkyl group has 1 to 4 carbon atoms and  $R^{11}$  is H.

15 5. A compound according to one or more of the precedings claims 1-4 wherein  $A^2$  is a structural fragment of formula IIIa.

6. A compound according to one or more of the preceding claims 1-4 wherein  $A^2$  is a structural fragment of formula IIIb.

20 7. A compound according to one or more of the preceding claims 1-6 wherein B is a structural fragment of formula IVa, wherein  $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $CH-C(NH)-NH_2$ , r is 1 and n is 1.

25 8. A compound according to one or more of the preceding claims 1-6 wherein B is a structural fragment of formula IVa, wherein  $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$ ,  $X^3$  is  $N-C(NH)-NH_2$ , r is 0 or 1 and n is 1 or 2.

30 9. A compound according to one or more of the preceding claims 1-6 wherein B is a structural fragment of formula IVb, wherein  $X^5$  is  $C(NH)-NH_2$  and  $R^6$  is H and n is 1.

35 10. A compound according to one or more of the preceding claims 1-6 wherein B is a structural element of formula IVa, wherein  $X^1$  and  $X^3$  are NH,  $X^2$  is  $C=NH$ ,  $X^4$



is  $\text{CH}_2$ ,  $r$  is 1 and  $n$  is 2.

11. A compound according to one or more of the preceding claims 1-6 wherein B is a structural element of formula IVa, wherein  $X^1$  is absent,  $X^2$  and  $X^4$  are  $\text{CH}_2$ ,  $X^3$  is  $\text{N-C(NH)-NH}_2$ ,  $r$  is 0 and  $n$  is 1 or 2.

12. A compound according to claims 1 or 2 in which  $n$  is 1 or 2,  $A^1$  is a structural fragment of formula IIa wherein  $k$  is 0 or 1,  $R^1$  represents  $\text{R}^{11}\text{OOC-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms,  $R^2$  represents H,  $R^3$  represents a cyclohexyl group,  $A^2$  represents a structural fragment of Formula IIIa wherein Y represents a methylene group, an ethylene group, or a  $n$ -propylene group and the resulting 6-membered ring may or may not carry in position 4 an alkyl group with 1 to 4 carbon atoms,  $R^5$  represents H, B represents a structural fragment of formula IVa wherein  $X^1$ ,  $X^2$  and  $X^4$  are  $\text{CH}_2$  and  $X^3$  is  $\text{CH-C(NH)-NH}_2$  or  $\text{N-C(NH)-NH}_2$ ,  $r$  is 0 or 1, or  $X^1$  and  $X^3$  is NH,  $X^2$  is  $\text{C=NH}$ ,  $X^4$  is  $\text{CH}_2$ ,  $r$  is 1, or  $X^1$  is absent,  $X^2$  and  $X^4$  are  $\text{CH}_2$ ,  $X^3$  is  $\text{N-C(NH)-NH}_2$ ,  $r$  is 0.

13. A compound according to claims 1 or 2 in which  $n$  is 1,  $A^1$  is a structural fragment of formula IIa wherein  $k$  is 0 or 1,  $R^1$  represents  $\text{R}^{11}\text{OOC-alkyl-}$ , where the alkyl group has 1 to 4 carbon atoms,  $R^2$  represents H,  $R^3$  represents a cyclohexyl group,  $A^2$  represents a structural fragment of Formula IIIa wherein Y represents a methylene group, an ethylene group, or a  $n$ -propylene group and the resulting 6-membered ring may or may not carry in position 4 an alkyl group with 1 to 4 carbon atoms,  $R^5$  represents H, B represents a structural fragment of formula IVb wherein  $X^5$  represents  $\text{C(NH)-NH}_2$  and  $R^6$  is H.

## 14. A compound selected from

- HOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 5 HOOC-CH<sub>2</sub>-(R) Cgl-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Pro-Pab  
 (HOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Cgl-Pro-Pab  
 H-(R) Cgl-Pic-Pab  
 HOOC-CH<sub>2</sub>-(R, S) CH(COOH)-(R) Cgl-Pic-Pab  
 10 H-(R) Cha-Aze-Pab  
 HOOC-CH<sub>2</sub>-(R) Cha-Aze-Pab  
 HOOC-CH<sub>2</sub>-(R, S) CH(COOH)-(R) Cha-Aze-Pab  
 HOOC-CH<sub>2</sub>-(RorS) CH(COOH)-(R) Cha-Aze-Pab/a  
 HOOC-CH<sub>2</sub>-(RorS) CH(COOH)-(R) Cha-Aze-Pab/b  
 15 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Pab  
 HOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R) Cha-Aze-Pab  
 H-(R) Cha-Pro-Pab  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pab  
 HOOC-CH<sub>2</sub>-(Me) (R) Cha-Pro-Pab  
 20 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(Me) (R) Cha-Pro-Pab  
 HOOC-CH<sub>2</sub>-(RorS) CH(COOH)-(R) Cha-Pro-Pab/a  
 HOOC-CH<sub>2</sub>-(RorS) CH(COOH)-(R) Cha-Pro-Pab/b  
 HOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R) Cha-Pro-Pab  
 25 EtOOC-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pab  
 Ph(4-COOH)-SO<sub>2</sub>-(R) Cha-Pro-Pab  
 H-(R) Cha-Pic-Pab  
 HOOC-CH<sub>2</sub>-(R) Cha-Pic-Pab  
 HOOC-CH<sub>2</sub>-(RorS) CH(COOH)-(R) Cha-Pic-Pab/a  
 30 HOOC-CH<sub>2</sub>-(RorS) CH(COOH)-(R) Cha-Pic-Pab/b  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pic-Pab  
 HOOC-CO-(R) Cha-Pic-Pab  
 HOOC-CH<sub>2</sub>-CO-(R) Cha-Pic-Pab  
 Me-OOC-CH<sub>2</sub>-CO-(R) Cha-Pic-Pab  
 35 H<sub>2</sub>N-CO-CH<sub>2</sub>-(R) Cha-Pic-Pab  
 Boc-(R) Cha-Pic-Pab  
 Ac-(R) Cha-Pic-Pab

- Me-SO<sub>2</sub>-(R) Cha-Pic-Pab  
 H-(R) Cha-(R, S) betaPic-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-(R, S) betaPic-Pab  
 HOOC-CH<sub>2</sub>-(R) Cha-Val-Pab  
 5 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Val-Pab  
 H-(R) Hoc-Aze-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Hoc-Aze-Pab  
 HOOC-CH<sub>2</sub>-(R, S) CH(COOH)-(R) Hoc-Pro-Pab  
 HOOC-CH<sub>2</sub>-(R) Hoc-Pic-Pab  
 10 (HOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Hoc-Pic-Pab  
 HOOC-CH<sub>2</sub>-(R) Pro(3-(S)Ph)-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Pro(3-(S)Ph)-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Tic-Pro-Pab  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Aze-Pig  
 15 HOOC-CH<sub>2</sub>-(R) Cgl-Pro-Pig  
 H-(R) Cha-Aze-Pig  
 HOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pac  
 H-(R) Cha-Pro-Pac  
 H-(R) Cgl-Ile-Pab  
 20 H-(R) Cgl-Aze-Pab  
 HOOC-(R, S) CH(Me)-(R) Cha-Pro-Pab  
 MeOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 EtOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
<sup>n</sup>BuOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 25 <sup>n</sup>HexOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
 H-(R) Cgl-Pro-Pac  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pac  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Pro-Pac  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Pac  
 30 HOOC-CH<sub>2</sub>-(R) Cha-Aze-Pig  
 HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pig  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pig  
 (HOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Cgl-Pro-Pig  
 HOOC-CH<sub>2</sub>-CH<sub>2</sub>(HOOC-CH<sub>2</sub>)-(R) Cha-Pro-Pig  
 35 HOOC-CH<sub>2</sub>-(R) Cgl-Aze-(R, S) Itp  
 HOOC-CH<sub>2</sub>-(R) Cha-Aze-(R, S) Itp  
 H-(R) Cha-Pic-(R, S) Itp

- HOOC-CH<sub>2</sub>-(R) Cha-Pic-(R,S) Itp  
H-(R) Cgl-Pro-(R,S) Hig  
HOOC-CH<sub>2</sub>-(R) Cgl-Pro-(R,S) Hig  
H-(R) Cha-Pro-(R,S) Hig  
5 H-(R) Cgl-Aze-Rig  
HOOC-CH<sub>2</sub>-(R) Cgl-Aze-Rig  
HOOC-CH<sub>2</sub>-(R) Cha-Pro-Rig  
HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Rig  
HOOC-CH<sub>2</sub>-(R) Cha-Pro-(S) Itp  
10 H-(R) Cha-Pro-(R,S) Nig  
H-(R) Cha-Pro-Mig  
H-(R) Cha-Pro-Dig  
H-(R) Cha-Aze-Dig
- 15 either as such or stereoisomer thereof or in the form  
of a physiologically acceptable salt.
15. A compound selected from
- 20 HOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Pab  
HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pab  
HOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pab  
HOOC-CH<sub>2</sub>-(R) Cha-Pic-Pab  
25 HOOC-CH<sub>2</sub>-(R) Cgl-Pro-Pig  
EtOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pab  
HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pac  
HOOC-CH<sub>2</sub>-(R) Cha-Pro-Pig
- 30 either as such or stereoisomer thereof or in the form  
of a physiologically acceptable salt.
16. A compound selected from
- 35 BnOOC-CH<sub>2</sub>-(R) Cgl-Az -Pab(Z)  
BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Aze-Pab(Z)  
BnOOC-CH<sub>2</sub>-(R) Cgl-Pr -Pab(Z)

- BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Pr -Pab(Z)  
 (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Cgl-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R, S) CH(COOBn)-(R) Cgl-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Aze-Pab(Z)  
 5 BnOOC-CH<sub>2</sub>-(R, S) CH(COOBn)-(R) Cha-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(RorS) CH(COOBn)-(R) Cha-Aze-Pab(Z) /a  
 BnOOC-CH<sub>2</sub>-(RorS) CH(COOBn)-(R) Cha-Aze-Pab(Z) /b  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R) Cha-Aze-Pab(Z)  
 10 BnOOC-CH<sub>2</sub>-(R) Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(Me) (R) Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(Me) (R) Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R, S) CH(COOBn)-(R) Cha-Pro-Pab(Z)  
 15 BnOOC-CH<sub>2</sub>-NH-CO-CH<sub>2</sub>-(R) Cha-Pro-Pab(Z)  
 Ph(4-COOH)-SO<sub>2</sub>-(R) Cha-Pro-Pab(Z)  
 Boc-(R) Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R, S) CH(COOBn)-(R) Cha-Pic-Pab(Z)  
 20 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-Pic-Pab(Z)  
 EtOOC-CO-(R) Cha-Pic-Pab(Z)  
 MeOOC-CH<sub>2</sub>-CO-(R) Cha-Pic-Pab(Z)  
 H<sub>2</sub>N-CO-CH<sub>2</sub>-(R) Cha-Pic-Pab(Z)  
 Ac-(R) Cha-Pic-Pab(Z)  
 25 Me-SO<sub>2</sub>-(R) Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Cha-Val-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cha-(R, S) Val-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Hoc-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R, S) CH(COOBn)-(R) Hoc-Pro-Pab(Z)  
 30 BnOOC-CH<sub>2</sub>-(R) Hoc-Pic-Pab(Z)  
 (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R) Hoc-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R) Pro(3-(S) Ph)-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Pro(3-(S) Ph)-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Tic-Pro-Pab(Z)  
 35 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R) Cgl-Aze-Pig(Z)<sub>2</sub>  
 BnOOC-CH<sub>2</sub>-(R) Cgl-Pro-Pig(Z)<sub>2</sub>  
 BnOOC-CH<sub>2</sub>-(R) Cgl-Aze-Pac(Z)

- BnOOC-(R,S)CH(Me)-(R)Cha-Pro-Pab(Z)  
 MeOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
 EtOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
<sup>n</sup>BuOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
 5 <sup>n</sup>HexOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cgl-Pro-Pac(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Aze-Pac(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Aze-Pig(Z)  
 10 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z)  
 (BnOOC-CH<sub>2</sub>)<sub>2</sub>-(R)Cgl-Pro-Pig(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>(BnOOC-CH<sub>2</sub>)-(R)Cha-Pro-Pig(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pic-(R,S)Itp(Z)  
 15 BnOOC-CH<sub>2</sub>-(R)Cgl-Pro-(R,S)Hig(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Rig(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Rig(Z)  
 BnOOC-CH<sub>2</sub>-CH<sub>2</sub>-(R)Cha-Aze-Rig(Z)
- 20 either as such or stereoisomer thereof or in the form  
 of a physiologically acceptable salt.
17. A compound selected from
- 25 BnOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pic-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cgl-Pro-Pig(Z)<sub>2</sub>  
 30 EtOOC-CH<sub>2</sub>-(R)Cgl-Aze-Pab(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pac(Z)  
 BnOOC-CH<sub>2</sub>-(R)Cha-Pro-Pig(Z)
- either as such or stereoisomer thereof or in the form  
 35 of a physiologically acceptable salt.

18. A compound selected from

H-(R)Pro-Phe-Pab

HOOC-CH<sub>2</sub>-(R)Pro-Phe-Pab

5 H-(R)Phe-Phe-Pab

HOOC-CH<sub>2</sub>-(R)Phe-Phe-Pab

HOOC-CO-(R)Phe-Phe-Pab

10 either as such or stereoisomer thereof or in the form of a physiologically acceptable salt.

19. A compound selected from

Boc-(R)Pro-Phe-Pab(Z)

15 BnOOC-CH<sub>2</sub>-(R)Pro-Phe-Pab(Z)

Boc-(R)Phe-Phe-Pab(Z)

MeOOC-CO-(R)Phe-Phe-Pab(Z)

BnOOC-CH<sub>2</sub>-(R)Phe-Phe-Pab(Z)

20 either as such or stereoisomer thereof or in the forms of a physiologically acceptable salt.

20. A process for preparing a compound according to any of claims 1-19, which process comprises coupling of an  
25 N-terminally protected amino acid or dipeptide or amino acid, when an N-terminally amino acid is used a second amino acid is added afterwards using standard methods to a compound

30




wherein n is an integer 0, 1, 2, 3 or 4, X is B or B-D,  
35 where B is as defined in formula I and D is as defined in formula V as such or having the guanidino or amidino nitrogens either mono or diprotonated with an amine

protecting group such as a benzyl oxy carbonyl- or  
 tertbutoxy carbonyl- or p-toluenesulphonyl group or X  
 is a group transferable into B followed by removal of  
 the protecting group(s) or deprotection of the N-  
 5 terminal nitrogen followed by alkylation of the N-  
 terminal nitrogen and if desired deprotection by known  
 methods and if desired forming a physiologically  
 acceptable salt, and in those cases where the reaction  
 results in a mixture of stereoisomers, these are  
 10 optionally separated by standard chromatographic or  
 re-crystallisation techniques, and if desired a single  
 stereoisomer is isolated.

21. A process according to claim 20 for preparing a  
 15 compound according to any of claims 1-19, which  
 process comprises:

a) (Method Ia) Coupling of an N-terminally protected  
 dipeptide, selected from A<sup>1</sup> and A<sup>2</sup> in Formulas I or V by  
 20 using standard peptide coupling, shown in the formula




35 wherein n is as defined in Formula I, W<sup>1</sup> is an N-  
 terminal amino protecting group such as tert-butyl oxy  
 carbonyl and benzyloxy carbonyl and Q<sup>1</sup> is -C(NH)-





$\text{NH}_2$ ,  $-\text{C}(\text{NW}^2)-\text{NH}-\text{W}^2$ ,  $-\text{C}(\text{NH})-\text{NH}-\text{W}^2$ ,  $-\text{NH}-\text{C}(\text{NH})-\text{NH}_2$ ,  $-\text{NH}-\text{C}(\text{NH})-\text{NH}-\text{W}^2$ ,  $-\text{N}(\text{W}^2)-\text{C}(\text{NH})-\text{NH}-\text{W}^2$  or  $-\text{NH}-\text{C}(\text{NW}^2)-\text{NH}-\text{W}^2$ , where  $\text{W}^2$  is an amine protecting group such as tert-butyloxy carbonyl or benzyloxy carbonyl, or  $\text{Q}^1$  is  $-\text{CN}$ ,  $-\text{CO}-\text{NH}_2$  or  $-\text{CS}-\text{NH}_2$ , where the group is subsequently transferred into an amidino group or  $\text{Q}^1$  is  $\text{NH}_2$  or  $\text{NH}-\text{W}^2$ , where  $\text{W}^2$  is as defined above, where the amino group is subsequently transferred into a guanidino group (giving  $\text{Q}^1 = -\text{NH}-\text{C}(\text{NH})-\text{NH}_2$ ), after deprotection of the  $\text{W}^2$ -group when  $\text{Q}^1$  is  $-\text{NH}-\text{W}^2$  ( $\text{W}^2$  in this case must be orthogonal to  $\text{W}^1$ ), by methods known in the art.

b) (Method Ib) Coupling of an N-terminally protected amino acid, selected from  $\text{A}^2$  in Formulas I or V and prepared by using standard peptide coupling, shown in the formula




wherein  $n$ ,  $\text{W}^1$ , and  $\text{Q}^1$  are as defined above followed by deprotection of the  $\text{W}^1$ -group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method Ia above,

c) (Method IIa) Coupling of an N-terminally protected dipeptide, selected from A<sup>1</sup> and A<sup>2</sup> in Formulas I or V by using standard peptide coupling, shown in the formula



wherein n is as defined in Formula I, W<sup>1</sup> is an N-terminal amino protecting group such as tert. butyloxy carbonyl and benzyloxy carbonyl and Q<sup>1</sup> is -C(NH)-NH<sub>2</sub>, -C(NW<sup>2</sup>)-NH-W<sup>2</sup>, -C(NH)-NH-W<sup>2</sup>, -NH-C(NH)-NH<sub>2</sub>, -NH-C(NH)-NH-W<sup>2</sup>, -N(W<sup>2</sup>)-C(NH)-NH-W<sup>2</sup> or -NH-C(NW<sup>2</sup>)-NH-W<sup>2</sup>, where W<sup>2</sup> is an amine protecting group such as tert. butyloxy carbonyl or benzyloxy carbonyl, or Q<sup>1</sup> is -CN, -CO-NH<sub>2</sub> or CS-NH<sub>2</sub>, where the group is subsequently transferred into a amidino group or Q<sup>1</sup> is NH<sub>2</sub> or NH-W<sup>2</sup>, where W<sup>2</sup> is as defined above, where the amino group is subsequently transferred into a guanidino group (giving Q<sup>1</sup> = -NH-C(NH)-NH<sub>2</sub>), after deprotection of the W<sup>2</sup>-group when Q<sup>1</sup> is -NH-W<sup>2</sup> (W<sup>2</sup> in this case must be orthogonal to W<sup>1</sup>), by methods known in the art.


d) (Method IIb) Coupling of an N-terminally protected amino acid, selected from A<sup>2</sup> in Formulas I or V by using standard peptide coupling, shown in the formula



**15**


wherein n, W<sup>1</sup> and Q<sup>1</sup> are as defined above followed by deprotection of the W<sup>1</sup>-group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method IIa above,

e) (Metod IIIa) Coupling of an N-terminally protected dipeptide, selected from A<sup>1</sup> and A<sup>2</sup> in Formulas I or V by using standard peptide coupling, shown in the formula




wherein n is as defined in Formula I and r is 0 or 1 when  $X^1$ ,  $X^2$  and  $X^4$  are  $CH_2$  or r is 0 when  $X^2$  and  $X^4$  are  $CH_2$  and  $X^1$  is absent,  $W^1$  is an N-terminal amino protecting group such as tert-butyloxy carbonyl and benzyloxy carbonyl and  $Q^2$  is  $-C(NH)-NH_2$ ,  $-C(NW^2)-NH-W^2$ , or  $-C(NH)-NH-W^2$ , where  $W^2$  is an amine protecting group such as tert-butyloxy carbonyl or benzyloxy carbonyl, or  $Q^2$  is equal to  $W^2$  where the amino group, after deprotection of the  $W^2$  group ( $W^2$  in this case must be orthogonal to  $W^1$ ), is subsequently transferred into a guanidino group using a unprotected, N-protected or N,N'-diprotected guanidation reagent by methods known in the art,

f) (Method IIIb) Coupling of an N-terminally protected amino acid, selected from  $A^2$  in Formulas I or V by using standard peptide coupling, shown in the formula




wherein n, r,  $X^1$ ,  $X^2$  and  $X^4$ ,  $W^1$ , and  $Q^2$  are as defined above followed by deprotection of the  $W^1$ -group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method IIIa above,

g) (Method IVa) Coupling of an N-terminally protected dipeptide, selected from A<sup>1</sup> and A<sup>2</sup> in Formulas I or V by using standard peptide coupling, shown in the formula



20 wherein n is as defined in Formula I, W<sup>1</sup> is an N-terminal amino protecting group such as tert-butyloxy carbonyl or benzyloxy carbonyl and W<sup>3</sup> is H or an amino protecting group such as aryl sulfonyl, benzyloxy carbonyl or tert-butyloxy carbonyl.

25 h) (Method IVb) Coupling of an N-terminally protected amino acid, selected from A<sup>2</sup> in Formulas I or V by using standard peptide coupling, shown in the formula



wherein  $n$ ,  $W^1$ , and  $W^3$  are as defined above followed by deprotection of the  $W^1$ -group and coupling with the N-terminal amino acid, in a protected form, leading to the protected peptide described in Method IVa,

5

the final compounds can be made in any of the following ways, depending on the nature of the  $Q^1$ - or  $Q^2$ -groups used: Removal of the protecting group(s) (when  $Q^1 = -C(NH)-NH_2$ ,  $-C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH_2$ ,  $-NH-C(NH)-NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$ ), or a selective deprotection of the  $W^1$ - group (e.g when  $Q^1$  or  $Q^2 = -C(NW^2)-NH-W^2$ ,  $-C(NH)-NH-W^2$ ,  $-NH-C(NH)-NH-W^2$ ,  $-N(W^2)-C(NH)-NH-W^2$  or  $-NH-C(NW^2)-NH-W^2$  ( $W^2$  in this case must be orthogonal to  $W^1$ ) followed by alkylation of the N-terminal nitrogen and if desired deprotection.

10

15

22. A compound according to any of claims 1-19 for use in therapy.

20

23. A compound according to any of claims 1-5 or 7-17 for use as an anticoagulant or antithrombotic agent.

25

24. A compound according to any of claims 1-4, 6-10 or 18-19 for use as an antiinflammatory agent.

30

25. A pharmaceutical preparation comprising an effective amount of a compound as outlined in claims 1-19 in conjunction with one or more pharmaceutical carriers.

35

26. A pharmaceutical preparation comprising an effective amount of a compound as outlined in any of claims 1-5 or 7-17 in conjunction with one or more pharmaceutical carriers for use as an anticoagulant or antithrombotic agent.

27. A pharmaceutical preparation comprising an effective amount of a compound as outlined in any of claims 1-4, 6-10 or 18-19 in conjunction with one or more pharmaceutical carriers for use as an antiinflammatory agent.

28. Use of compound according to any of claims 1-5 or 7-17 as an active ingredient for manufacture of a pharmaceutical preparation for inhibition of thrombin in a human or animal organism.

29. Use of compound according to any of claims 1-4, 6-10 or 18-19 as an active ingredient for manufacture of a pharmaceutical preparation for inhibition of kininogenases in a human or animal organism.

30. A method for obtaining inhibition of thrombin in a human or animal organism in need of such inhibition, comprising administering to said organism an inhibitory effective amount of a compound claimed in any of claims 1-5 or 7-17.


31. A method for obtaining inhibition of kininogenases in a human or animal organism in need of such inhibition, comprising administering to said organism an inhibitory effective amount of a compound claimed in any of claims 1-4, 6-10 or 18-19.

32. Use of a compound of the formula:



either as such or having the amidino group either mono- or diprotected at the nitrogens with a protective group, or in the form of a salt, as a starting material in synthesis of a peptidic serine protease inhibitor, and in particular in synthesis of a peptidic thrombin inhibitor or a peptidic kininogenases inhibitor.


33. A structural fragment of the formula



as a structural element in a pharmaceutically active compound, especially a peptidic compound.

34. A compound 4-aminomethyl-1-(N-benzyloxycarbonylamidino) benzene either as such, in the form of a salt or having a protection with a benzyloxycarbonyl group at the other nitrogen.

35. Use of a compound of the formula:




either as such or having the amidino group either mono- or diprotected at the nitrogens with a protective group, or in the form of a salt, as a starting material in synthesis of a thrombin inhibitor, and in particular in synthesis of a peptidic thrombin inhibitor.



36. A structural fragment of the formula

5



as a structural element in a thrombin inhibitor,  
especially a peptidic thrombin inhibitor.


10

37. A compound 4-aminomethyl-1-(N-benzyloxy-  
carbonylamidino) cyclohexane either as such, in the  
form of a salt or having a protection with a  
benzyloxycarbonyl group at the other nitrogen.

15

38. A compound of the formula:

20




either as such or having the amidino group either mono-  
or diprotected at the nitrogens with a protective  
group, or in the form of a salt, .

25

39. Use of a compound as described in claim 38, as a  
starting material in the synthesis of a serine protease  
inhibitor.

30


40. A structural fragment of the formula



as a structural element in a thrombin inhibitor,  
especially a peptidic thrombin inhibitor.

41. A compound of the formula:

5



10

either as such or having the amidino group either mono-  
or diprotected at the nitrogens with a protective  
group, or in the form of a salt.


15

42. Use of a compound as described in claim 41, as a  
starting material in the synthesis of a serine protease  
inhibitor.

20

43. A structural fragment of the formula

25



30


as a structural element in a thrombin inhibitor,  
especially a peptidic thrombin inhibitor.

44. A compound 4-aminoethyl-1-benzyloxy-carbonylamidino  
piperidine either as such, in the form of a salt or  
having a protection with a benzyloxycarbonyl group at  
the nitrogen.

35

257

45. A compound of the formula:




where  $n$  is 1 or 2 and  $s$  is 0 or 1,  
either as such or having the amidino group either mono-  
or diprotected at the nitrogens with a protective  
15 group, or in the form of a salt.

46. Use of a compound as described in claim 45, as a  
starting material in the synthesis of a serine protease  
inhibitor.

20

47. A structural fragment of the formula



where  $n$  is 1 or 2 and  $s$  is 0 or 1,  
35 as a structural element in a thrombin inhibitor,  
especially a peptidic thrombin inhibitor.

48. A compound which is (3RS)-1-(N-benzyloxycarbonyl-amidino)-3-aminomethyl pyrrolidine, (3RS)-1-(N-benzyl-oxycarbonylamidino)-3-aminoethyl pyrrolidine, 3-amino-methyl-1-(N-benzyloxycarbonyl-amidino) azetidine,  
5 3-aminoethyl-1-(N-benzyloxycarbonyl-amidino) azetidine or either as such, in the form of a salt or having a protection with a benzyloxycarbonyl group at the other nitrogen.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/00535

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC 6 C07K 5/06, C07K 5/02, A61K 38/55, C07C 257/18, C07C 257/16, C07D 239/14, C07D 211/26, C07D 205/04, C07D 207/09 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: A61K, C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
BIOSIS, MEDLINE, EMBASE, CA, WPI, CLAIMS		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, A1, 0530167 (AKTIEBOLAGET ASTRA), 3 March 1993 (03.03.93) —	1-29
A	EP, A1, 0513543 (BEHRINGWERKE AKTIENGESELLSCHAFT), 19 November 1992 (19.11.92) —	1-29
A	EP, A1, 0236163 (SANOFI), 9 Sept 1987 (09.09.87) —	1-29
A	EP, A2, 0542525 (ELI LILLY AND COMPANY), 19 May 1993 (19.05.93) —	1-29
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "B" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
27 October 1994		31 -10- 1994
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer  Elisabeth Carlberg Telephone No. +46 8 782 25 00

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/00535

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	THROMBOSIS ET DIATHESIS HAEMORRHAGICA, Volume 23, No 3, 1970, j.D. Geratz, "Inhibition of Thrombin, Plasmin and Plasminogen Activation by Amidino Compounds" page 486 - page 499  --	32,34
X	THROMBOSIS ET DIATHESIS HAEMORRHAGICA, Volume 31, 1974, Erika Glusa et al, "The Influence of Benzamidine Derivatives on Human Platelet Function" page 172 - page 178  --	32,34
X	BIOCHEMICAL PHARMACOLOGY, Volume 23, 1974, Fritz Markwardt et al, "Synthetic Low Molecular Weight Inhibitors of Serum Kallikrein" page 2247 - page 2256  --	32,34
X	PHARMAZIE, Volume 34, No 10, 1979, D. Labes et al, "Hansch-Analyse der Hemmwirkung von 3- und 4-substituierten Benzamidinen gegenüber Thrombin, Plasmin und Trypsin" page 649 - page 653  --	132,34
X	EP, A1, 0001774 (BAYER AKTIENGESELLSCHAFT ZENTRALBEREICH PATENTE), 16 May 1979 (16.05.79), page 32, line 16 - page 33, line 14; page 51, line 1 - line 9  -----	35,37

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 94/00535

**Box I** Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 30 and 31  
because they relate to subject matter not required to be searched by this Authority, namely:  
See PCT Rule 39.1 (iv): Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.
2. ☒ Claims Nos.: 33, 36, 40, 43 and 47  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
  
\*\* see extra sheet
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II** Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

- a) claims 1-29, 33, 36, 40, 43 and 47,
- b) claims 32, 34, 35 and 37,
- c) claims 38 and 39,
- d) claims 41, 42, 44-46 and 48

\* see extra sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ A protest accompanied the payment of additional search fees.

- \* The subjects of invention as listed below are so different from each other, that no technical relationship can be appreciated to be present so as to form a single general inventive concept.

Claims 1-29, 33, 36, 40, 43 and 47 relate to compounds, to processes for preparing them, to pharmaceutical preparations containing the compounds, to the use of the compounds for manufacture of pharmaceutical preparations and to structural fragments, which have been regarded as being part of the compounds disclosed in claim 1.

Claims 32, 34, 35 and 37 relate to structurally similar compounds (intermediates) and to the use of such compounds as starting material in synthesis of enzyme inhibitors.

Claims 38 and 39 relate to a compound (intermediate) and to its use as starting material in the synthesis of a serine protease inhibitor.

Claims 41, 42, 44-46 and 48 relate to compounds (intermediates) and to their use as starting material in the synthesis of a serine protease inhibitor.

- \*\* The wording of claims 33, 36, 40, 43 and 47, which indicates several possible end compounds with a structural fragment, does not define one solution of one technical problem. Therefore, the claims are not searchable.



**INTERNATIONAL SEARCH REPORT**

Information on patent family members

01/10/94

International application No.

PCT/SE 94/00535

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A1- 0530167	03/03/93	AP-A- 313	10/02/94
		AU-A- 2499092	05/04/93
		CA-A- 2116527	18/03/93
		EP-A- 0605462	13/07/94
		FI-A,D- 940945	28/02/94
		HU-D- 9400589	00/00/00
		NZ-A- 243675	27/06/94
		WO-A- 9305069	18/03/93
EP-A1- 0513543	19/11/92	AU-B- 651196	14/07/94
		AU-A- 1608692	12/11/92
		CN-A- 1067249	23/12/92
		DE-A- 4115468	12/11/92
		JP-A- 5155898	22/06/93
		NZ-A- 242668	27/06/94
EP-A1- 0236163	09/09/87	FR-A,B- 2593812	07/08/87
		JP-A- 62228050	06/10/87
		US-A- 4977168	11/12/90
		EP-A,B- 0236164	09/09/87
		FR-A,B- 2593814	07/08/87
		US-A- 4791102	13/12/88
EP-A2- 0542525	19/05/93	AU-B- 646767	03/03/94
		AU-A- 2829892	13/05/93
		AU-A- 2831292	13/05/93
		CA-A- 2082748	13/05/93
		CN-A- 1074438	21/07/93
		FI-A- 925124	13/05/93
		HU-A- 65858	28/07/94
		JP-A- 5221964	31/08/93
		US-A- 5252566	12/10/93
EP-A1- 0001774	16/05/79	DE-A- 2748295	03/05/79
		JP-A- 54073702	13/06/79